


LECTURES ON q -ANALOGUES OF CARTAN DOMAINS AND ASSOCIATED HARISH-CHANDRA MODULES

L. Vaksman (Ed.)



Kharkov, Ukraine — 2001

To Vladimir Drinfeld

Contents

PREFACE	5
INTRODUCTION	
<i>S. Sinel'shchikov and L. Vaksman</i>	6
Part I THE SIMPLEST EXAMPLE	12
A NON-COMMUTATIVE ANALOGUE OF THE FUNCTION THEORY IN THE UNIT DISC	
<i>D. Shklyarov, S. Sinel'shchikov, and L. Vaksman</i>	13
QUANTUM DISC: THE BASIC STRUCTURES	
<i>D. Shklyarov, S. Sinel'shchikov, and L. Vaksman</i>	30
QUANTUM DISC: THE CLIFFORD ALGEBRA AND THE DIRAC OPERATOR	
<i>K. Schmüdgen, S. Sinel'shchikov, and L. Vaksman</i>	40
ON UNIQUENESS OF COVARIANT DEFORMATION WITH SEPARATION OF VARIABLES OF THE QUANTUM DISC	
<i>D. Shklyarov</i>	51
Part II UNITARY REPRESENTATIONS AND NON-COMMUTATIVE HARMONIC ANALYSIS	62
HARISH-CHANDRA EMBEDDING AND q-ANALOGUES OF BOUNDED SYMMETRIC DOMAINS	
<i>S. Sinel'shchikov and L. Vaksman</i>	63
q-ANALOGUES OF SOME BOUNDED SYMMETRIC DOMAINS	
<i>D. Shklyarov, S. Sinel'shchikov, and L. Vaksman</i>	67
ON A q-ANALOGUE OF THE FOCK INNER PRODUCT	
<i>D. Shklyarov</i>	73
GEOMETRIC REALIZATIONS FOR SOME SERIES OF REPRESENTATIONS OF THE QUANTUM GROUP $SU_{2,2}$	
<i>D. Shklyarov, S. Sinel'shchikov, and L. Vaksman</i>	94
Part III QUANTUM HARISH-CHANDRA MODULES ASSOCIATED TO q-CARTAN DOMAINS	111
NON-COMPACT QUANTUM GROUPS AND HARISH-CHANDRA MODULES	
<i>D. Shklyarov, S. Sinel'shchikov, A. Stolin, and L. Vaksman</i>	112
ON A q-ANALOGUES OF THE PENROSE TRANSFORM	
<i>D. Shklyarov, S. Sinel'shchikov, A. Stolin, and L. Vaksman</i>	117
SPHERICAL PRINCIPAL NON-DEGENERATE SERIES OF REPRESENTATIONS FOR THE QUANTUM GROUP $SU_{2,2}$	
<i>S. Sinel'shchikov, A. Stolin, and L. Vaksman</i>	125
Part IV ADDITIONAL RESULTS ON SOME QUANTUM VECTOR SPACES	135

**HIDDEN SYMMETRY OF THE DIFFERENTIAL CALCULUS ON THE
QUANTUM MATRIX SPACE**

S. Sinel'shchikov and L. Vaksman **136**

**q-ANALOGS OF CERTAIN PREHOMOGENEOUS VECTOR SPACES:
COMPARISON OF SEVERAL APPROACHES**

D. Shklyarov **141**

**HIDDEN SYMMETRY OF SOME ALGEBRAS OF q-DIFFERENTIAL
OPERATORS**

D. Shklyarov, S. Sinel'shchikov, and L. Vaksman **155**

PREFACE

This volume contains a mildly expanded version of lectures and talks at seminars and conferences, as well as review papers on subjects listed in the title of the volume. A great deal of these texts have already been published or sent to press. However, the only way to provide a good exposition of the field we are interested in is to collect all of those papers together.

Bounded symmetric domains form a favorite subject of research in function theory, non-commutative harmonic analysis, and representation theory of real reductive Lie groups. The authors introduce the notions of q -analogues of bounded symmetric domains and q -Harish-Chandra modules. For that, they follow the traditions of quantum group theory in replacing Lie groups with the quantum universal enveloping Drinfeld-Jimbo algebras and representations of groups with associated Harish-Chandra modules.

The purpose of this volume is to convince the reader in exceptional attraction of the deduced class of quantum homogeneous spaces and the related class of modules over quantum universal enveloping algebras.

The first part of the volume contains 4 lectures on the quantum unit disc. The first of them does not assume the knowledge of quantum group theory. It contains a list of problems which we find specific for this part of 'non-commutative function theory', together with explanations about what kind of results would be reasonable to expect when studying q -analogues of bounded symmetric domains.

The second part of the volume contains 3 lectures devoted to 'function theory' in q -analogues of bounded symmetric domains and the related problems of non-commutative harmonic analysis. A feature of these fields is that the algebras they deal with are involutive, and the modules over universal enveloping algebras involved are normally unitarisable.

The third part of the volume includes 3 lectures devoted to q -analogues of Harish-Chandra modules. These modules arise naturally as soon as one disregards involution and unitarisability. It is worthwhile to emphasize an important open problem in the quantum group theory, the problem of construction and classification of simple quantum Harish-Chandra modules discussed in the first lecture of this part.

The fourth part of the volume collects some additional and auxiliary results. In particular, the last lecture contains a discussion on equivalence of several approaches to construction of q -analogues for vector spaces which admit natural embeddings of the bounded symmetric domains in question.

These lectures were delivered within the period 1996 – 2001. In some of them it would be reasonable to provide references to papers which contain complete proofs of the results announced in the lectures. Also, in some other parts of the volume, we would like to mention some applications of our results. This information is attached as notes of the Editor to particular lectures.

Editor

INTRODUCTION

S. Sinel'shchikov

L. Vaksman

Institute for Low Temperature Physics & Engineering
47 Lenin Avenue, 61103 Kharkov, Ukraine

A great deal of a background of the quantum group theory was worked out by V. Drinfeld and expounded in his report at International Congress of Mathematicians [4].

Just after its appearance in 80-th the theory of quantum groups attracts an attention of a large number of specialists due to substantial new problems that arise here as well as due to perfectly surprising applications and links to various fields of mathematics and theoretical physics. By now, dozens of monographs and hundreds of research papers have been dedicated to the theory of quantum groups and related topics. It is too cumbersome to describe all the areas in quantum group theory. We are going to dwell here on the theory of 'functions' in q-Cartan domains.

Consider a bounded domain $\mathcal{D}$ in a finite dimensional complex vector space. It is said to be symmetric if every point $p \in \mathcal{D}$ is an isolated fixed point for a biholomorphic involutive automorphism

$$\varphi_p : \mathcal{D} \rightarrow \mathcal{D}, \quad \varphi_p \circ \varphi_p = \text{id}.$$

$\mathcal{D}$ is called reducible if it is biholomorphically isomorphic to a Cartesian product of two nontrivial domains. A classification of irreducible bounded symmetric domains up to isomorphism was obtained by Eli Cartan. There exist four series of such domains (A, C, D, BD) and the two exceptional domains. It was demonstrated by Harish-Chandra that every irreducible bounded symmetric domain admits a so called *standard* realization as the unit ball in a finite dimensional *normed* vector space. The irreducible bounded symmetric domains of this sort are called Cartan domains.

SERIES A. Let $m, n \in \mathbb{N}$ and $m \leq n$. Consider the vector space Mat_{mn} of complex matrices with m rows and n columns. Every such matrix $\mathbf{z}$ determines a linear map of Hermitian spaces $\mathbb{C}^n \rightarrow \mathbb{C}^m$. Equip Mat_{mn} with the operator norm $\|\mathbf{z}\|$. The unit ball $\mathcal{D} = \{\mathbf{z} \in \text{Mat}_{mn} \mid \|\mathbf{z}\| < 1\}$ is a Cartan domain of series A. It is easy to produce an automorphism with a fixed point $\mathbf{z}_0$ in the special case $\mathbf{z}_0 = 0$ (just $\mathbf{z} \mapsto -\mathbf{z}$). The general case reduces to this special case by considering an action of the group $\text{Aut}(\mathcal{D})$. This group admits an easy description if one passes from matrices $\mathbf{z}$ to graphs of the associated linear maps, thus using the embedding $\mathcal{D} \hookrightarrow \text{Gr}_m(\mathbb{C}^{m+n})$ (similar embeddings exist for all Cartan domains; these are called Borel embeddings).

EXAMPLE (SERIES C, D). The series C and D are defined in a similar way, except that $m = n$, and in the case of series C one should consider the symmetric matrices ($\mathbf{z} = \mathbf{z}^t$) while in the case of series D antisymmetric matrices ($\mathbf{z} = -\mathbf{z}^t$). For all those series the

This lecture has been delivered at the conference dedicated to the 40-th anniversary of the Institute for Low Temperature Physics and Engineering, August 2000, Kharkov

automorphisms of domains are just fractionally linear maps

$$\mathbf{z} \mapsto (\mathbf{az} + \mathbf{b})(\mathbf{cz} + \mathbf{d})^{-1}.$$

The following areas attract a permanent attention of specialists in function theory and theory of representations of real reductive Lie groups (see a list of references in [1]):

- construction of unitary representations of $\text{Aut}(\mathcal{D})$ in Hilbert spaces of functions in the domain $\mathcal{D}$ or on its boundary,
- the theory of Toeplitz and Hankel operators in those Hilbert spaces,
- non-commutative harmonic analysis,
- description of algebras of $\text{Aut}(\mathcal{D})$ -invariant differential operators and their common eigenfunctions,
- studying of generalized hypergeometric functions related to Cartan domains.

Our principal observation is that the methods of quantum group theory allows one to embed a great deal of the results on Cartan domains into a one-parameter family of 'q-analogues' (everywhere in the sequel $q \in (0, 1)$).

The term 'q-analogue' can be easily illustrated by an example of a q-analogue of the binomial formula:

$$(a + b)^k = \sum_{j=0}^k \binom{k}{j} b^j a^{k-j}.$$

Consider the free algebra $\mathbb{C}\langle a, b \rangle$ over the field of complex numbers with two generators a, b , together with the two-sided ideal generated by $ab - qba$. The associated factor algebra $\mathbb{C}[a, b]_q$ is called the algebra of functions on the quantum plane. The monomials $\{b^j a^l\}_{j, l \in \mathbb{Z}_+}$ form a basis in the vector space $\mathbb{C}[a, b]_q$. Thus we get well defined numbers $\begin{bmatrix} k; q \\ j \end{bmatrix}$ such that

$$(a + b)^k = \sum_{j=0}^k \begin{bmatrix} k; q \\ j \end{bmatrix} b^j a^{k-j}.$$

Of course, $\lim_{q \rightarrow 1} \begin{bmatrix} k; q \\ j \end{bmatrix} = \binom{k}{j}$. The numbers $\begin{bmatrix} k; q \\ j \end{bmatrix}$ are called q-binomial coefficients or Gaussian polynomials as they are really polynomials of the indeterminate q . They are well studied and are among the simplest q-special functions [5].

After introducing 'Planck's constant' h by $q = e^{-h/2}$, we observe a similarity between the passage from the relation $ab - ba = 0$ to $ab - e^{-h/2}ba = 0$ and the quantization procedure.

The quantum group theory [3] allows one to obtain non-commutative algebras which are q-analogues for algebras of functions in Cartan domains and q-analogues of some results of function theory in those domains. To rephrase, the quantum group theory leads to a variety of results of non-commutative function theory in q-Cartan domains.

Consider a Cartan domain $\mathcal{D} \subset V$. The algebra $\text{Pol}(V)_q$ of 'polynomial functions' on the quantum vector space V for $0 < q < 1$ is introduced as a *-algebra determined by its

generators $z_1, z_2, \dots, z_M$ and the relations

$$\begin{aligned} z_i z_j &= q^{k_{ij}} z_j z_i + \sum_{M \geq i > i' > j' > j \geq 1} a_{ii'jj'}(q) z_{j'} z_{i'}, \quad i > j \\ z_i^* z_j &= \sum_{i', j'=1}^M b_{ii'jj'}(q) z_{j'} z_{i'}^* + (1 - q^2) \delta_{ij}. \end{aligned}$$

The choice of k_{ij} , $a_{ii'jj'}(q)$, $b_{ii'jj'}(q)$ is determined by quantum symmetry arguments; in particular, $\lim_{q \rightarrow 1} a_{ii'jj'}(q) = 0$, $\lim_{q \rightarrow 1} b_{ii'jj'}(q) = \delta_{ii'} \delta_{jj'}$.

EXAMPLE. The simplest Cartan domain is just the unit disc in $\mathbb{C}$. In this special case the $*$ -algebra $\text{Pol}(\mathbb{C})_q$ is determined by a single generator z and a single relation:

$$z^* z = q^2 z z^* + 1 - q^2.$$

In the classical case (i.e. for $q = 1$) the $*$ -algebra $\text{Pol}(V)$ can be equipped with the norm

$$\|f\| = \max_{z \in \overline{\mathcal{D}}} |f(z)|,$$

where $\overline{\mathcal{D}}$ is the closure of the domain $\mathcal{D}$. Each element $z_0 \in V$ determines a $*$ -homomorphism

$$\text{Pol}(V) \rightarrow \mathbb{C}, \quad f(z) \mapsto f(z_0).$$

In this setting the points of $\overline{\mathcal{D}}$ are exactly the points which determine *bounded* linear functionals.

That is, one can use $\|f\|$ in $\text{Pol}(V)$ instead of $\|z\|$ in V to distinguish the domain $\mathcal{D} \subset V$. This approach can be transferred onto the quantum case. By definition,

$$\|f\| = \sup_{\tilde{T}} \|\tilde{T}(f)\|,$$

with $\tilde{T}$ is any irreducible $*$ -representation of $\text{Pol}(V)_q$ in a Hilbert space. (An experience shows that there exists a unique¹ faithful irreducible $*$ -representation T . This representation provides a supremum in the above equality. In the example $\text{Pol}(\mathbb{C})_q$ such $*$ -representation can be defined in $l^2(\mathbb{Z}_+)$ by $T(z)e_k = (1 - q^{(2k+1)})^{1/2} e_{k+1}$, $T(z^*)e_k = (1 - q^{2k})^{1/2} e_{k-1}$, $k \neq 0$, $T(z^*)e_0 = 0$.)

To conclude, we restrict ourselves to the case of unit disc in $\mathbb{C}$ in presenting a list of those formulae of classical function theory and classical harmonic analysis where we were lucky to 'insert the index q '. First, recall that

$$\begin{aligned} &\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad z \mapsto \frac{az + b}{cz + d}, \\ &\mathcal{D} = \{z \in \mathbb{C} \mid |z| < 1\}, \\ &\text{Aut}(\mathcal{D}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| \bar{a} = d, \bar{b} = c, ad - bc = 1 \right\} / \left\{ \pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\}, \\ &\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad z \mapsto \frac{az + b}{cz + d}. \end{aligned}$$

¹Up to unitary equivalence

The ring of $\text{Aut}(\mathcal{D})$ -invariant differential operators is generated by

$$\square = -(1 - |z|^2)^2 \frac{\partial}{\partial z} \cdot \frac{\partial}{\partial \bar{z}}$$

(the invariant Laplace operator). An invariant integral is of the form

$$d\nu = (1 - |z|^2)^{-2} \cdot \frac{d\text{Re}z d\text{Im}z}{\pi}$$

(the normalizing multiple is not essential here).

The spectrum of $\square$ in $L^2(d\nu)$ is purely continuous and fills the semiaxis $(-\infty, -\frac{1}{4}]$. Recall an explicit form of an eigenfunction expansion for $\square$.

The function $P(z, \zeta) = \frac{1 - |z|^2}{|1 - z\bar{\zeta}|^2}$ is called the Poisson kernel. If f is a 'good' function on the circle $\partial\mathcal{D} = \{z \in \mathbb{C} \mid |z| = 1\}$, then for all $\rho \in \mathbb{R}$ the function

$$u(z) = \int_{\partial\mathcal{D}} P(z, e^{i\theta})^{i\rho + \frac{1}{2}} f(e^{i\theta}) \frac{d\theta}{2\pi}$$

is a solution of the differential equation:

$$\square u = \left(-\frac{1}{4} - \rho^2\right) u.$$

An eigenfunction expansion of a 'good' function in the disc $\mathcal{D}$ has the form

$$u(z) = \int_0^\infty \left\{ \int_0^{2\pi} P(z, e^{i\theta})^{i\rho + \frac{1}{2}} \tilde{u}(\rho, e^{i\theta}) \frac{d\theta}{2\pi} \right\} s(\rho) d\rho,$$

with $\tilde{u}(\rho, e^{i\theta})$ being an analogue of the Fourier transform,

$$\tilde{u}(\rho, e^{i\theta}) = \int_{\mathcal{D}} P(z, e^{i\theta})^{-i\rho + \frac{1}{2}} u(z) d\nu(z),$$

and $s(\rho) d\rho = 2\rho \text{th}(\pi\rho) d\rho$ the Plancherel measure.

One more result is related to the invertibility of (the closure of) the operator $\square$ in $L^2(d\nu)$:

$$(\square^{-1}f)(z) = \int_{\mathcal{D}} G(z, \zeta) f(\zeta) d\nu(\zeta).$$

Here $G(z, \zeta) = \ln \left(\left| \frac{z - \zeta}{1 - z\bar{\zeta}} \right|^2 \right)$ is the Green function of $\square$.

For a 'good' function $u(z)$ in the disc $\mathcal{D}$ one has

$$u(z) = \frac{1}{2\pi i} \int_{\mathcal{D}} \frac{1}{(1 - \bar{\zeta}z)^2} u(\zeta) d\bar{\zeta} \wedge d\zeta + \frac{1}{2\pi i} \int_{\mathcal{D}} \frac{1 - |z|^2}{(z - \zeta)(1 - \bar{\zeta}z)} \cdot \frac{\partial u}{\partial \bar{\zeta}} d\bar{\zeta} \wedge d\zeta.$$

All the above results admit q -analogues. Here is the idea of some of the proofs. For $q = 1$ the radial part $\square^{(0)}$ of the operator $\square$ is a differential operator in a suitable Hilbert space of functions on the interval $(0, 1)$. For $0 < q < 1$ the spectrum of the operator $T(1 - zz^*)$ is a union of $\{0\}$ and the geometric progression $\{1, q^2, q^4, q^6, \dots\}$. This progression substitutes the interval $(0, 1)$, and the operator $\square^{(0)}$ is replaced by a difference operator (a difference approximation of the differential operator $\square^{(0)}$) in the space of functions on the above progression. What remains is to obtain formulae for eigenfunction expansion for this difference operator. A passage from the radial part $\square^{(0)}$ to the entire operator $\square$ is accessible via an application of quantum symmetry arguments.

Describe briefly a passage from the disc to generic Cartan domains and their q -analogues. Let $K \subset \text{Aut}(\mathcal{D})$ be a subgroup of all linear automorphisms of $\mathcal{D}$. The subalgebra of K -invariant elements of the algebra of continuous functions in $\overline{\mathcal{D}}$ admits a q -analogue. One can prove that this subalgebra is commutative, and its space of maximal ideals works as the set $\text{spec } T = \{0\} \cup \{1, q^2, q^4, q^6, \dots\}$. In this setting a single difference operator is replaced by a family of pairwise commuting self-adjoint difference operators in the associated Hilbert space. The problems in question are just the expansion problems in the common eigenfunctions.

References

- [1] J. Arazy. A survey of invariant Hilbert space of analytic functions on bounded symmetric domains, in: Contemporary Mathematics, **185**, 1995, p. 7 – 65.
- [2] R. J. Baston, M. G. Eastwood. The Penrose Transform. Its Interaction with Representation Theory, Clarendon Press, Oxford, 1989.
- [3] V. Chari, A. Pressley. A Guide to Quantum Groups, Cambridge Univ. Press, 1995.
- [4] V. Drinfeld. Quantum groups, in: Proceedings of the International Congress of Mathematicians, Berkeley, 1986, A. M. Gleason (ed), AMS, Providence RI, 1987, pp. 798 – 820.
- [5] G. Gasper, M. Rahman. Basic Hypergeometric Series, Cambridge Univ. Press, Cambridge, 1990.

Notes of the Editor

The recent works [1, 2] deals with the $*$ -representations of $\text{Pol}(V)_q$ in the special case of series A, that is $V = \text{Mat}_{m,n}$. The results of those works are applied in some of the lectures of this volume as well as in [3, 4]. The work [3] contains a description of all irreducible $*$ -representations of $\text{Pol}(\text{Mat}_{2,2})_q$ up to a unitary equivalence. In [4] one can find a proof of the fact that for $f \in \text{Pol}(\text{Mat}_{2,2})_q$ the norms $\|f\|$, $\|Tf\|$ are the same: $\|f\| = \|Tf\|$. These norms and the identity have been discussed in the introduction.

References

- [1] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035.
- [2] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum matrix ball: The Bergman kernel*, math.QA/9909036.

- [3] L. Turowska, *On Representations of a q -Analogue of the $*$ -Algebra $\text{Pol}(\text{Mat}_2)$* , J. Phys. **A34** (2001), 2063 – 2070, math.OA/0005243.
- [4] D. Proskurin, L. Turowska, *On the $*$ -algebra associated with $\text{Pol}(\text{Mat}_{2,2})_q$* , Methods of Functional Analysis and Topology, **7** (2001), No 1, 88 – 92.

PART I
THE SIMPLEST EXAMPLE

A NON-COMMUTATIVE ANALOGUE OF THE FUNCTION THEORY IN THE UNIT DISC

D. Shklyarov

S. Sinel'shchikov

L. Vaksman

Institute for Low Temperature Physics & Engineering

National Academy of Sciences of Ukraine

e-mail: sinelshchikov@ilt.kharkov.ua, vaksman@ilt.kharkov.ua

Abstract. The present work considers one of the simplest homogeneous spaces of the quantum group $SU(1,1)$, the q -analogue of the unit disc in $\mathbb{C}$. We present q -analogues of Cauchy-Green formulae, integral representations of eigenfunctions of the Laplace-Beltrami operator, Green functions for Poisson equation and an inversion formula for Fourier transform. It is also demonstrated that the two-parameter quantization of the disc introduced before by S. Klimek and A. Lesniewski, can be derived via the Berezin method.

1 Introduction

The theory of von Neumann algebras is a non-commutative analogue of the function theory of real variable and its far-reaching generalization. This approach is also applicable for developing a non-commutative analogue of the function theory of complex variable, as one can see in the classical work of W. Arveson [1]. However, a progress in non-commutative complex analysis was inhibited by the absence of substantial examples. The break was made possible by the fundamental works of L. Faddeev and his collaborators, V. Drinfeld, M. Jimbo, S. Woronovicz in quantum group theory. The subject of this paper is the simplest example of a quantum homogeneous complex manifold, the quantum disc. Meanwhile, a part of the exposed results are extensible onto the case of quantum bounded symmetric domains [16].

We obtain non-commutative analogues for some integral representations of functions in the disc and find a non-commutative analogue of the Plancherel measure. The Berezin method is applied to produce a formal deformation for our quantum disc. Our results provide an essential supplement to those of the well known work [13] in this field. The initial sections contain the necessary material on differential and integral calculi in the quantum disc.

In the cases when the statement in question to be proved via the quantum group theory is available in the mathematical literature (in particular, in our electronic preprints [17],[18],[19],[20]), we restrict ourselves to making exact references. This approach allows us to expound all the main results without applying the quantum group theory, and thus making this text intelligible for a broader class of readers.

The authors are grateful to V. Drinfeld for helpful discussions of a draft version of this work.

2 Functions in the quantum disc

We assume in the sequel all the vector spaces to be complex, and let q stand for a real number, $0 < q < 1$.

The works [15, 13, 9, 6] consider the involutive algebra $\text{Pol}(\mathbb{C})_q$ given by its generator z and the commutation relation

$$z^*z = q^2zz^* + 1 - q^2.$$

It arises both in studying the algebras of functions in the quantum disc [15, 13, 9] and in studying the q -analogues of the Weil algebra (oscillation algebra) [6].

Any element $f \in \text{Pol}(\mathbb{C})_q$ admits a unique decomposition

$$f = \sum_{jk} a_{jk}(f) z^j z^{*k}, \quad a_{jk} \in \mathbb{C}.$$

Moreover, at the limit $q \rightarrow 1$ we have $z^*z = zz^*$. This allows one to treat $\text{Pol}(\mathbb{C})_q$ as a polynomial algebra on the quantum plane.

It is worthwhile to note that the passage to the generators $a^+ = (1 - q^2)^{-1/2}z^*$, $a = (1 - q^2)^{-1/2}z$ realizes an isomorphism between the $*$ -algebra $\text{Pol}(\mathbb{C})_q$ and the q -analogue of the Weil algebra considered in [6]: $a^+a - q^2aa^+ = 1$.

Impose the notation $y = 1 - zz^*$. It is straightforward that

$$z^*y = q^2yz^*, \quad zy = q^{-2}yz. \quad (2.1)$$

It is also easy to show that any element $f \in \text{Pol}(\mathbb{C})_q$ admits a unique decomposition

$$f = \sum_{m>0} z^m \psi_m(y) + \psi_0(y) + \sum_{m>0} \psi_{-m}(y) z^{*m}. \quad (2.2)$$

The passage to the decomposition (2.2) is similar to the passage from Cartesian coordinates on the plane $\mathbb{C} \simeq \mathbb{R}^2$ to polar coordinates.

We follow [13, 15] in completing the vector space $\text{Pol}(\mathbb{C})_q$ to obtain the function space in the quantum disc.

Consider the $\text{Pol}(\mathbb{C})_q$ -module H determined by its generator v_0 and the relation $z^*v_0 = 0$. Let $T : \text{Pol}(\mathbb{C})_q \rightarrow \text{End}_{\mathbb{C}}(H)$ be the representation of $\text{Pol}(\mathbb{C})_q$ in H , and $l_{m,n}(f)$, $m, n \in \mathbb{Z}_+$, stand for the matrix elements of the operator $T(f)$ in the basis $\{z^m v_0\}_{m \in \mathbb{Z}_+}$.

Impose three topologies in the vector space $\text{Pol}(\mathbb{C})_q$ and prove their equivalence.

Let $\mathcal{T}_1$ be the weakest among the topologies in which all the linear functionals $l'_{j,k} : f \mapsto a_{jk}(f)$, $j, k \in \mathbb{Z}_+$, are continuous,

$\mathcal{T}_2$ the weakest among the topologies in which all the linear functionals $l''_{m,n} : f \mapsto \psi_m(q^{2n})$, $m \in \mathbb{Z}$, $n \in \mathbb{Z}_+$, are continuous,

$\mathcal{T}_3$ the weakest among the topologies in which all the linear functionals $l_{m,n}$, $m, n \in \mathbb{Z}_+$, are continuous.

Proposition 2.1 *The topologies $\mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_3$ are equivalent.*

Proof. Remind the standard notation:

$$(t; q)_m = \prod_{j=0}^{m-1} (1 - tq^j); \quad (t; q)_\infty = \prod_{j=0}^{\infty} (1 - tq^j).$$

It follows from the definitions that with $m \geq k$

$$z^{*k} z^m v_0 = (q^{2m}; q^{-2})_k \cdot z^{m-k} v_0.$$

Hence

$$l_{m,n}(f) = \sum_{j=0}^{\min(m,n)} (q^{2m}; q^{-2})_{m-j} \cdot l'_{n-j, m-j}; \quad m, n \in \mathbb{Z}_+.$$

Thus the topology $\mathcal{T}_1$ is stronger than $\mathcal{T}_3$. In fact, these topologies are equivalent since the linear span of the functionals $\{l_{m-j, n-j}\}_{j=0}^{\min(m,n)}$ coincides with that of $\{l'_{m-j, n-j}\}_{j=0}^{\min(m,n)}$. The equivalence of $\mathcal{T}_2$ and $\mathcal{T}_3$ follows from the relations $l_{j,k} = (q^{2j}; q^{-2})_{\max(0, j-k)} l'''_{k-j, \min(j,k)}$. $\square$

We denote the completion of the Hausdorff topological vector space $\text{Pol}(\mathbb{C})_q$ by $D(U)'_q$ and call it the space of distributions in the quantum disc. $D(U)'_q$ may be identified with the space of formal series of the form (2.2) whose coefficients $\psi_j(y)$ are defined in $q^{2\mathbb{Z}_+}$. The linear functionals $l_{m,n}$ are extendable by continuity onto the topological vector space $D(U)'_q$. Associate to each distribution $f \in D(U)'_q$ the infinite matrix $T(f) = (l_{m,n}(f))_{m,n \in \mathbb{Z}_+}$.

A distribution $f \in D(U)'_q$ is said to be finite if $\#\{(j,k) \mid \psi_j(q^{2k}) \neq 0\} < \infty$. Evidently, a distribution f is finite iff the matrix $T(f)$ has only finitely many non-zero entries. The vector space of finite functions in the quantum disc is denoted by $D(U)_q$. There exists a non-degenerate pairing

$$D(U)'_q \times D(U)_q \rightarrow \mathbb{C}; \quad f_1 \times f_2 \mapsto \text{tr } T(f_1)T(f_2).$$

The extension by continuity procedure allows one to equip $D(U)_q$ with a structure of $*$ -algebra, and $D(U)'_q$ with a structure of $D(U)_q$ -bimodule.

Consider the algebra $\text{Pol}(\mathbb{C})_q^{op}$ derived from $\text{Pol}(\mathbb{C})_q$ via a replacement of the multiplication law by the opposite one. The elements $z \otimes 1$, $z^* \otimes 1$, $1 \otimes z$, $1 \otimes z^*$ of $\text{Pol}(\mathbb{C})_q^{op} \otimes \text{Pol}(\mathbb{C})_q$ are denoted respectively by z, z^*, ζ, ζ^* . To avoid confusion in the notation, we use braces to denote the multiplication in $\text{Pol}(\mathbb{C})_q^{op} \otimes \text{Pol}(\mathbb{C})_q$, e.g. $\{zz^*\} = q^2\{z^*z\} + 1 - q^2$. The module H^{op} over $\text{Pol}(\mathbb{C})_q^{op}$ is defined by its generator v_0^{op} and the relation $zv_0^{op} = 0$. Apply the above argument to $\text{Pol}(\mathbb{C})_q^{op} \otimes \text{Pol}(\mathbb{C})_q$ -module $H^{op} \otimes H$ in order to introduce the algebra $D(U \times U)_q$ of finite functions in the Cartesian product of quantum discs, together with $D(U \times U)_q$ -bimodule $D(U \times U)'_q$.

The reason for replacement the multiplication law in $\text{Pol}(\mathbb{C})_q$ by the opposite one is cleared in [17, 22, 23].

The linear functional

$$\mu(f) = (1 - q^2) \sum_{m \in \mathbb{Z}_+} \psi_0(q^{2m}) q^{2m}, \quad f \in D(U)_q,$$

is called the (normalized) Lebesgue integral in the quantum disc, since under the formal passage to the limit as $q \rightarrow 1$ one has $\mu(f) \rightarrow \frac{1}{\pi} \int_U f d\text{Im } z \cdot d\text{Re } z$.

Let $K \in D(U \times U)'_q$; the integral operator $f \mapsto \text{id} \otimes \mu(K(1 \otimes f))$ with kernel K maps $D(U)_q$ into $D(U)'_q$. We are interested in solving an inverse problem which is in finding out the explicit formulae for kernels $K \in D(U \times U)'_q$ of well known linear operators. In

this field, an analogue of the Bergman kernel for the quantum disc was obtained in the work of S. Klimek and A. Lesniewski [13]:

$$K_q(z, \zeta) = (1 - z\zeta^*)^{-1}(1 - q^2 z\zeta^*)^{-1}.$$

Finally, equip H with the structure of a pre-Hilbert space by setting

$$(z^j v_0, z^m v_0) = \delta_{jm}(q^2; q^2)_m; \quad j, m \in \mathbb{Z}_+.$$

It is easy to show that $T(z^*) = T(z)^*$, $I - T(z)T(z^*) \geq 0$. Thus we get a $*$ -representation of $\text{Pol}(\mathbb{C})_q$ in the completion $\overline{H}$ of H (see [13]).

3 Differential forms and $\overline{\partial}$ -problem

Let $\Omega_q(\mathbb{C})$ stand for the involutive algebra determined by its generators z, dz and the relations

$$\begin{aligned} 1 - z^* \cdot z &= q^2(1 - z \cdot z^*), & dz \cdot z^* &= q^{-2}z^* \cdot dz, & dz \cdot z &= q^2z \cdot dz, \\ dz \cdot dz^* &= -q^{-2}dz^* \cdot dz, & dz \cdot dz &= 0. \end{aligned}$$

(Also, an application of the involution $*$ to the above yields

$$dz^* \cdot dz^* = 0, \quad dz^* \cdot z = q^2z \cdot dz^*, \quad dz^* \cdot z^* = q^{-2}z^* \cdot dz^*.)$$

Equip $\Omega_q(\mathbb{C})$ with the grading as follows:

$$\deg z = \deg z^* = 0, \quad \deg dz = \deg dz^* = 1.$$

There exists a unique linear map $d : \Omega_q(\mathbb{C}) \rightarrow \Omega_q(\mathbb{C})$ such that

$$d : z \mapsto dz, \quad d : z^* \mapsto dz^*, \quad d : dz \mapsto 0, \quad d : dz^* \mapsto 0,$$

and

$$d(\omega' \cdot \omega'') = d\omega' \cdot \omega'' + (-1)^{\deg \omega'} \cdot \omega' d\omega''; \quad \omega', \omega'' \in \Omega_q(\mathbb{C}).$$

Evidently, $d^2 = 0$, and $(d\omega)^* = d\omega^*$ for all $\omega \in \Omega_q(\mathbb{C})$.

Turn to a construction of operators $\partial, \overline{\partial}$. For that, we need a bigrading in $\Omega_q(\mathbb{C})$:

$$\deg z = \deg z^* = (0, 0); \quad \deg(dz) = (1, 0); \quad \deg(dz^*) = (0, 1).$$

Now d has a degree 1 and admits a unique decomposition into a sum $d = \partial + \overline{\partial}$ of operators $\partial, \overline{\partial}$ with bidegrees respectively $(1, 0)$ and $(0, 1)$. A standard argument allows one to deduce from $d^2 = 0$ that $\partial^2 = \overline{\partial}^2 = \partial\overline{\partial} + \overline{\partial}\partial = 0$. It is also easy to show that $(\partial\omega)^* = \overline{\partial}\omega^*$ for all $\omega \in \Omega_q(\mathbb{C})$.

Each element $\omega \in \Omega_q(\mathbb{C})$ is uniquely decomposable into a sum

$$\omega = f_{00} + dz f_{10} + f_{01} dz^* + dz f_{11} dz^*, \quad f_{ij} \in \text{Pol}(\mathbb{C})_q, \quad i, j = 0, 1.$$

Equip $\Omega_q(\mathbb{C})$ with a topology corresponding to this decomposition:

$$\Omega_q(\mathbb{C}) \simeq \text{Pol}(\mathbb{C})_q \oplus \text{Pol}(\mathbb{C})_q \oplus \text{Pol}(\mathbb{C})_q \oplus \text{Pol}(\mathbb{C})_q.$$

Pass as above via a completion procedure from $\text{Pol}(\mathbb{C})_q$ to the space of distributions $D(U)'_q$ and then to the space of finite functions to obtain the bigraded algebra $\Omega(U)_q$.

The operators $d, \partial, \bar{\partial}$ are transferred by continuity from $\Omega_q(\mathbb{C})$ onto the algebra $\Omega(U)_q$ of differential forms with finite coefficients in the quantum disc.

The subsequent constructions involve essentially q -analogues of type $(0,*)$ differential forms with coefficients in sections of holomorphic bundles. The latter carry a structure of bimodules over algebras of type $(0,*)$ differential forms as above. Remind the notion of a differentiation for such a bimodule.

Let Ω be a $\mathbb{Z}_+$ -graded algebra and M a $\mathbb{Z}_+$ -graded Ω -bimodule. A degree 1 operator is said to be a differentiation if for all $m \in M$, $\omega \in \Omega$ one has $\bar{\partial}(m\omega) = (\bar{\partial}m)\omega + (-1)^{\deg m}m \cdot \bar{\partial}\omega$, $\bar{\partial}(\omega m) = (\bar{\partial}\omega) \cdot m + (-1)^{\deg \omega}\omega \cdot \bar{\partial}m$.

Let $\lambda \in \mathbb{C}$. Consider the graded bimodule over $\Omega(\mathbb{C})_q^{(0,*)} = \Omega(\mathbb{C})_q^{(0,0)} + \Omega(\mathbb{C})_q^{(0,1)}$ determined by its generator v_λ with $\deg(v_\lambda) = 0$ and the relations

$$z \cdot v_\lambda = q^{-\lambda}v_\lambda \cdot z, \quad z^* \cdot v_\lambda = q^\lambda v_\lambda \cdot z^*, \quad dz^* \cdot v_\lambda = q^\lambda v_\lambda \cdot dz^*.$$

Denote this graded bimodule by $\Omega(\mathbb{C})_{\lambda,q}^{(0,*)}$. It possesses a unique degree 1 differentiation $\bar{\partial}$ such that $\bar{\partial}v_\lambda = 0$. Pass (via an extension by continuity) from polynomial coefficients to finite ones to obtain the graded bimodule $\Omega(U)_{\lambda,q}^{(0,*)}$ over $\Omega(U)_q^{(0,*)}$, together with its differentiation $\bar{\partial}$.

We restrict ourselves to the case $\lambda \in \mathbb{R}$ and equip the spaces $\Omega(U)_{\lambda,q}^{(0,0)}$, $\Omega(U)_{\lambda,q}^{(0,1)}$ with the scalar products

$$(f_1 \cdot v_\lambda, f_2 \cdot v_\lambda) = \int_{U_q} f_2^* f_1 (1 - zz^*)^{\lambda-2} d\mu, \quad (3.1)$$

$$(f_1 v_\lambda dz^*, f_2 v_\lambda dz^*) = \int_{U_q} f_2^* f_1 (1 - zz^*)^\lambda d\mu. \quad (3.2)$$

The completions of the pre-Hilbert spaces $\Omega(U)_{\lambda,q}^{(0,0)}$ and $\Omega(U)_{\lambda,q}^{(0,1)}$ can be used in the formulation of $\bar{\partial}$ -problem. Specifically, we mean finding a solution of the equation $\bar{\partial}u = f$ in the orthogonal complement to the kernel of $\bar{\partial}$. In the classical case ($q = 1$) such a formulation is standard [2], and the solution is very well known. If λ stand for a real number and $\lambda > 1$ than one has

$$u(z) = \frac{1}{2\pi i} \int_U \frac{1}{z - \zeta} \left(\frac{1 - |\zeta|^2}{1 - \bar{\zeta}z} \right)^{\lambda-1} f(\zeta) d\bar{\zeta} \wedge d\zeta. \quad (3.3)$$

(3.3) implies the "Cauchy-Green formula":

$$\begin{aligned} u(z) = \frac{\lambda - 1}{2\pi i} \int_U \frac{(1 - |\zeta|^2)^{\lambda-2}}{(1 - \bar{\zeta}z)^\lambda} u(\zeta) d\bar{\zeta} \wedge d\zeta + \\ + \frac{1}{2\pi i} \int_U \frac{(1 - |\zeta|^2)^{\lambda-1}}{(z - \zeta)(1 - \bar{\zeta}z)^{\lambda-1}} \frac{\partial u}{\partial \bar{\zeta}} d\bar{\zeta} \wedge d\zeta. \end{aligned} \quad (3.4)$$

Our purpose is to obtain the q -analogues of (3.3), (3.4) for $\lambda = 2$.

The standard way of solving the $\bar{\partial}$ -problem is to solve first the Poisson equation $\square\omega = f$ with $\square = -\bar{\partial}^*\bar{\partial}$. In the case $\lambda = 2$ the kernel in (3.3) is derived by a differentiation in z

of the Green function

$$G(z, \zeta) = \frac{1}{\pi} \ln(|z - \zeta|^2) - \frac{1}{\pi} \ln(|1 - z\bar{\zeta}|^2). \quad (3.5)$$

In its turn, (3.5) can be obtained by the d'Alembert method: the first term is contributed by a real source, and the second one is coming from an imaginary source.

Note that the differential calculus for the quantum disc we use here is well known (see, for example, [24]). Its generalization onto the case of an arbitrary bounded symmetric domain was obtained in [16] via an application of a quantum analogue of the Harish-Chandra embedding [11].

4 Green function for Poisson equation

With $q = 1$ the measure $d\nu = (1 - |z|^2)^{-2} d\mu$ is invariant with respect to the Möbius transformations. In the case $q \in (0, 1)$ impose an "invariant integral" $\nu : D(U)_q \rightarrow \mathbb{C}$, $f \mapsto \int_{U_q} f d\nu$ by setting

$$\int_{U_q} f d\nu \stackrel{\text{def}}{=} \int_{U_q} f \cdot (1 - z\bar{z}^*)^{-2} d\mu.$$

The Hilbert spaces $L^2(d\nu)_q$, $L^2(d\mu)_q$ are defined as completions of the vector spaces $D(U)_q = \Omega(U)_q^{(0,0)}$, $\Omega(U)_q^{(0,1)}$ with respect to the norms

$$\|f\| = \left(\int_{U_q} f^* f d\nu \right)^{1/2}, \quad \|f dz^*\| = \left(\int_{U_q} f^* f d\mu \right)^{1/2}.$$

Proofs of the following statements are to be found in [17, Proposition 5.7, Corollary 5.8], [19, Corollary 4.2].

Lemma 4.1 *There exist $0 < c_1 \leq c_2$ such that*

$$c_1 \leq \bar{\partial}^* \bar{\partial} \leq c_2. \quad (4.1)$$

Proposition 4.2 *The exact estimates for $\bar{\partial}^* \bar{\partial}$ are of the form*

$$\frac{1}{(1+q)^2} \leq \bar{\partial}^* \bar{\partial} \leq \frac{1}{(1-q)^2}. \quad (4.2)$$

The inequalities (4.1) allow one to extend by continuity the operators $\bar{\partial}$, $\square = -\bar{\partial}^* \bar{\partial}$ from the dense subspace of finite functions $D(U)_q$ onto the entire $L^2(d\nu)_q$. They also imply that for any $f \in L^2(d\nu)_q$ there exists a unique solution u of Poisson equation $\square u = f$. Now it follows from (4.2) that $\|u\| \leq (1+q)^2 \|f\|$.

One obtains easily from the definitions

Lemma 4.3 *The series*

$$\sum_{i,j \in \mathbb{Z}_+} z^{*i} \zeta^i \psi_{ij}(y, \eta) z \zeta^{*j}, \quad y = 1 - z^* z, \quad \eta = 1 - \zeta \zeta^*$$

converges in $D(U \times U)_q'$ for any family $\{\psi_{ij}(y, \eta)\}_{i,j \in \mathbb{Z}_+}$ of functions defined on $q^{2\mathbb{Z}_+} \times q^{2\mathbb{Z}_+}$.

Corollary 4.4 *For all $m \geq 0$ there exists a well defined generalized kernel*

$$G_m = \{((1 - \zeta\zeta^*)(1 - z^*\zeta)^{-1})^m ((1 - z^*z)(1 - z\zeta^*)^{-1})^m\}. \quad (4.3)$$

To state the principal result of the section, we need an expansion of the Green function (3.5):

$$\begin{aligned} \ln \frac{|z - \zeta|^2}{|1 - z\bar{\zeta}|^2} &= \ln \left(1 - \frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - z\bar{\zeta}|^2} \right) = \\ &= - \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - z\bar{\zeta}|^2} \right)^m. \end{aligned}$$

Evidently, a formal passage to a limit yields

$$\lim_{q \rightarrow 1} G_m = \left(\frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - z\bar{\zeta}|^2} \right)^m.$$

A proof of the following result one can find in [19, Theorem 1.2].

Theorem 4.5 *The continuous operator $\square^{-1}$ in $L^2(d\nu)_q$ coincides on the dense linear subspace $D(U)_q \subset L^2(d\nu)_q$ with the integral operator whose kernel is $G = - \sum_{m=1}^{\infty} \frac{q^{-2}-1}{q^{-2m}-1} G_m$:*

$$\square^{-1}f = \int_{U_q} G(z, \zeta) f(\zeta) d\nu.$$

Here $G_m \in D(U \times U)'_q$ is given by (4.3).

Note in conclusion that the operators $\partial, \bar{\partial}, \square$ admit an extension by continuity onto the space $D(U)'_q$ of distributions in the quantum disc.

5 The Cauchy-Green formula

One can use the differentials $\partial : \Omega_q^{(0,0)} \rightarrow \Omega_q^{(1,0)}, \bar{\partial} : \Omega_q^{(0,0)} \rightarrow \Omega_q^{(0,1)}$ to define the partial derivatives $\frac{\partial^{(l)}}{\partial z}, \frac{\partial^{(r)}}{\partial z}, \frac{\partial^{(l)}}{\partial z^*}, \frac{\partial^{(r)}}{\partial z^*}$. Specifically, we set up $\partial f = dz \cdot \frac{\partial^{(l)}f}{\partial z} = \frac{\partial^{(r)}f}{\partial z} dz, \bar{\partial} f = dz^* \cdot \frac{\partial^{(l)}f}{\partial z^*} = \frac{\partial^{(r)}f}{\partial z^*} dz^*$. It is easy to show that these operators admit extensions by continuity from $D(U)_q$ onto $D(U)'_q$.

Let $f \in D(U)_q$. Define the integral of the (1,1)-form $dz \cdot f \cdot dz^*$ over the quantum disc by $\int_{U_q} dz \cdot f \cdot dz^* = -2i\pi \int_{U_q} f d\mu$.

A proof of the following proposition can be found in [19, Theorem 2.1].

Proposition 5.1 *Let $f \in D(U)_q$. Then*

1. *There exists a unique solution $u \in L^2(d\mu)_q$ of the $\bar{\partial}$ -problem $\bar{\partial}u = f$, which is orthogonal to the kernel of $\bar{\partial}$.*
2. *$u = \frac{1}{2\pi i} \int_{U_q} d\zeta \frac{\partial^{(l)}}{\partial z} G(z, \zeta) f d\zeta^*$, with $G \in D(U \times U)'_q$ being the Green function of the Poisson equation.*
3. *$f = -\frac{1}{2\pi i} \int_{U_q} (1 - z\zeta^*)^{-1} (1 - q^{-2}z\zeta^*)^{-1} d\zeta f(\zeta) d\zeta^* - \frac{1}{2\pi i} \int_{U_q} d\zeta \frac{\partial^{(l)}}{\partial z} G(z, \zeta) \cdot \frac{\partial^{(r)}f}{\partial \zeta^*} d\zeta^*$.*

6 Eigenfunctions of the operator $\square$

Let $\mathbb{C}[\partial U]_q$ stand for the algebra of finite sums of the form

$$\sum_{m \in \mathbb{Z}} a_m e^{im\theta}, \quad \theta \in \mathbb{R}/2\pi\mathbb{Z} \quad (6.1)$$

with complex coefficients. The $\mathbb{C}[\partial U]_q$ -module of formal series of the form (6.1) is denoted by $\mathbb{C}[[\partial U]]_q$. We also denote the algebra of finite sums like (6.1) with coefficients from $D(U)_q$ by $D(U \times \partial U)_q$, and the module of formal series (6.1) with coefficients from $D(U)'_q$ by $D(U \times \partial U)'_q$. This vector space will be equipped by the topology of coefficientwise convergence.

The use of the index q in the notation for the above vector spaces is justified by the fact that, as one can show, they are in fact modules over the quantum universal enveloping algebra.

Recall the notations [10]:

$$(a; q^2)_\infty = \prod_{j \in \mathbb{Z}_+} (1 - aq^{2j}), \quad (a; q^2)_\gamma = \frac{(a; q^2)_\infty}{(aq^{2\gamma}; q^2)_\infty}, \quad \gamma \in \mathbb{C}.$$

With $q = 1$, the integral

$$u(z) = \int_{\partial U} \left(\frac{1 - |z|^2}{(1 - z\bar{\zeta})(1 - \bar{z}\zeta)} \right)^{l+1} f(\zeta) d\nu, \quad d\nu = \frac{d\theta}{2\pi},$$

represents an eigenfunction of $\square$ (see [12]):

$$\square u = \lambda(l)u, \quad \lambda(l) = \left(l + \frac{1}{2} \right)^2 - \frac{1}{4}.$$

With $q \in (0, 1)$, the power P^γ of the Poisson kernel $P = \frac{1 - |z|^2}{|1 - z\bar{\zeta}|^2}$ is replaced by the element $P_\gamma \in D(U \times \partial U)_q$:

$$P_\gamma = (1 - zz^*)^\gamma (z\zeta^*; q^2)_{-\gamma} \cdot (q^2 z^* \zeta; q^2)_{-\gamma}. \quad (6.2)$$

Here $(z\zeta^*; q^2)_{-\gamma}$, $(q^2 z^* \zeta; q^2)_{-\gamma}$ are the q -analogues of the powers $(1 - z\bar{\zeta})^{-\gamma}$, $(1 - \bar{z}\zeta)^{-\gamma}$, and the q -binomial theorem (see [10]) implies

$$\begin{aligned} (z\zeta^*; q^2)_{-\gamma} &= \sum_{n \in \mathbb{Z}_+} \frac{(q^{2\gamma}; q^2)_n}{(q^2; q^2)_n} (q^{-2\gamma} z\zeta^*)^n, \\ (q^2 z^* \zeta; q^2)_{-\gamma} &= \sum_{n \in \mathbb{Z}_+} \frac{(q^{2\gamma}; q^2)_n}{(q^2; q^2)_n} (q^{2-2\gamma} z^* \zeta)^n. \end{aligned}$$

The following proposition is proved in [19, Theorem 3.1].

Proposition 6.1 *For all $f \in \mathbb{C}[\partial U]_q$ the element*

$$u = \int_{\partial U} P_{l+1}(z, e^{i\theta}) f(e^{i\theta}) \frac{d\theta}{2\pi} \quad (6.3)$$

of $D(U)'_q$ is an eigenvector of $\square$:

$$\square u = \lambda(l)u, \quad \lambda(l) = -\frac{(1 - q^{-2l})(1 - q^{2l+2})}{(1 - q^2)^2}.$$

We need the following standard notation ([10]):

$$\begin{aligned} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; q; z \\ b_1, \dots, b_s \end{matrix} \right] &= \\ &= \sum_{n \in \mathbb{Z}_+} \frac{(a_1; q)_n \cdot (a_2; q)_n \cdot \dots \cdot (a_r; q)_n}{(b_1; q)_n \cdot (b_2; q)_n \cdot \dots \cdot (b_s; q)_n (q; q)_n} \left((-1)^n \cdot q^{\frac{n(n-1)}{2}} \right)^{1+s-r} \cdot z^n. \end{aligned}$$

Corollary 6.2 (cf. [22]). *The series*

$$\varphi_l(y) = {}_3\Phi_2 \left[\begin{matrix} y^{-1}, q^{-2l}, q^{2(l+1)}; q^2; q^2 \\ q^2; 0 \end{matrix} \right]$$

converges in $D(U)'_q$, and its sum is an eigenfunction of $\square$: $\square \varphi_l = \lambda(l) \varphi_l$.

Proof. The convergence of the series is due to the fact that it breaks for each $y \in q^{2\mathbb{Z}_+}$. So it suffices to establish the relation

$$\varphi_l(y) = \int_{\partial U} P_{l+1}(z, \zeta) d\nu.$$

It follows from the definitions that the above integral equals to

$$\begin{aligned} \sum_{n \in \mathbb{Z}_+} \frac{(q^{2l+2}; q^2)_n^2}{(q^2; q^2)_n^2} q^{-2(2l+1)n} y^{l+1} z^n z^{*n} &= \\ &= y^{l+1} {}_3\Phi_1 \left[\begin{matrix} q^{2+2l}, q^{2+2l}, y^{-1}; q^2; q^{-2(2l+1)}y \\ q^2 \end{matrix} \right]. \end{aligned}$$

Now it remains to apply the identity (see [10]):

$$b^n {}_3\Phi_1 \left[\begin{matrix} q^{-n}, b, \frac{q}{z}; q, \frac{z}{c} \\ \frac{bq^{1-n}}{c} \end{matrix} \right] = {}_3\Phi_2 \left[\begin{matrix} q^{-n}, b, \frac{bzq^{-n}}{c}; q, q \\ \frac{bq^{1-n}}{c}, 0 \end{matrix} \right],$$

with q being replaced by q^2 , y by q^{2n} , b by q^{2l+2} , z by q^{-2l} , and c by $q^{2+2l-2n}$. $\square$

Note that $\varphi_l(y)$ is a q -analogue of a spherical function on a hyperbolic plane (see [12]).

For each $l \in \mathbb{C}$ a linear operator has been constructed from $\mathbb{C}[\partial U]_q$ into the eigenspace of $\square$, associated to the eigenvalue $\lambda(l)$. Now we try to invert this linear operator.

For that, we need a q -analogue of the operator $b_r : f(z) \mapsto f(re^{i\theta})$ which restricts the function in the disc onto the circle $|z| = r$ of radius $r \in (0, 1)$. Let $r > 0$, $1 - r^2 \in q^{2\mathbb{Z}_+}$. Define a linear operator $b_r : D(U)'_q \rightarrow \mathbb{C}[[\partial U]]_q$ by

$$\begin{aligned} b_r : \sum_{j>0} z^j \cdot \psi_j(y) + \psi_0(y) + \sum_{j>0} \psi_{-j}(y) \cdot z^{*j} &\mapsto \\ &\sum_{j>0} (re^{i\theta})^j \cdot \psi_j(1 - r^2) + \psi_0(1 - r^2) + \sum_{j>0} \psi_{-j}(q^{-2j} \cdot (1 - r^2))(re^{-i\theta})^j. \end{aligned}$$

(It is implicit that the functions $\psi_j(y)$, $j \in \mathbb{Z}$, vanish at $y \notin q^{2\mathbb{Z}_+}$).

Recall the definition of the q -gamma-function ([10]):

$$\Gamma_q(x) = \frac{(q; q)_\infty}{(q^x; q)_\infty} (1 - q)^{1-x}.$$

One may assume without loss of generality that

$$0 \leq \operatorname{Im} l < \frac{\pi}{2 \ln(q^{-1})}, \quad \operatorname{Re} l \geq -\frac{1}{2}.$$

Proposition 6.3 *Let $\operatorname{Re} l > -\frac{1}{2}$, and $u \in D(U)_q'$ is an eigenfunction of $\square$ given by (6.3). Then in $\mathbb{C}[\partial U]_q$ one has*

$$f = \frac{\Gamma_{q^2}^2(l+1)}{\Gamma_{q^2}(2l+1)} \lim_{1-r^2 \in q^{2\mathbb{Z}_+}, r \rightarrow 1} (1-r^2)^l b_r u.$$

The proof of this proposition is based on the following result which was communicated to the authors by L. I. Korogodsky:

Lemma 6.4

$$1). \quad \lim_{x \in q^{-2\mathbb{Z}_+}, x \rightarrow \infty} \varphi_l \left(\frac{1}{x} \right) \bigg/ \left(\frac{\Gamma_{q^2}(2l+1)}{\Gamma_{q^2}^2(l+1)} x^l \right) = 1$$

if $\operatorname{Re} l > -\frac{1}{2}$.

$$2). \quad \lim_{x \in q^{-2\mathbb{Z}_+}, x \rightarrow \infty} \varphi_l \left(\frac{1}{x} \right) \bigg/ \left(\frac{\Gamma_{q^2}(-2l-1)}{\Gamma_{q^2}^2(-l)} x^{-l-1} \right) = 1$$

if $\operatorname{Re} l < -\frac{1}{2}$.

Proof. It follows from the relation $\varphi_l(y) = \varphi_{-1-l}(y)$ that one may restrict oneself to the case $\operatorname{Re} l > -\frac{1}{2}$. An application of the identity ([10])

$${}_2\Phi_1 \left[\begin{matrix} q^{-n}, b; q; z \\ c \end{matrix} \right] = \frac{(\frac{c}{b}; q)_n}{(c; q)_n} {}_3\Phi_2 \left[\begin{matrix} q^{-n}, b, \frac{bzq^{-n}}{c}; q; q \\ \frac{bq^{1-n}}{c}, 0 \end{matrix} \right],$$

with q, b, c, z being replaced respectively by $q^2, q^{-2l}, q^{-2l-2n}, q^{2l+2}$, yields

$$\begin{aligned} \varphi_l(q^{2n}) &= \frac{(q^{-2l-2n}; q^2)_n}{(q^{-2n}; q^2)_n} \cdot {}_2\Phi_1 \left[\begin{matrix} q^{-2n}; q^{-2l}; q^2; q^{2l+2} \\ q^{-2l-2n} \end{matrix} \right] \sim \\ &\sim q^{-2nl} \frac{(q^{2(l+1)}; q^2)_\infty}{(q^2; q^2)_\infty} \cdot {}_1\Phi_0[q^{-2l}; q^2; q^{2(l+1)}] = \\ &= q^{-2nl} \frac{(q^{2(l+1)}; q^2)_\infty}{(q^2; q^2)_\infty} \cdot \frac{(q^{2(l+1)}; q^2)_\infty}{(q^{2(2l+1)}; q^2)_\infty}. \end{aligned}$$

Now it remains to refer to the definition of the q -gamma-function. $\square$

In the special case $f = 1$ proposition 6.3 follows from lemma 6.4. The general case reduces to the above special case via an application of a quantum symmetry argument, which is described in [19, Theorem 3.7].

7 Fourier transform

It follows from the definitions that the integral operator with kernel $K = \sum_i k_i'' \otimes k_i'$ is conjugate to the integral operator with the kernel $K^t = \sum_i k_i'^* \otimes k_i''^*$. Note that the conjugate to the unitary is an inverse operator.

Recall [12] the heuristic argument that leads to the Fourier transform. Proposition 6.1 allows one to obtain eigenfunctions of $\square$. It is natural to expect that "any" function u admits a decomposition in eigenfunctions of $\square$, and that the associated Fourier operator is unitary.

Impose the notations: $h = -2 \ln q$,

$$\begin{aligned} P_\gamma^t &= (q^2 z^* \zeta; q^2)_{-\gamma} (z \zeta^*; q^2)_{-\gamma} (1 - \zeta \zeta^*)^\gamma, \\ c(l) &= \Gamma_{q^2}(2l+1) / (\Gamma_{q^2}(l+1))^2. \end{aligned}$$

It is shown in [19, section 5] that, just as in the standard representation theory (see [12]), one has

Proposition 7.1 *Consider the Borel measure $d\sigma$ on $[0, \frac{\pi}{h}]$, given by*

$$d\sigma(\rho) = \frac{1}{2\pi} \cdot \frac{h \cdot e^h}{e^h - 1} c(-\frac{1}{2} + i\rho)^{-1} \cdot c(-\frac{1}{2} - i\rho)^{-1} d\rho.$$

The integral operators

$$\begin{aligned} u(z) &\mapsto \int_{U_q} P_{\frac{1}{2}-i\rho}^t(z, \zeta) u(\zeta) d\nu, \\ f(e^{i\theta}, \rho) &\mapsto \int_0^{\pi/h} \int_0^{2\pi} P_{\frac{1}{2}+i\rho}(z, e^{i\theta}) f(e^{i\theta}, \rho) \frac{d\theta}{2\pi} d\sigma(\rho) \end{aligned}$$

are extendable by continuity from the dense linear subspaces

$$D(U)_q \subset L^2(d\nu)_q, \quad C^\infty[0, \frac{\pi}{h}] \otimes \mathbb{C}[\partial U]_q \subset L^2(d\sigma) \otimes L^2(\frac{d\theta}{2\pi})$$

up to mutually inverse unitaries F, F^{-1} .

REMARK. The function $c(l)$, the measure $d\sigma(\rho)$ and the operator F are the quantum analogues for c -function of Harish-Chandra, Plancherel measure and Fourier transform respectively (see [12]).

8 The Berezin deformation of the quantum disc

We are going to use in the sequel bilinear operators $L : D(U)_q \times D(U)_q \rightarrow D(U)_q$ of the form

$$L : f_1 \times f_2 \rightarrow \sum_{ijk m=0}^{N(L)} a_{ijk m} \left(\left(\frac{\partial(r)}{\partial z^*} \right)^i f_1 \right) z^{*j} z^k \left(\left(\frac{\partial(l)}{\partial z} \right)^m f_2 \right), \quad (8.1)$$

with $a_{ijk m} \in \mathbb{C}$. Such operators will be called q -bidifferential.

Our principal purpose is to construct the formal deformation of the multiplication law in $D(U)_q$. The new multiplication is to be a bilinear map

$$\begin{aligned} * : D(U)_q \times D(U)_q &\rightarrow D(U)_q[[t]], \\ * : f_1 \times f_2 &\mapsto f_1 \cdot f_2 + \sum_{i=1}^{\infty} t^i C_i(f_1, f_2), \end{aligned}$$

which satisfies the formal associativity condition

$$\sum_{i+k=m} C_i(f_1, C_k(f_2, f_3)) = \sum_{i+k=m} C_i(C_k(f_1, f_2), f_3)$$

(cf. [14]). When producing the new multiplication $*$, we follow F. Berezin [3]. The bilinear operators $C_j : D(U)_q \times D(U)_q \rightarrow D(U)_q$, $j \in \mathbb{N}$, will turn out to be q -bidifferential, and we shall give explicit formulae for them.

To begin with, choose a positive α and consider a linear functional $\nu_\alpha : \text{Pol}(\mathbb{C})_q \rightarrow \mathbb{C}$;

$$\int_{U_q} f d\nu_\alpha \stackrel{\text{def}}{=} \frac{1 - q^{4\alpha}}{1 - q^2} \cdot \int_{U_q} f \cdot (1 - zz^*)^{2\alpha+1} d\nu = (1 - q^{4\alpha}) \text{tr} T(f \cdot (1 - zz^*)^{2\alpha}).$$

Impose a norm $\|f\|_\alpha = \left(\int_{U_q} f^* f d\nu_\alpha \right)^{1/2}$ on $\text{Pol}(\mathbb{C})_q$. Let $L_{q,\alpha}^2$ stand for the completion of $\text{Pol}(\mathbb{C})_q$ with respect to the above norm, and $H_{q,\alpha}^2$ for the linear span of monomials $z^j \in L_{q,\alpha}^2$, $j \in \mathbb{Z}_+$.

Lemma 8.1 *The monomials $\{z^m\}_{m \in \mathbb{Z}_+}$ are pairwise orthogonal in $H_{q,\alpha}^2$, and $\|z^m\|_\alpha = ((q^2; q^2)_m / (q^{4\alpha+2}; q^2)_m)^{1/2}$.*

Proof. The pairwise orthogonality of the monomials z^m is obvious;

$$\begin{aligned} \|z^m\|_\alpha^2 &= (1 - q^{4\alpha}) \cdot \text{tr} T(z^{*m} z^m (1 - zz^*)^{2\alpha}) = \\ &= \frac{1 - q^{4\alpha}}{1 - q^2} \int_0^1 (q^2 y; q^2)_m \cdot y^{2\alpha-1} d_{q^2} y = \\ &= \frac{1 - q^{4\alpha}}{1 - q^2} \cdot \frac{\Gamma_{q^2}(2\alpha) \cdot \Gamma_{q^2}(m+1)}{\Gamma_{q^2}(m+2\alpha+1)} = \frac{(q^2; q^2)_m}{(q^{4\alpha+2}; q^2)_m}. \end{aligned}$$

We have used the well known [10, §1.11] identity

$$\int_0^1 t^{\beta-1} \cdot (tq^2; q^2)_{\alpha-1} d_{q^2} t = \frac{\Gamma_{q^2}(\beta) \Gamma_{q^2}(\alpha)}{\Gamma_{q^2}(\alpha + \beta)}.$$

□

Corollary 8.2 *Let $\widehat{z}$ be the operator of multiplication by z in $H_{q,\alpha}^2$, and $\widehat{z}^*$ the conjugate operator. Then $\widehat{z}$, $\widehat{z}^*$ are bounded, and*

$$\widehat{z}^* \widehat{z} = q^2 \widehat{z} \widehat{z}^* + 1 - q^2 + q^{4\alpha} \cdot \frac{1 - q^2}{1 - q^{4\alpha}} (1 - \widehat{z} \widehat{z}^*) (1 - \widehat{z}^* \widehat{z}). \quad (8.2)$$

Proof follows from

$$\begin{aligned}\widehat{z} : z^m &\mapsto z^{m+1}, & m \in \mathbb{Z}_+; \\ \widehat{z}^* : 1 &\mapsto 0, & \widehat{z}^* : z^m \mapsto \frac{1 - q^{2m}}{1 - q^{4\alpha+2m}} z^{m-1}, & m \in \mathbb{N}.\end{aligned}$$

In fact,

$$\begin{aligned}(1 - \widehat{z}\widehat{z}^*)^{-1} : z^m &\mapsto ((q^{-2m} - q^{4\alpha})/(1 - q^{4\alpha}))z^m, \\ (1 - \widehat{z}^*\widehat{z})^{-1} : z^m &\mapsto ((q^{-2m-2} - q^{4\alpha})/(1 - q^{4\alpha}))z^m.\end{aligned}$$

Hence $(1 - \widehat{z}\widehat{z}^*)^{-1} = q^2(1 - \widehat{z}^*\widehat{z})^{-1} - q^{4\alpha} \frac{1-q^2}{1-q^{4\alpha}}$. $\square$

Lemma 8.1 and corollary 8.2 were proved in the work by S. Klimek and A. Lesniewski [13] on two-parameter quantization of the disc beyond the framework of perturbation theory.

To every element $f = \sum a_{ij} z^i z^{*j} \in \text{Pol}(\mathbb{C})$ we associate the linear operator $\widehat{f} = \sum a_{ij} \widehat{z}^i \widehat{z}^{*j}$ in $H_{q,\alpha}^2$. The formal deformation of the multiplication law in the algebra of functions in the quantum disc will be derived via an application of "Berezin quantization procedure" $f \mapsto \widehat{f}$ to the ordinary multiplication in the algebra of linear operators.

More exactly, (8.2) allows one to get a formal asymptotic expansion

$$\widehat{f}_1 \cdot \widehat{f}_2 = \widehat{f_1 \cdot f_2} + \sum_{k=1}^{\infty} q^{4\alpha k} C_k(\widehat{f_1}, \widehat{f_2}). \quad f_1, f_2 \in \text{Pol}(\mathbb{C})_q,$$

with $C_k : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q \rightarrow \text{Pol}(\mathbb{C})_q$, $k \in \mathbb{N}$, bilinear maps. In this way, we get a formal deformation

$$\begin{aligned}* : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q &\rightarrow \text{Pol}(\mathbb{C})_q[[t]]; \\ f_1 * f_2 &= f_1 \cdot f_2 + \sum_{k=1}^{\infty} t^k \cdot C_k(f_1, f_2); \quad f_1, f_2 \in \text{Pol}(\mathbb{C})_q.\end{aligned}$$

We present an explicit formula for the multiplication $*$, and thus also for bilinear maps C_k , $k \in \mathbb{N}$. Let $\widetilde{\square}$ be a linear operator in $\text{Pol}(\mathbb{C})_q^{op} \otimes \text{Pol}(\mathbb{C})_q$ given by

$$\widetilde{\square} = q^{-2}(1 - (1 + q^{-2})z^* \otimes z + q^{-2}z^{*2} \otimes z^2) \frac{\partial^{(r)}}{\partial z^*} \otimes \frac{\partial^{(l)}}{\partial z},$$

and $m : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q \rightarrow \text{Pol}(\mathbb{C})_q$, $m : \psi_1 \otimes \psi_2 \rightarrow \psi_1 \psi_2$ the multiplication in $\text{Pol}(\mathbb{C})_q$.

Theorem 8.3 *For all $f_1, f_2 \in \text{Pol}(\mathbb{C})_q$*

$$f_1 * f_2 = (1 - t) \sum_{j \in \mathbb{Z}_+} t^j \cdot m(p_j(\widetilde{\square}) f_1 \otimes f_2),$$

with

$$p_j(\widetilde{\square}) = \sum_{k=0}^j \frac{(q^{-2j}; q^2)_k}{(q^2; q^2)_k^2} q^{2k} \cdot \prod_{i=0}^{k-1} (1 - q^{2i}((1 - q^2)^2 \cdot \widetilde{\square} + 1 + q^2) + q^{4i+2}).$$

The proof can be found in [20, Theorem 8.4].

EXAMPLE. For all $f_1, f_2 \in \text{Pol}(\mathbb{C})_q$

$$f_1 * f_2 = f_2 \cdot f_1 + t \cdot (q^{-2} - 1) \frac{\partial^{(r)} f_1}{\partial z^*} (1 - z^* z)^2 \frac{\partial^{(l)} f_2}{\partial z} + O(t^2).$$

It is worthwhile to note that the formal associativity of the multiplication $*$ follows from the associativity of multiplication in the algebra of linear operators.

Corollary 8.4 *The bilinear operators C_k are of the form (8.1) and are extendable by continuity up to q -bidifferential operators $C_k : D(U)_q \times D(U)_q \rightarrow D(U)_q$.*

The above q -bidifferential operators determine a formal deformation of the multiplication in $D(U)_q$. The formal associativity of the newly formed multiplication $* : D(U)_q \times D(U)_q \rightarrow D(U)_q[[t]]$ follows from the formal associativity of the previous multiplication $* : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q \rightarrow \text{Pol}(\mathbb{C})_q[[t]]$.

Finally, note that our proof of theorem 8.3 is based on the properties of some q -analogue for Berezin transform [21].

9 Appendix. On q -analogue of the Green formula

Consider the two-sided ideal $J \in \text{Pol}(\mathbb{C})_q$ generated by the element $1 - zz^* \in \text{Pol}(\mathbb{C})_q$, and the commutative quotient algebra $\mathbb{C}[\partial U]_q \stackrel{\text{def}}{=} \text{Pol}(\mathbb{C})_q / J$. Its elements will be identified with the corresponding polynomials on the circle ∂U . The image $f|_{\partial U}$ of $f \in \text{Pol}(\mathbb{C})_q$ under the canonical homomorphism $\text{Pol}(\mathbb{C})_q \rightarrow \mathbb{C}[\partial U]_q$ will be called a restriction of f onto the boundary of the quantum disc.

Define the integral $\Omega(\mathbb{C})_q^{(1,0)} \rightarrow \mathbb{C}$ by

$$\int_{\partial U} dz \cdot f \stackrel{\text{def}}{=} 2\pi i \int_{\partial U} (z \cdot f)|_{\partial U} d\nu, \quad f \in \text{Pol}(\mathbb{C})_q,$$

with

$$\int_{\partial U} \psi d\nu \stackrel{\text{def}}{=} \int_0^{2\pi} \psi(e^{i\theta}) \frac{d\theta}{2\pi}.$$

Proposition 9.1 *For all $\psi \in \Omega(\mathbb{C})_q^{(0,1)}$ one has*

$$\int_{U_q} \bar{\partial} \psi = \int_{\partial U} \psi. \tag{9.1}$$

REMARK. The integral $\int_{U_q} dz \cdot f \cdot dz^* = -2i\pi \int_{U_q} f d\mu$ introduced in section 4 for $f \in D(U)_q$, is extendable by continuity onto all (1,1)-forms $dz \cdot f \cdot dz^*$ with

$$f = \sum_{m>0} z^m \psi_m(y) + \psi_0(y) + \sum_{m>0} \psi_{-m}(y) z^{*m} \in D(U)_q',$$

such that $\sum_{m \in \mathbb{Z}_+} |\psi_0(q^{2m})| q^{2m} < \infty$. Under these assumptions one also has

$$\int_{U_q} dz dz^* f = \int_{U_q} dz \cdot f \cdot dz^* = \int_{U_q} f dz dz^*. \quad (9.2)$$

Proof. Recall that (see (2.2))

$$\psi = dz \left(\sum_{m>0} z^m \psi_m(y) + \psi_0(y) + \sum_{m>0} \psi_{-m}(y) z^{*m} \right). \quad (9.3)$$

We can restrict ourselves to the case $\psi = dz \psi_{-1}(y) z^*$, since this is the only term in (9.3) which could make a non-zero contribution to (9.1).

It follows from the definitions that $\bar{\partial} \psi = dz \cdot f(y) dz^*$, with

$$f(y) = \psi_{-1}(y) - q^{-2} \frac{\psi_{-1}(q^{-2}y) - \psi_{-1}(y)}{q^{-2}y - y} (1 - y).$$

In fact, $\bar{\partial} y = \bar{\partial}(1 - zz^*) = -z dz^*$. Hence $\bar{\partial} y^m = \sum_{j=0}^{m-1} y^j (-z dz^*) y^{m-1-j} = -\frac{1-q^{2m}}{1-q^2} z y^{m-1} dz^*$.

That is, for any polynomial $p(y)$ one has

$$\bar{\partial} p(y) = -z \frac{p(y) - p(q^2 y)}{y - q^2 y} \cdot dz^*. \quad (9.4)$$

(Note that the validity of (9.4) for polynomials already implies its validity for all distributions). Finally,

$$\bar{\partial}(dz \psi_{-1}(y) z^*) = dz \left(-z \frac{\psi_{-1}(y) - \psi_{-1}(q^2 y)}{y - q^2 y} z^* + \psi_{-1}(y) \right) dz^*.$$

On the other hand, $-z \frac{\psi_{-1}(y) - \psi_{-1}(q^2 y)}{y - q^2 y} z^* + \psi_{-1}(y) = f(y)$, since $zy = q^{-2}yz$, $zz^* = 1 - y$.

If one assumes $\psi_{-1}(0) = 0$, it is easy to show that $\sum_{n \in \mathbb{Z}_+} f(q^{2n}) q^{2n} = 0$. Hence, in this case $\int_{U_q} \bar{\partial} \psi = \int_{\partial U} \psi = 0$. Thus, Proposition 9.1 is proved for all $(1,0)$ -forms from some linear subspace of codimensionality 1. Now it remains to prove (9.1) in the special case $\psi = dz \cdot z^*$. $\square$

Corollary 9.2 *If $\psi \in \Omega(U)_q^{(1,0)}$, then $\int_{U_q} \bar{\partial} \psi = 0$.*

References

- [1] W. B. Arveson, "Subalgebras of C^* -algebras", *Acta Mathematicae*, 1969, **123**, No 3-4, 141-224.
- [2] M. Beals, C. Fefferman, R. Grossman, "Strictly pseudoconvex domains in $\mathbb{C}^n$ ", *Bull. Amer. Math. Soc.*, **8** (1983), No 2, 125 - 322.

- [3] F. Berezin, "General concept of quantization", *Commun. Math. Phys.*, **40** (1975), 153 – 174.
- [4] B. Blackadar. "K-theory for Operator Algebras", Springer-Verlag, New York, Heidelberg, Berlin, 1986.
- [5] O. Bratteli, D. Robinson. "Operator Algebras and Quantum Statistical Mechanics I", Springer-Verlag, New York, Heidelberg, Berlin, 1979.
- [6] M. Chaichian, P. P. Kulish. "Quantum group covariant systems", q-alg/9512017.
- [7] V. Chari, A. Pressley. "A Guide to Quantum Groups", Cambridge Univ. Press, 1995.
- [8] V. G. Drinfeld, "Quantum groups", in *Proceedings of the International Congress of Mathematicians*, Berkeley, 1986, A. M. Gleason (ed.), 1987, American Mathematical Society, Providence RI, 798 – 820.
- [9] P. G. A. Floris. "Addition Formula for q -adic Polynomials", Report W94-24, Mathematical Institute, University of Leiden, 10 p.
- [10] G. Gasper, M. Rahman. "Basic Hypergeometric Series", Cambridge University Press, Cambridge, 1990.
- [11] S. Helgason. "Differential Geometry and Symmetric Spaces", Acad. Press, N.Y. – London, 1962.
- [12] S. Helgason. "Groups and Geometric Analysis", Academic Press Inc., N.Y. – London, 1984.
- [13] S. Klimek, A. Lesniewski, "A two-parameter quantum deformation of the unit disc", *J. Funct. Anal.* **115**, (1993), 1 – 23.
- [14] A. Lichnerowicz. "Deformation of algebras associated with a symplectic manifold", in *Differential Geometry and Mathematical Physics*, M. Cahen et al.(eds.), D. Reidel Publishing Company, 1983, 69 – 83.
- [15] G. Nagy, A. Nica. "On the "quantum disc" and a "non-commutative circle", in: *Algebraic Methods on Operator Theory*, R. E. Curto, P. E. T. Jorgensen (eds.), Birkhauser, Boston, 1994, 276 – 290.
- [16] S. Sinel'shchikov, L. Vaksman. "On q -analogues of Bounded Symmetric Domains and Dolbeault Complexes", *Mathematical Physics, Analysis and Geometry*, Kluwer Academic Publishers, V.1, No.1, 1998, 75–100, q-alg 9703005.
- [17] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. "On function theory in quantum disc: covariance", math.QA/9808037.
- [18] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. "On function theory in quantum disc: invariant kernels", math.QA/9808047.
- [19] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. "On function theory in quantum disc: q -differential equations and Fourier transform", math.QA/9809002.
- [20] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. "On function theory in quantum disc: q -analogue of Berezin transform", math.QA/9809018.
- [21] A. Unterberger, H. Upmeyer, "The Berezin transform and invariant differential operators", *Comm. Math. Phys.*, **164** (1994), 563 – 598.
- [22] L. L. Vaksman, "Intertwining operators and quantum homogeneous spaces", *Matematicheskaya Fizika. Analiz. Geometriya*, **1** (1994), n 3/4, 392 – 409; q-alg/9511007.
- [23] L. L. Vaksman, "Integral intertwining operators and quantum homogeneous spaces", *Theoretical and Mathematical Physics*, **105** (1995), No 3, 355 – 363.

- [24] Chong-Sun Chu, Pei-Ming Ho, Bruno Zumino. "Some complex quantum manifolds and their geometry", hep-th/9608188, 1996, 49 p.p.

QUANTUM DISC: THE BASIC STRUCTURES

D. Shklyarov

S. Sinel'shchikov

L. Vaksman

1 Introduction

The theory of quantum groups and their homogeneous spaces is in an extensive progress since mid-80's [4]. An important class of such homogeneous spaces is formed by q -analogues of Cartan domains. The simplest Cartan domain is the unit disc $\mathbb{U}$ in $\mathbb{C}$; the present work is devoted to its q -analogue, the quantum disc. We intend to introduce the basic notions of the theory of q -Cartan domains while restricting ourselves to this simplest case. We hope this text could facilitate reading works on general q -Cartan domains and will allow better understanding the results of non-commutative function theory in quantum disc [13]. The concluding part of our work contains references to the papers of other authors devoted to the related topics. Among those one should emphasize the work by K. Schmüdgen and A. Schüler [12] which introduces the differential calculus in quantum disc which we use below.

The next section recalls the basic notions of the quantum group theory after the lecture notes [8] and the monograph [2].

2 The $\ast$ -Hopf algebra $U_q\mathfrak{su}_{1,1}$

Everywhere in the sequel we assume $0 < q < 1$, and $\mathbb{C}$ is implicit as a ground field. The algebras under consideration are supposed to be unital, unless the contrary is stated explicitly.

Recall the definition of the Hopf algebra $U_q\mathfrak{sl}_2$. It is determined by its generators E, F, K, K^{-1} and the following relations:

$$\begin{aligned} KK^{-1} &= K^{-1}K = 1, \\ K^{\pm 1}E &= q^{\pm 2}EK^{\pm 1}, & K^{\pm 1}F &= q^{\mp 2}FK^{\pm 1}, \\ EF - FE &= \frac{K - K^{-1}}{q - q^{-1}}. \end{aligned}$$

A $U_q\mathfrak{sl}_2$ -module V is said to be $\mathbb{Z}$ -weight if

$$V = \bigoplus_{\mu \in \mathbb{Z}} V_{\mu}, \quad V_{\mu} = \{v \in V \mid K^{\pm 1}v = q^{\pm \mu}v\}.$$

We restrict our considerations to $U_q\mathfrak{sl}_2$ -modules of the above form and define a linear operator H by $H|_{V_{\mu}} = \mu$. Obviously, $K^{\pm 1}v = q^{\pm H}v$, $v \in V$, and the following relations

This is an expanded version of a lecture delivered at a seminar for undergraduate students in Kharkov, April 2000.

are valid in the algebra of linear operators:

$$\begin{aligned} HE - EH &= 2E, & HF - FH &= -2F, \\ EF - FE &= \frac{q^H - q^{-H}}{q - q^{-1}}. \end{aligned}$$

A formal passage to a limit as $q \rightarrow 1$ leads to the determining relations of the ordinary universal enveloping algebra $U\mathfrak{sl}_2$.

Within the category of modules over an algebra A , the operation of tensor product is not defined; it is also not clear which A -module is to be treated as trivial and how to define an A -module V^* dual to an A -module V . These three problems are solvable via equipping A with the structure of Hopf algebra. Specifically, one has to distinguish

- a homomorphism $\Delta : A \rightarrow A \otimes A$, called a comultiplication and used for producing a tensor product of A -modules V', V'' :

$$A \xrightarrow{\Delta} A \otimes A \rightarrow \text{End}(V') \otimes \text{End}(V'') \simeq \text{End}(V' \otimes V'');$$

- a homomorphism $\varepsilon : A \rightarrow \mathbb{C}$, called a counit, to be used for producing the trivial A -module $\mathbb{C}$;
- an antihomomorphism $S : A \rightarrow A$, called an antipode, to be used for producing a dual A -module:

$$(\xi f)(v) \stackrel{\text{def}}{=} f(S(\xi)v), \quad \xi \in A, \quad v \in V, \quad f \in V^*.$$

Of course, the definition of a Hopf algebra includes several assumptions on $(A, \Delta, \varepsilon, S)$ which provide the habitual properties of the operations of tensor product and passage to a dual module.

Introduce a comultiplication $\Delta : U_q\mathfrak{sl}_2 \rightarrow U_q\mathfrak{sl}_2 \otimes U_q\mathfrak{sl}_2$, a counit $\varepsilon : U_q\mathfrak{sl}_2 \rightarrow \mathbb{C}$, and an antipode $S : U_q\mathfrak{sl}_2 \rightarrow U_q\mathfrak{sl}_2$ as follows:

$$\begin{aligned} \Delta(E) &= E \otimes 1 + K \otimes E, & \Delta(F) &= F \otimes K^{-1} + 1 \otimes E, & \Delta(K^{\pm 1}) &= K^{\pm 1} \otimes K^{\pm 1}; \\ \varepsilon(E) &= \varepsilon(F) = 0, & \varepsilon(K^{\pm 1}) &= 1; \\ S(E) &= -K^{-1}E, & S(F) &= -FK, & S(K^{\pm 1}) &= K^{\mp 1}. \end{aligned}$$

Equip $U_q\mathfrak{sl}_2$ with an antilinear involution $*$: $U_q\mathfrak{sl}_2 \rightarrow U_q\mathfrak{sl}_2$ given by

$$E^* = -KF, \quad F^* = -EK^{-1}, \quad K^* = K.$$

It follows from the definitions that

$$\Delta(\xi^*) = \Delta(\xi)^{* \otimes *}, \quad \xi \in U_q\mathfrak{sl}_2.$$

The $*$ -Hopf-algebra $(U_q\mathfrak{sl}_2, *)$ we thus obtain will be denoted by $U_q\mathfrak{su}_{1,1}$. It is a q -analogue of the $*$ -Hopf-algebra $U\mathfrak{su}_{1,1} \otimes_{\mathbb{R}} \mathbb{C}$. A $U_q\mathfrak{su}_{1,1}$ -module V is said to be unitarizable if for some scalar product (i.e. positive sesquilinear form) in V

$$(\xi v_1, v_2) = (v_1, \xi^* v_2), \quad \xi \in U_q\mathfrak{su}_{1,1}, \quad v_1, v_2 \in V.$$

Obviously, a tensor product of unitarizable $U_q\mathfrak{su}_{1,1}$ -modules is a unitarizable $U_q\mathfrak{su}_{1,1}$ -module.

3 Examples of $U_q\mathfrak{sl}_2$ -module algebras: $\mathbb{C}[z]_q$, Λ_q , $A_{1,q}$

Consider a $U_q\mathfrak{sl}_2$ -module V and its element v . This element is called $U_q\mathfrak{sl}_2$ -invariant if the linear map $\mathbb{C} \rightarrow V$, $z \mapsto zv$, is a morphism of $U_q\mathfrak{sl}_2$ -modules (equivalently, $Ev = Fv = Hv = 0$).

Consider an algebra $\mathcal{F}$ and a linear map $m : \mathcal{F} \otimes \mathcal{F} \rightarrow \mathcal{F}$, $m : f_1 \otimes f_2 \mapsto f_1 f_2$ determined by the multiplication law in $\mathcal{F}$: $f_1 \times f_2 \mapsto f_1 f_2$. $\mathcal{F}$ is called a $U_q\mathfrak{sl}_2$ -module algebra if it is equipped with a structure of $U_q\mathfrak{sl}_2$ -module and the multiplication $m : \mathcal{F} \otimes \mathcal{F} \rightarrow \mathcal{F}$ is a morphism of $U_q\mathfrak{sl}_2$ -modules (equivalently, the following q -analogue of the Leibnitz rule is valid:

$$\begin{aligned} E(f_1 f_2) &= (E f_1) f_2 + (q^H f_1)(E f_2), \\ F(f_1 f_2) &= (F f_1)(q^{-H} f_2) + f_1(F f_2), \\ H(f_1 f_2) &= (H f_1) f_2 + f_1(H f_2). \end{aligned}$$

Under the presence of the unit element $1 \in \mathcal{F}$ there is an additional requirement of its $U_q\mathfrak{sl}_2$ -invariance: $E1 = F1 = H1 = 0$.

In a similar way, the notion of $U_q\mathfrak{sl}_2^{\text{op}}$ -module coalgebra is introduced, with $U_q\mathfrak{sl}_2^{\text{op}}$ being the Hopf algebra deduced from $U_q\mathfrak{sl}_2$ via replacing its comultiplication Δ with the opposite one Δ^{op} (if $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$ then $\Delta^{\text{op}}(\xi) = \sum_j \xi''_j \otimes \xi'_j$).

An important example of a $U_q\mathfrak{sl}_2^{\text{op}}$ -module coalgebra is built from the Verma module of zero weight $M(0)$. This module is determined by its generator $v(0) \in M(0)$ and the relations $Ev(0) = 0$, $K^{\pm 1}v(0) = v(0)$. Obviously, there exists a unique morphism of $U_q\mathfrak{sl}_2^{\text{op}}$ -modules $\delta : M(0) \rightarrow M(0) \otimes M(0)$ such that $\delta v(0) = v(0) \otimes v(0)$. Coassociativity of this comultiplication follows from

$$((v(0) \otimes v(0)) \otimes v(0) = v(0) \otimes (v(0) \otimes v(0))).$$

Note that the vector spaces $M(0)$, $M(0) \otimes M(0)$ are graded:

$$\begin{aligned} M(0) &= \bigoplus_j M(0)_j, \quad M(0)_j = \{v \in M(0) \mid Hv = 2jv\}, \\ (M(0) \otimes M(0))_j &= \bigoplus_{i+k=j} M(0)_i \otimes M(0)_k, \end{aligned}$$

and that the comultiplication preserves the degree of homogeneity:

$$\Delta : M(0)_j \rightarrow (M(0) \otimes M(0))_j,$$

with $M(0)_0 = \mathbb{C}v(0)$. That is why the dual graded vector space $M(0)^* \stackrel{\text{def}}{=} \bigoplus_j (M(0)_j)^*$ is a unital $\mathbb{Z}$ -graded algebra. By a construction, $M(0)$ is a $U_q\mathfrak{sl}_2^{\text{op}}$ -module coalgebra and $M(0)^*$ is a $U_q\mathfrak{sl}_2$ -module algebra. We are going to describe it by explicit formulae.

The elements $v(0)$, $S(F)v(0)$, $S(F^2)v(0)$, $\dots$, form a basis of the vector space $M(0)$. Hence there exists a unique element $z \in M(0)^*$ such that

$$z(S(F^j)v(0)) = \begin{cases} q^{1/2}, & j = 1, \\ 0, & j \neq 1. \end{cases}$$

It is easy to prove that the algebra $M(0)^*$ is isomorphic to $\mathbb{C}[z]$ and

$$Fz = q^{1/2}, \quad Ez = -q^{1/2}z^2, \quad K^{\pm 1}z = q^{\pm 2}z. \quad (3.1)$$

Thus we have equipped the polynomial algebra $\mathbb{C}[z]$ with a structure of $U_q\mathfrak{sl}_2$ -module algebra, to be denoted $\mathbb{C}[z]_q$ in the sequel.

Turn to producing a differential calculus on $\mathbb{C}[z]_q$. A duality argument like that we used to derive $\mathbb{C}[z]_q$ allows one to produce a differential $U_q\mathfrak{sl}_2$ -module algebra (Λ_q, d) . In this setting, the embeddings $\mathbb{C}[z]_q \hookrightarrow \Lambda_q$ and the differential $d : \Lambda_q \rightarrow \Lambda_q$ are morphisms of $U_q\mathfrak{sl}_2$ -modules. One can also prove that

$$dz \cdot dz = 0, \quad dz \cdot z = q^2 z \cdot dz.$$

See [16] for details. Note that this differential calculus is well known and can be derived from the Wess-Zumino differential calculus on the quantum $\mathbb{C}^2$ ($t_1 t_2 = q t_2 t_1$) via the localization procedure $z = t_2^{-1} t_1$ and further restriction onto the subalgebra of zero homogeneity degree elements.

To conclude, describe one more $U_q\mathfrak{sl}_2$ -module algebra, which is a q -analogue of the Weyl algebra A_1 .

Consider the linear operator $\frac{d}{dz}$ in $\mathbb{C}[z]_q$ given by $df = dz \cdot \left(\frac{d}{dz}f\right)$, $f \in \mathbb{C}[z]_q$. It is easy to demonstrate that

$$\frac{d}{dz} \cdot \widehat{z} - q^{-2} \widehat{z} \cdot \frac{d}{dz} = 1, \quad (3.2)$$

with $\widehat{z}$ being the left multiplication operator by z ($\widehat{z} : f \mapsto zf$) in $\mathbb{C}[z]_q$. The algebra determined by the two generators $\frac{d}{dz}$ and $\widehat{z}$ and the above relations is called the quantum Weyl algebra (q -oscillatory algebra) $A_{1,q}$. Our purpose is to equip it with a structure of a $U_q\mathfrak{sl}_2$ -module algebra. We use the fact that the algebra $\text{End}_{\mathbb{C}}(\mathbb{C}[z]_q)$ of all linear operators in $\mathbb{C}[z]_q$ is $U_q\mathfrak{sl}_2$ -module:

$$\xi(T) = \sum_j \xi'_j \cdot T \cdot S(\xi''_j) \quad \text{for} \quad \Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j.$$

We are to prove that the image of $A_{1,q}$ under the canonical embedding into $\text{End}_{\mathbb{C}}(\mathbb{C}[z]_q)$ appears to be a submodule of the $U_q\mathfrak{sl}_2$ -module $\text{End}_{\mathbb{C}}(\mathbb{C}[z]_q)$. This is a consequence of the following relations which describe the action of the generators $E, F, K^{\pm 1}$ on the generators $\widehat{z}, \frac{d}{dz}$:

$$\begin{aligned} E(\widehat{z}) &= -q^{1/2} \widehat{z}^2, & F(\widehat{z}) &= q^{1/2}, & K^{\pm 1}(\widehat{z}) &= q^{\pm 2} \widehat{z}, \\ E\left(\frac{d}{dz}\right) &= q^{-3/2}(q^{-2} + 1) \widehat{z} \frac{d}{dz}, & F\left(\frac{d}{dz}\right) &= 0, & K^{\pm 1}\left(\frac{d}{dz}\right) &= q^{\mp 2} \frac{d}{dz}. \end{aligned} \quad (3.3)$$

We restrict ourselves to proving the most intricate of these relations. It follows from the definitions that

$$\begin{aligned} \frac{df(z)}{dz} &= \frac{f(q^{-2}z) - f(z)}{q^{-2}z - z}, & Ef(z) &= -q^{1/2}z^2 \frac{f(z) - f(q^2z)}{z - q^2z}, \\ Ff(z) &= q^{1/2} \frac{f(q^{-2}z) - f(z)}{q^{-2}z - z}, & K^{\pm 1}f(z) &= f(q^{\pm 2}z). \end{aligned}$$

Therefore,

$$\begin{aligned} E \frac{d}{dz} f(z) &= -\frac{q^{1/2}}{(1-q^2)^2} (q^2 f(q^{-2}z) - (1+q^2)f(z) + f(q^2z)), \\ \frac{d}{dz} E f(z) &= -\frac{q^{1/2}}{(1-q^2)^2} (f(q^{-2}z) - (1+q^2)f(z) + q^2 f(q^2z)). \end{aligned}$$

It follows from the definition of $E \left(\frac{d}{dz} \right)$ that

$$E \left(\frac{d}{dz} \right) = E \frac{d}{dz} - K \frac{d}{dz} K^{-1} E = E \frac{d}{dz} - q^{-2} \frac{d}{dz} E,$$

hence

$$E \left(\frac{d}{dz} \right) f(z) = -\frac{q^{-3/2}}{(q^{-2}-1)^2} (1-q^{-4})(f(q^{-2}z) - f(z)) = q^{-3/2}(1+q^{-2}) \hat{z} \frac{df(z)}{dz}.$$

It follows from the above observations that $A_{1,q}$ is a $U_q \mathfrak{sl}_2$ -module algebra and the action of $U_q \mathfrak{sl}_2$ in $A_{1,q}$ is given by (3.3).

4 The $U_q \mathfrak{su}_{1,1}$ -module algebras $\text{Pol}(\mathbb{C})_q$, $\mathcal{D}(\mathbb{U})_q$

Consider a $U_q \mathfrak{sl}_2$ -module algebra F with an involution $*$. F is called a $U_q \mathfrak{su}_{1,1}$ -module algebra if the involutions agree:

$$(\xi f)^* = (S(\xi))^* f^*, \quad \xi \in U_q \mathfrak{su}_{1,1}, \quad f \in F.$$

We present below examples of such algebras.

Note first that

$$(S(E))^* = q^{-2} F, \quad (S(F))^* = q^2 E, \quad (S(K^{\pm 1}))^* = K^{\mp 1}.$$

Equip the vector space $\text{Pol}(\mathbb{C})_q = \mathbb{C}[z]_q \otimes \mathbb{C}[z^*]_q$ with an involution: $(f_2(z) \otimes f_1(z^*))^* = \overline{f_2}(z) \otimes \overline{f_1}(z^*)$, where bar denotes complex conjugation for the coefficients of polynomials. The involutions in $\text{Pol}(\mathbb{C})_q$ and $U_q \mathfrak{su}_{1,1}$ agree in the above sense. What remains is to equip $\text{Pol}(\mathbb{C})_q$ with a multiplication $m : \text{Pol}(\mathbb{C})_q \otimes \text{Pol}(\mathbb{C})_q \rightarrow \text{Pol}(\mathbb{C})_q$ which agree with the action of $U_q \mathfrak{sl}_2$ and the involution $*$ in $\text{Pol}(\mathbb{C})_q$. It was demonstrated in [16] that such multiplication can be derived from the morphism of $U_q \mathfrak{sl}_2$ -modules

$$\check{R} : \mathbb{C}[z^*]_q \otimes \mathbb{C}[z]_q \rightarrow \mathbb{C}[z]_q \otimes \mathbb{C}[z^*]_q$$

determined by the action of the universal R-matrix in $\mathbb{C}[z^*]_q \otimes \mathbb{C}[z]_q$ and a subsequent ordinary flip of tensor multiples. More precisely, $m = m_+ \otimes m_- (\text{id} \otimes \check{R} \otimes \text{id})$, with $m_{\pm}$ being the multiplications in $\mathbb{C}[z]_q$ and $\mathbb{C}[z^*]_q$, respectively.

To describe the action of the universal R-matrix one has well known Drinfeld's formulae [4]:

$$R = \exp_{q^2}((q^{-1} - q)E \otimes F) q^{-H \otimes H/2},$$

$$\text{with } \exp_t(x) = \sum_{n=1}^{\infty} x^n \left(\prod_{j=1}^n ((1-t^j)/(1-t)) \right)^{-1}.$$

An application of this relation allows one to prove (see [16]) that

$$z^*z = q^2zz^* + 1 - q^2. \quad (4.1)$$

Furthermore, $\text{Pol}(\mathbb{C})_q$ can be determined by the generators z, z^* and the relation (4.1), and the action of $E, F, K^{\pm 1}$ on z, z^* can be given by (3.1).

We need a positive invariant integral in the quantum disc. The problem is that in the classical case ($q = 1$) the positive $SU_{1,1}$ -invariant measure in the unit disc such integral is not defined on the polynomial algebra since

$$\int_{\mathbb{U}} (1 - |z|^2)^{-2} d\text{Re}z \wedge d\text{Im}z = \infty.$$

To produce an invariant integral, one uses an extension $\text{Fun}(\mathbb{U})_q$ of the $U_q\mathfrak{su}_{1,1}$ -module $*$ -algebra $\text{Pol}(\mathbb{C})_q$, derived by adding an element f_0 with the following properties:

$$z^*f_0 = 0, \quad f_0z = 0, \quad f_0^2 = f_0. \quad (4.2)$$

The involution is extended from $\text{Pol}(\mathbb{C})_q$ onto $\text{Fun}(\mathbb{U})_q$ in such a way that f_0 becomes a selfadjoint element: $f_0^* = f_0$.

To motivate this definition, let us consider a faithful irreducible $*$ -representation T of the $*$ -algebra $\text{Pol}(\mathbb{C})_q$. Such $*$ -representation is unique up to a unitary equivalence. In the standard basis $\{e_k\}_{k=0}^{\infty}$ of the Hilbert space $l^2(\mathbb{Z}_+)$ it can be given by

$$T(z)e_k = (1 - q^{2(k+1)})^{1/2} e_{k+1}, \quad T(z^*)e_k = \begin{cases} (1 - q^{2k})^{1/2} e_{k-1}, & k > 0, \\ 0, & k = 0. \end{cases}$$

It follows from the definition of the $*$ -algebra $\text{Fun}(\mathbb{U})_q$ that T is extendable up to a $*$ -representation $\tilde{T}$ of $\text{Fun}(\mathbb{U})_q$ and $\tilde{T}(f_0)$ appears to be an orthogonal projection onto the 'vacuum subspace' $\mathbb{C}e_0$.

Note that the system of equation $z^*f_0 = 0, f_0z = 0$ has a solution in the space of formal series $\mathcal{D}(\mathbb{U})'_q = \left\{ \sum_{j,k=0}^{\infty} c_{jk} z^j z^{*k} \mid c_{jk} \in \mathbb{C} \right\}$. Specifically,

$$f_0 = \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k-1)}}{(1 - q^2)(1 - q^4) \dots (1 - q^{2k})} z^k z^{*k}.$$

We thus obtain an embedding of vector spaces $\text{Fun}(\mathbb{U})_q \subset \mathcal{D}(\mathbb{U})'_q$. The vector space $\mathcal{D}(\mathbb{U})'_q$ with the coefficientwise convergence topology will be called the distribution space in the quantum disc. One can demonstrate that the involution and the $U_q\mathfrak{sl}_2$ -action are extendable by a continuity from $\text{Pol}(\mathbb{C})_q$ onto $\mathcal{D}(\mathbb{U})'_q$ and thus equip $\text{Fun}(\mathbb{U})_q$ with a structure of $U_q\mathfrak{su}_{1,1}$ -module algebra. The action of $E, F, K^{\pm 1}$ onto f_0 is described by the explicit formulae:

$$Ef_0 = -\frac{q^{1/2}}{1 - q^2} z f_0, \quad Ff_0 = -\frac{q^{1/2}}{q^{-2} - 1} f_0 z^*, \quad K^{\pm 1} f_0 = f_0.$$

The two-sided ideal $\mathcal{D}(\mathbb{U})_q = \text{Fun}(\mathbb{U})_q f_0 \text{Fun}(\mathbb{U})_q$ works as an algebra of finite functions in the quantum disc. Note that there exists a non-degenerate pairing of $\mathcal{D}(\mathbb{U})_q$ and $\mathcal{D}(\mathbb{U})'_q$.

Of course, the subalgebra $\mathcal{D}(\mathbb{U})_q$ is a $U_q\mathfrak{su}_{1,1}$ -module algebra. It is just the domain where a positive $U_q\mathfrak{sl}_2$ -invariant integral is to be defined.

Consider the linear span $\mathcal{L} \subset l^2(\mathbb{Z}_+)$ of the standard basis $\{e_k\}_{k=0}^\infty$ together with a linear operator Γ in $\mathcal{L}$,

$$\Gamma e_k = q^{-2k} e_k, \quad k \in \mathbb{Z}_+.$$

The linear functional

$$\int_{\mathbb{U}_q} f d\nu \stackrel{\text{def}}{=} (1 - q^2) \text{Tr}(\tilde{T}(f)\Gamma)$$

is well defined on $\mathcal{D}(\mathbb{U})_q$; it is positive and $U_q\mathfrak{sl}_2$ -invariant (more precisely,

$$\begin{aligned} \int_{\mathbb{U}_q} f^* f d\nu &> 0 \quad \text{for } f \neq 0, \quad \text{and} \\ \int_{\mathbb{U}_q} (\xi f) d\nu &= \varepsilon(\xi) \int_{\mathbb{U}_q} f d\nu, \quad f \in \mathcal{D}(\mathbb{U})_q, \quad \xi \in U_q\mathfrak{sl}_2. \end{aligned}$$

Note finally an outward similarity of the above invariant integral in $\mathcal{D}(\mathbb{U})_q$ and the well known in the quantum group theory invariant integral $\text{tr}_q : \text{End}_{\mathbb{C}}(V) \rightarrow \mathbb{C}$, with V being a $U_q\mathfrak{sl}_2$ -module. This similarity is due to the fact that $\mathcal{D}(\mathbb{U})_q$ is embedable into a one-parameter family of $U_q\mathfrak{su}_{1,1}$ -module algebras $\text{End}_{\mathbb{C}} V_t$ as a limit object. In turn, the origin of the new deformation parameter t is in the Berezin quantization [14].

5 Differential calculi on $\text{Pol}(\mathbb{C})_q$, $\text{Fun}(\mathbb{U})_q$

A universal R-matrix was used in the previous section to pass from the $U_q\mathfrak{sl}_2$ -module algebra $\mathbb{C}[z]_q$ to the $U_q\mathfrak{su}_{1,1}$ -module algebra $\text{Pol}(\mathbb{C})_q$. A similar construction described in [16] allows one to pass from the $U_q\mathfrak{sl}_2$ -module algebra (Λ_q, d) as in section 3 to the $U_q\mathfrak{su}_{1,1}$ -module differential algebra described below.

Let Ω_q be the algebra determined by the generators z, z^*, dz, dz^* and the relations

$$\begin{aligned} z^* z &= q^2 z z^* + 1 - q^2, \\ dz \cdot z &= q^2 z \cdot dz, & dz^* \cdot z^* &= q^{-2} z^* \cdot dz^*, \\ dz \cdot z^* &= q^{-2} z^* \cdot dz, & dz^* \cdot z &= q^2 z \cdot dz^*, \\ dz \cdot dz &= dz^* \cdot dz^* = 0, & dz^* \cdot dz &= -q^2 dz \cdot dz^*. \end{aligned}$$

Equip this algebra with an involution $*$: $z \mapsto z^*$; $*$: $dz \mapsto dz^*$ and a $\mathbb{Z}_2$ -grading $\deg(z) = \deg(z^*) = 0$, $\deg(dz) = \deg(dz^*) = 1$. There exists a unique differentiation d of the superalgebra Ω_q such that $d^2 = 0$ and $d : z \mapsto dz$, $d : z^* \mapsto dz^*$. The $*$ -algebra $\text{Pol}(\mathbb{C})_q$ is uniquely embedable into Ω_q . There exists a unique extension of the structure of $U_q\mathfrak{su}_{1,1}$ -module algebra from $\text{Pol}(\mathbb{C})_q$ onto Ω_q in such a way that the differential d appears to be a morphism of $U_q\mathfrak{su}_{1,1}$ -modules.

The $*$ -algebra $\text{Fun}(\mathbb{U})_q$ was derived from $\text{Pol}(\mathbb{C})_q$ via adding the element f_0 which satisfies (4.2). Besides that, one should also include to the definition of the algebra of differential forms the relations $f_0 \cdot dz = dz \cdot f_0$, $f_0 \cdot dz^* = dz^* \cdot f_0$; the definition of the differential d should also include the equality

$$df_0 = -\frac{1}{1 - q^2} (dz f_0 z^* + z f_0 dz^*).$$

In [15, section 5] these relations and the latter equality were obtained via the embedding $\text{Fun}(\mathbb{U})_q \subset \mathcal{D}(\mathbb{U})'_q$. Also, it was proved there that the structure of $U_q\mathfrak{su}_{1,1}$ -module algebra is uniquely extendable from $\text{Fun}(\mathbb{U})_q$ onto the above differential $*$ -algebra.

Note that, similarly to the case $q = 1$, one has a bigrading of the algebra of differential forms:

$$\deg(z) = \deg(z^*) = \deg(f_0) = (0, 0), \quad \deg(dz) = (1, 0), \quad \deg(dz^*) = (0, 1).$$

The differential d admits a decomposition

$$d = \partial + \bar{\partial}$$

as a sum of holomorphic and antiholomorphic differentials, to be defined in the standard way. For example,

$$\partial f_0 = -\frac{1}{1-q^2} dz \cdot f_0 \cdot z^*, \quad \bar{\partial} f_0 = -\frac{1}{1-q^2} z \cdot f_0 \cdot dz^*.$$

Of course, $\partial^2 = \partial\bar{\partial} + \bar{\partial}\partial = \bar{\partial}^2 = 0$.

Define the operators $\frac{\partial}{\partial z}, \frac{\partial}{\partial z^*}$ in $\text{Pol}(\mathbb{C})_q$ by

$$\partial f = dz \cdot \frac{\partial f}{\partial z}, \quad \bar{\partial} f = dz^* \cdot \frac{\partial f}{\partial z^*},$$

and the operators $\widehat{z}, \widehat{z}^*$ by $\widehat{z}f = zf, \widehat{z}^*f = z^*f$. These linear operators are extendable by a continuity onto the entire space of distributions $\mathcal{D}(\mathbb{U})'_q$ and satisfies the following commutation relations:

$$\begin{aligned} \widehat{z}^*\widehat{z} &= q^2\widehat{z}\widehat{z}^* + 1 - q^2, \\ \frac{\partial}{\partial z}\widehat{z} - q^{-2}\widehat{z}\frac{\partial}{\partial z} &= 1, & \frac{\partial}{\partial z}\widehat{z}^* &= q^2\widehat{z}^*\frac{\partial}{\partial z}, \\ \frac{\partial}{\partial z^*}\widehat{z} &= q^{-2}\widehat{z}\frac{\partial}{\partial z^*}, & \frac{\partial}{\partial z^*}\widehat{z}^* - q^2\widehat{z}^*\frac{\partial}{\partial z^*} &= 1, \\ \frac{\partial}{\partial z^*}\frac{\partial}{\partial z} &= q^2\frac{\partial}{\partial z}\frac{\partial}{\partial z^*}. \end{aligned}$$

These commutation relations follow from

$$\begin{aligned} z^*z &= q^2zz^* + 1 - q^2, \\ \partial(zf) &= dz \cdot f + zdz\frac{\partial f}{\partial z} = dz\left(f + q^{-2}z\frac{\partial f}{\partial z}\right), \\ \partial(z^*f) &= z^*dz\frac{\partial f}{\partial z} = q^2dz \cdot z^*\frac{\partial f}{\partial z}, \\ \bar{\partial}(zf) &= zdz^*\frac{\partial f}{\partial z^*} = q^{-2}dz^* \cdot z \cdot \frac{\partial f}{\partial z^*}, \\ \bar{\partial}(z^*f) &= dz^* \cdot f + z^* \cdot dz^* \cdot \frac{\partial f}{\partial z^*} = dz^*\left(f + q^2z^*\frac{\partial f}{\partial z^*}\right), \\ \partial\bar{\partial}f &= \partial\left(dz^*\frac{\partial f}{\partial z^*}\right) = -dz^*dz\frac{\partial}{\partial z}\frac{\partial}{\partial z^*}f = q^2dzdz^*\frac{\partial}{\partial z}\frac{\partial}{\partial z^*}f, \\ \bar{\partial}\partial f &= \bar{\partial}\left(dz\frac{\partial f}{\partial z}\right) = -dzdz^*\frac{\partial}{\partial z^*}\frac{\partial}{\partial z}f, & \partial\bar{\partial} + \bar{\partial}\partial &= 0. \end{aligned}$$

We thus obtain the algebra of differential operators in the quantum disc. An example of such operator is a q -analogue of the Laplace operator: $(1 - zz^*)^2 \frac{\partial}{\partial z} \frac{\partial}{\partial z^*}$.

6 Concluding notes

The results described above constitute a background for producing non-commutative function theory in the quantum disc [13]. On the other hand, these results admit an extension onto the case of generic q -Cartan domains introduced in [16].

Applications to non-commutative function theory use essentially the fact that f_0 generate the $U_q \mathfrak{sl}_2$ -module $\mathcal{D}(\mathbb{U})_q$. This allows one to reduce proofs of some relations in algebras of intertwining operators to their verification on f_0 .

A great deal of the algebras we consider in this work were mentioned in the literature (sometimes in different contexts). Nevertheless, most of works on those algebras either do not mention presence of a $U_q \mathfrak{sl}_2$ -module structure or do not use involution and positive invariant integral. That is why we treat the authors of [9, 11] as our direct predecessors.

Our approach to constructing the differential calculus is inspired by the classical result which establishes a duality between the covariant differential operators and morphisms of generalized Verma modules (see [1, §11.1], [6]).

Among other works on quantum algebras, related to this text, one should mention [3, 5, 10, 7].

References

- [1] Baston R., Eastwood M. The Penrose Transform. Its Interaction with Representation Theory. – Clarendon Press, Oxford, 1989.
- [2] V. Chari and A. Pressley. A Guide to Quantum Groups, Cambridge Univ. Press, 1995.
- [3] M. Chaichian, P. P. Kulish. "Quantum group covariant systems", E-print: q-alg/9512017.
- [4] V. G. Drinfeld, "Quantum groups", in *Proceedings of the International Congress of Mathematicians*, Berkeley, 1986, A. M. Gleason (ed.), 1987, American Mathematical Society, Providence, R. I., 798 – 820.
- [5] P. G. A. Floris. "Addition Formula for q -adic Polynomials", Report W94-24, Mathematical Institute, University of Leiden, 10 p.
- [6] M. Harris, H. P. Jakobsen, *Covariant differential operators*, in: Group Theoretical Methods in Physics, Istanbul 1982, Springer Lecture Notes in Physics **180**, Berlin-Heidelberg-New York-Tokyo, 1983.
- [7] H. P. Jakobsen. 'Quantized Hermitian Symmetric Spaces', In "Lie theory and its applications in physics" (Clausthal, 1995), 105 – 116.
- [8] J. C. Jantzen. Lectures on Quantum Groups, Amer. Math. Soc., Providence RI, 1996.
- [9] S. Klimek, A. Lesniewski, "A two-parameter quantum deformation of the unit disc", *J. Funct. Anal.* **115**, (1993), 1 – 23.
- [10] A. Kamita, Y. Morita, and T. Tanisaki. 'Quantum deformations of certain prehomogeneous spaces I', Hiroshima Math. J., **28** (1998), 527 – 540.

- [11] G. Nagy, A. Nica. "On the "quantum disc" and a "non-commutative circle", in: *Algebraic Methods on Operator Theory*, R. E. Curto, P. E. T. Jorgensen (eds.), Birkhauser, Boston, 1994, 276 – 290.
- [12] K. Schmüdgen, A. Schüler, *Covariant differential calculi on quantum spaces and quantum groups*, C. R. Acad. Sci. Paris, Ser I **316** (1993), 1155 – 1160.
- [13] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *On function theory in quantum disc: integral representations*, math.QA/9808015, 17 p.
- [14] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *q-Analogue of the Berezin quantization method*, math.QA/9904173, 9p.
- [15] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum matrix ball: the Bergman kernel*, math.QA/9909036.
- [16] S. Sinel'shchikov and L. Vaksman, *On q-analogues of bounded symmetric domains and Dolbeault complexes*, Math. Phys., Anal., and Geom., **1** (1998), 75 – 100.

QUANTUM DISC: THE CLIFFORD ALGEBRA AND THE DIRAC OPERATOR

K. Schmüdgen

S. Sinel'shchikov

L. Vaksman

1 Introduction

In his non-commutative geometry A. Connes [1] has developed methods and notions in order to define and to study "non-commutative Riemannian manifolds". A crucial role in this theory is played by the Dirac operators and its non-commutative analogs.

One of the simplest (non-compact) Riemannian manifolds of a constant curvature is the unit disc $\mathbb{U} \subset \mathbb{C}$ whose metric is given by

$$(1 - x^2 - y^2)^{-2} dx dy$$

(the Poincaré model of Lobachevskian geometry). A non-commutative analogue of this Riemannian manifold was extensively studied recently within the framework of the quantum group theory [5]. The aim of the present paper is to develop the Clifford algebra and the Dirac operator for this quantum disc.

2 A q -Clifford bundle

In what follows $q \in (0, 1)$ and $\mathbb{C}$ is the ground field. We assume some acquaintance with the basics of quantum group theory [5, 3]. All algebras in this section are supposed to be unital.

We use the standard notation for the quantum universal enveloping algebra $U_q \mathfrak{sl}_2$, its generators E, F, K, K^{-1} , the comultiplication Δ , the counit ε , and the antipode S [3, 8]. Throughout we shall use the Sweedler notation $\Delta(f) = f_{(1)} \otimes f_{(2)}$ instead of $\Delta(f) = \sum_i f'_i \otimes f''_i$. The $*$ -Hopf algebra $U_q \mathfrak{su}_{1,1} = (U_q \mathfrak{sl}_2, *)$ is defined by

$$E^* = -KF, \quad F^* = -EK^{-1}, \quad (K^{\pm 1})^* = K^{\pm 1}.$$

Primarily we are interested in $U_q \mathfrak{sl}_2$ -module algebras and $U_q \mathfrak{su}_{1,1}$ -module $*$ -algebras¹. We follow [8] in starting with the $*$ -algebra $\text{Pol}(\mathbb{C})_q$ with single generator z and defining relation

$$z^* z = q^2 z z^* + 1 - q^2. \quad (2.1)$$

The structure of the $U_q \mathfrak{su}_{1,1}$ -module algebra used by the authors in [8] attracted an attention about 10 years ago [4]. It can be determined by

$$Fz = q^{1/2}, \quad Ez = -q^{1/2} z^2, \quad K^{\pm 1} z = q^{\pm 2} z. \quad (2.2)$$

This research was supported in part by of the US Civilian Research & Development Foundation (Award No UM1-2091) and by Swedish Academy of Sciences (project No 11293562).

¹A $U_q \mathfrak{sl}_2$ -module $*$ -algebra F is called $U_q \mathfrak{su}_{1,1}$ -module algebra if the following compatibility condition on involutions is valid:

$$(\xi f)^* = (S(\xi))^* f^*, \quad \xi \in U_q \mathfrak{su}_{1,1}, \quad f \in F.$$

We follow [8] in recalling the description of a covariant differential calculus over the $U_q\mathfrak{su}_{1,1}$ -module algebra $\text{Pol}(\mathbb{C})_q$ which initially appeared in [7]. This is a $U_q\mathfrak{su}_{1,1}$ -module algebra $\Omega_q \supset \text{Pol}(\mathbb{C})_q$ equipped with a (super)differentiation $d : \Omega_q \rightarrow \Omega_q$ which commutes with an involution $*$ and is a morphism of $U_q\mathfrak{su}_{1,1}$ -modules.

The list of relations which determine the $U_q\mathfrak{su}_{1,1}$ -module algebra Ω_q consists of the above relations determining the $U_q\mathfrak{su}_{1,1}$ -module algebra $\text{Pol}(\mathbb{C})_q$, and the additional relations

$$dz \cdot z = q^2 z \cdot dz, \quad dz \cdot z^* = q^{-2} z^* \cdot dz, \quad (2.3)$$

$$dz \cdot dz = 0, \quad dz^* \cdot dz^* = 0, \quad (2.4)$$

$$dz^* \cdot dz + q^2 dz \cdot dz^* = 0. \quad (2.5)$$

In the classical case $q = 1$ the passage from the algebra of exterior differential forms with polynomial coefficients to the algebra of polynomial sections of the Clifford bundle (associated to an invariant metric) reduces to a replacement of the relations

$$d\bar{z}dz + dzd\bar{z} = 0 \quad \left(\Leftrightarrow \frac{d\bar{z}dz + dzd\bar{z}}{(1 - |z|^2)^2} = 0 \right)$$

and

$$d\bar{z}dz + dzd\bar{z} = (1 - |z|^2)^2 \quad \left(\Leftrightarrow \frac{d\bar{z}dz + dzd\bar{z}}{(1 - |z|^2)^2} = 1 \right).$$

A similar argument for $q \in (0, 1)$ leads to the following definition of the 'algebra of polynomial sections' of a q -Clifford bundle.

DEFINITION. Let Cl_q denote the $*$ -algebra with generators z , dz and determining relations (2.1), (2.3), (2.4), and

$$dz^* \cdot dz + q^2 dz \cdot dz^* = (1 - zz^*)^2. \quad (2.6)$$

We equip Cl_q with the standard filtration given by $\deg(z) = \deg(z^*) = 0$, $\deg(dz) = \deg(dz^*) = 1$.

It follows from the definition of Cl_q that, just as in the classical case $q = 1$, one has

Proposition 2.1 *The graded algebra of Cl_q associated to its standard filtration is isomorphic to Ω_q .*

Consider the $U_q\mathfrak{sl}_2$ -module subalgebras

$$\begin{aligned} \Omega_q^{(*,0)} &= \{f + dz \cdot g \mid f, g \in \text{Pol}(\mathbb{C})_q\} \subset \Omega_q, \\ \Omega_q^{(0,*)} &= \{f + g \cdot dz^* \mid f, g \in \text{Pol}(\mathbb{C})_q\} \subset \Omega_q. \end{aligned}$$

As a consequence of the definition of Cl_q one has

$$\Omega_q^{(*,0)} \hookrightarrow \text{Cl}_q, \quad \Omega_q^{(0,*)} \hookrightarrow \text{Cl}_q. \quad (2.7)$$

Proposition 2.2 *There exists a unique structure of $U_q\mathfrak{su}_{1,1}$ -module algebra in Cl_q such that the embeddings (2.7) appear to be morphisms of $U_q\mathfrak{sl}_2$ -modules.*

Proof. The only non-trivial statement here is about the existence of a structure of $U_q\mathfrak{sl}_2$ -module algebra in Cl_q such that the embeddings (2.7) are morphisms of $U_q\mathfrak{sl}_2$ -modules.

Consider the $*$ -algebra $\mathcal{F}$ with generators z, dz and determining relations (2.1), (2.3), (2.4). $\mathcal{F}$ is a $U_q\mathfrak{su}_{1,1}$ -module algebra, and the canonical homomorphism $\mathcal{F} \rightarrow \Omega_q$ is a morphism of $U_q\mathfrak{su}_{1,1}$ -module algebras. One has to prove that the two-sided ideal J of $\mathcal{F}$ generated by $dz^* \cdot dz + q^2 dz \cdot dz^* - (1 - zz^*)^2$, is a submodule of the $U_q\mathfrak{su}_{1,1}$ -module $\mathcal{F}$. Obviously, $K^{\pm 1}J \subset J$, one needs only to prove that $EJ \subset J$, $FJ \subset J$. The proofs of these inclusions are similar. We restrict ourselves to demonstrating the first of them. It follows from the following statement.

Lemma 2.3

- i) $E(dzdz^*) = -q^{1/2} (1 + q^2) z dz dz^*$,
- ii) $E(dz^* dz) = -q^{1/2} (1 + q^2) z dz^* dz$,
- iii) $E((1 - zz^*)^2) = -q^{1/2} (1 + q^2) z(1 - zz^*)^2$.

Proof of lemma 2.3.

$$\begin{aligned} i) \ E(dzdz^*) &= (Edz)dz^* = d(Ez)dz^* = d(-q^{1/2}z^2)dz^* = -q^{1/2}(dz \cdot z + z \cdot dz)dz^* = \\ &= -q^{1/2} (1 + q^2) z dz dz^*. \end{aligned}$$

$$ii) \ E(dz^* dz) = (Kdz^*)(Edz) = q^{-2}dz^* (-q^{1/2} (1 + q^2) z dz) = -q^{1/2} (1 + q^2) z dz^* dz.$$

iii) Note first that $Ez^* = q^{-3/2}$. (In fact, $(Fz)^* = (S(F))^* z^*$. Hence $q^{1/2} = (-FK)^* z^* = -K^* (-EK^{-1}) z^* = q^2 Ez^*$). Furthermore,

$$E(1 - zz^*) = q^{1/2} zz^* - q^2 z q^{-3/2} = -q^{1/2} z(1 - zz^*).$$

Finally,

$$E((1 - zz^*)^2) = -q^{1/2} z(1 - zz^*)^2 - q^{1/2} (1 - zz^*) z(1 - zz^*) = -q^{1/2} (1 + q^2) z(1 - zz^*)^2. \quad \square$$

REMARK. Propositions 2.1 and 2.2 suggest that an invariant Riemannian metric in the quantum disc should be defined by an expression of the form

$$(1 - zz^*)^{-2} (dz^* \otimes dz + q^2 dz \otimes dz^*). \quad (2.8)$$

To begin with, consider the multiplicative closed subset $(1 - zz^*)^{\mathbb{N}} \subset \text{Pol}(\mathbb{C})_q \subset \Omega_q$. It follows from the relations

$$\begin{aligned} (1 - zz^*)z &= q^2 z(1 - zz^*), & (1 - zz^*)z^* &= q^{-2} z^*(1 - zz^*), \\ (1 - zz^*)dz &= dz(1 - zz^*), & (1 - zz^*)dz^* &= dz^*(1 - zz^*), \end{aligned}$$

that this multiplicative closed system is an Ore set. Consider the associated localizations $\widetilde{\text{Pol}}(\mathbb{C})_q$ and $\widetilde{\Omega}_q$ of $\text{Pol}(\mathbb{C})_q$ and Ω_q , respectively. It is well-known that there embeddings $\text{Pol}(\mathbb{C})_q \hookrightarrow \widetilde{\text{Pol}}(\mathbb{C})_q$ and $\Omega_q \hookrightarrow \widetilde{\Omega}_q$. One can also prove just as in [12] that the structure of $U_q\mathfrak{su}_{1,1}$ -module algebra is uniquely extended from Ω_q onto $\widetilde{\Omega}_q$.

Certainly, the grading mentioned above is canonically extendable onto $\widetilde{\Omega}_q$ and

$$\widetilde{\Omega}_q^1 = \left\{ f' dz + f'' dz^* \mid f', f'' \in \widetilde{\text{Pol}}(\mathbb{C})_q \right\}$$

is a $U_q\mathfrak{sl}_2$ -module $\widetilde{\text{Pol}}(\mathbb{C})_q$ -bimodule, equipped with the involution $*$ inherited from $\widetilde{\Omega}_q$.

Now we are in a position to introduce the $U_q\mathfrak{sl}_2$ -module $\widetilde{\Omega}_q \otimes_{\widetilde{\text{Pol}}(\mathbb{C})_q} \widetilde{\Omega}_q$, with a 'metric tensor' (2.8) chosen among the $U_q\mathfrak{sl}_2$ -invariant elements of this module.

3 A q-Dirac bundle

Consider the $U_q\mathfrak{sl}_2$ -module algebra of 'differential forms' $\Omega_q^{(0,*)}$, together with its $U_q\mathfrak{sl}_2$ -module subalgebra $\text{Pol}(\mathbb{C})_q$. Obviously, $\Omega_q^{(0,*)}$ is a free (left) $\text{Pol}(\mathbb{C})_q$ -module with generators $1, dz^*$. Of course, $\Omega_q^{(0,*)}$ is a $U_q\mathfrak{sl}_2$ -module $\text{Pol}(\mathbb{C})_q$ -module (a q-analogue of the space of polynomial sections of the vector bundle $T^{(0,*)}\mathbb{C}$).

In the classical case ($q = 1$) the Clifford algebra acts in the fibers of $T^{(0,*)}\mathbb{C}$, which allows us to speak about the Dirac bundle. We are going to describe a quantum analogue of this action. It follows from the definition of Cl_q that, in the category of $U_q\mathfrak{sl}_2$ -module $\text{Pol}(\mathbb{C})_q$ -modules there exists a canonical isomorphism

$$\Omega_q^{(0,*)} \simeq \text{Cl}_q / \text{Cl}_q \cdot dz.$$

In particular, $\Omega_q^{(0,*)}$ is a $U_q\mathfrak{sl}_2$ -module Cl_q -module.

Consider a sesquilinear map $\mathbf{h} : \Omega_q^{(0,*)} \times \Omega_q^{(0,*)} \rightarrow \text{Pol}(\mathbb{C})_q$,

$$\mathbf{h}(\varphi_0 + dz^*\varphi_1, \psi_0 + dz^*\psi_1) = \psi_0^*\varphi_0 + q^{-2}\psi_1^*(1 - zz^*)^2\varphi_1,$$

for $\varphi_0, \varphi_1, \psi_0, \psi_1 \in \text{Pol}(\mathbb{C})_q$.

We are going to demonstrate that the 'Hermitian metric' $\mathbf{h}$ is positive definite, invariant and 'respects' the Cl_q -action. That is, it can be used to produce a q-analogue of the Dirac operator.

Proposition 3.1

- i) $\mathbf{h}(\omega, \omega) \geq 0$ for all $\omega \in \Omega_q^{(0,*)}$ and $\mathbf{h}(\omega, \omega) = 0 \Rightarrow \omega = 0$,
- ii) $\xi \mathbf{h}(\omega_1, \omega_2) = \mathbf{h}(\xi_{(2)}(\omega_1), (S(\xi_{(1)}))^*\omega_2)$ for all $\xi \in U_q\mathfrak{su}_{1,1}$, $\omega_1, \omega_2 \in \Omega_q^{(0,*)}$.²
- iii) $\mathbf{h}(c\omega_1, \omega_2) = \mathbf{h}(\omega_1, c^*\omega_2)$ for all $c \in \text{Cl}_q$, $\omega_1, \omega_2 \in \Omega_q^{(0,*)}$. (3.1)

Proof. The first statement is evident. To prove the second statement, it suffices to establish $U_q\mathfrak{su}_{1,1}$ -invariance of the Hermitian forms

$$\begin{aligned} \mathbf{h}_1 : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q &\rightarrow \text{Pol}(\mathbb{C})_q, & \mathbf{h}_1 : f_1 \times f_2 &\mapsto f_2^*f_1, \\ \mathbf{h}_2 : \Omega_q^{(0,1)} \times \Omega_q^{(0,1)} &\rightarrow \text{Pol}(\mathbb{C})_q, & \mathbf{h}_2 : dz^*f_1 \times dz^*f_2 &\mapsto f_2^*(1 - zz^*)^2f_1. \end{aligned}$$

On the other hand, since Ω_q is a $U_q\mathfrak{su}_{1,1}$ -module algebra, the forms $\mathbf{h}_1$ and

$$\mathbf{h}_3 : \Omega_q^{(0,1)} \times \Omega_q^{(0,1)} \rightarrow \Omega_q^{(1,1)}, \quad \mathbf{h}_3 : \omega \times \omega \mapsto \omega^*\omega$$

are $U_q\mathfrak{su}_{1,1}$ -invariant. (Here $\Omega_q^{(1,1)} = \{f dz dz^* \mid f \in \text{Pol}(\mathbb{C})_q\} \subset \Omega_q$.) The $U_q\mathfrak{su}_{1,1}$ -invariance of $\mathbf{h}_2$ follows from the invariance of $\mathbf{h}_3$ and the invariance of the element

$$(1 - zz^*)^{-2} dz dz^* = dz dz^* (1 - zz^*)^{-2} \in \tilde{\Omega}_q. \quad (3.2)$$

²Recall that we use the Sweedler notation. The condition ii) is just a rephrasing of a $U_q\mathfrak{su}_{1,1}$ -invariance of the Hermitian metric $\mathbf{h}$. The flip of tensor multiples $\xi_{(1)}, \xi_{(2)}$ in ii) is due to the fact that a similar flip is normally used implicitly while constructing scalar products in the spaces of functions and differential forms. For example, for functions

$$f_1 \otimes f_2 \mapsto f_2 \otimes f_1 \mapsto f_2^* \otimes f_1 \mapsto f_2^* f_1 \mapsto \int f_2^* f_1 d\nu.$$

One can use a similar argument to prove (3.1) in the special case $c \in \text{Pol}(\mathbb{C})_q$. (The passage from $\mathbf{h}_3$ to $\mathbf{h}_2$ involves the fact that the 'Kähler form' (3.2) is in the center of $\widetilde{\Omega}_q$.)

We prove (3.1) in the general case. In view of the relations

$$\mathbf{h}(\omega_1 f, \omega_2) = \mathbf{h}(\omega_1, \omega_2) f, \quad \mathbf{h}(\omega_1, \omega_2 f) = f^* \mathbf{h}(\omega_1, \omega_2), \quad \omega_1, \omega_2 \in \Omega_q^{(0,*)}, \quad f \in \text{Pol}(\mathbb{C})_q,$$

it remains only to consider the case $\omega_1, \omega_2 \in \{1, dz^*\}$, $c = dz^*$. In this case (3.1) follows from

$$\begin{aligned} \mathbf{h}(dz dz^*, 1) &= \mathbf{h}(1, dz dz^*) = q^{-2}(1 - zz^*)^2, \\ \mathbf{h}(dz^*, dz^*) &= q^{-2}(1 - zz^*)^2. \end{aligned}$$

The first relation here follows from the fact that in the Cl_q -module $\Omega_q^{(0,*)} \simeq \text{Cl}_q / \text{Cl}_q \cdot dz$

$$dz \cdot dz^* = q^{-2}(1 - zz^*)^2,$$

while the second one is obvious. □

4 An important *-representation of Cl_q

Every positive linear functional $\mu : \text{Pol}(\mathbb{C})_q \rightarrow \mathbb{C}$ determines a *-representation of Cl_q in the pre-Hilbert space $\Omega_q^{(0,*)}$ with scalar product

$$(\omega_1, \omega_2) = \int \mathbf{h}(\omega_1, \omega_2) d\mu. \quad (4.1)$$

However, to produce a Dirac operator we need a $U_q \mathfrak{su}_{1,1}$ -invariant scalar product. We first recall the corresponding definition.

Consider a $U_q \mathfrak{su}_{1,1}$ -module V and a sesquilinear form $V \times V \rightarrow \mathbb{C}$, $v_1 \times v_2 \mapsto (v_1, v_2)$. This sesquilinear form is called $U_q \mathfrak{su}_{1,1}$ -invariant if for all $\xi \in U_q \mathfrak{su}_{1,1}$, $v_1, v_2 \in V$

$$\varepsilon(\xi) \cdot (v_1, v_2) = (\xi_{(2)} v_1, (S(\xi_{(1)}))^* v_2).$$

The invariance of a sesquilinear map $\mathbf{h}$ indicates a naive way of producing the required scalar product. That is, formula (4.1) suggests to define a $U_q \mathfrak{su}_{1,1}$ -invariant integral μ by

$$\int (\xi f) d\mu = \varepsilon(\xi) \int f d\mu, \quad \xi \in U_q \mathfrak{su}_{1,1}, \quad f \in \text{Pol}(\mathbb{C})_q.$$

In fact this fails even in the classical case ($q = 1$), since the positive $SU_{1,1}$ -invariant measure in the unit disc has the form

$$d\nu = \text{const} (1 - |z|^2)^{-2} dz d\bar{z},$$

and the associated integral is not defined on the polynomial algebra (unless $\text{const} = 0$). That is why in producing an integral calculus, the algebra $\mathcal{D}(\mathbb{U})$ of smooth functions with support inside the disc $\mathbb{U}$ is used. In the quantum case one can also replace the $U_q \mathfrak{su}_{1,1}$ -module algebra $\text{Pol}(\mathbb{C})_q$ by the $U_q \mathfrak{su}_{1,1}$ -module algebra $\mathcal{D}(\mathbb{U})_q$ to produce a $U_q \mathfrak{su}_{1,1}$ -invariant integral $\nu : \mathcal{D}(\mathbb{U})_q \rightarrow \mathbb{C}$ (see [8]).

Recall the definition of the $U_q \mathfrak{su}_{1,1}$ -module algebra $\mathcal{D}(\mathbb{U})_q$. Consider the *-algebra $\text{Fun}(\mathbb{U})_q$ with generators z, f_0 and determining relations (2.1) and

$$z^* f_0 = f_0 z = 0, \quad f_0^2 = f_0, \quad f_0^* = f_0. \quad (4.2)$$

Obviously, we have an embedding $\text{Pol}(\mathbb{C})_q \hookrightarrow \text{Fun}(\mathbb{U})_q$.

An argument used in [8] leads to (4.2) and to the relations

$$E f_0 = -\frac{q^{1/2}}{1-q^2} z f_0, \quad F f_0 = -\frac{q^{1/2}}{q^{-2}-1} f_0 z^*, \quad K^{\pm 1} f_0 = f_0. \quad (4.3)$$

This argument also allows us to prove that the structure of $U_q \mathfrak{su}_{1,1}$ -module algebra admits an extension from $\text{Pol}(\mathbb{C})_q$ to $\text{Fun}(\mathbb{U})_q$ which satisfies (4.3). The uniqueness of this extension is obvious. The $U_q \mathfrak{su}_{1,1}$ -module algebra $\mathcal{D}(\mathbb{U})_q$ is defined as a two-sided ideal of $\text{Fun}(\mathbb{U})_q$ generated by f_0 :

$$\mathcal{D}(\mathbb{U})_q = \text{Fun}(\mathbb{U})_q \cdot f_0 \cdot \text{Fun}(\mathbb{U})_q.$$

Recall the explicit form of the invariant integral ν .

Let Q be the linear operator in $l^2(\mathbb{Z}_+)$ given by $Q e_k = (1 - q^2) q^{-2k} e_k$, with $\{e_k\}$ being the standard basis in $l^2(\mathbb{Z}_+)$. Consider the $*$ -representation T of $\text{Fun}(\mathbb{U})_q$ in the Hilbert space $l^2(\mathbb{Z}_+)$ given by

$$T(z) e_k = (1 - q^{2(k+1)})^{1/2} e_{k+1}, \quad T(z^*) e_k = \begin{cases} (1 - q^{2k})^{1/2} e_{k-1}, & k > 0, \\ 0, & k = 0. \end{cases},$$

$$T(f_0) e_k = \begin{cases} e_k, & k = 0, \\ 0, & k \neq 0. \end{cases}$$

Note that the only non-zero matrix element of the operator $T(z^j f_0 z^{*k})$ is in line j column k . Hence the operators $T(z^j f_0 z^{*k})$ are linear independent. From this one deduces that the elements $z^j f_0 z^{*k}$, $j, k \in \mathbb{Z}_+$, form a basis in $\mathcal{D}(\mathbb{U})_q$ and that the representation T of $\mathcal{D}(\mathbb{U})_q$ is faithful.

It is easy to prove (see [9]):

Proposition 4.1 *The linear functional*

$$\int_{\mathbb{U}_q} f d\nu \stackrel{\text{def}}{=} \text{tr}(T(f)Q), \quad f \in \mathcal{D}(\mathbb{U})_q,$$

on the $U_q \mathfrak{su}_{1,1}$ -module $$ -algebra $\mathcal{D}(\mathbb{U})_q$ is well defined, positive definite, and $U_q \mathfrak{su}_{1,1}$ -invariant.*

We have thus derived a $U_q \mathfrak{su}_{1,1}$ -module algebra $\text{Fun}(\mathbb{U})_q$ from $\text{Pol}(\mathbb{C})_q$ by adding a generator and additional relations (4.2), (4.3).

In a similar way one can extend the $U_q \mathfrak{su}_{1,1}$ -module $*$ -algebras Ω_q and Cl_q (see [8]). For that, one has to add f_0 to the list of generators and to complete the list of relations with (4.2), (4.3) and

$$f_0 dz = dz \cdot f_0, \quad f_0 \cdot dz^* = dz^* \cdot f_0. \quad (4.4)$$

Let $\Omega(\mathbb{U})_q$, $\text{Cl}(\mathbb{U})_q$ be the two-sided ideals of those algebras generated by f_0 . We call $\Omega(\mathbb{U})_q$ the algebra of differential forms with finite coefficients in the quantum disc, and $\text{Cl}(\mathbb{U})_q$ the algebra of finite sections of the q -Clifford bundle.

Obviously, $\Omega(\mathbb{U})_q$ is a $U_q\mathfrak{su}_{1,1}$ -module Ω_q -bimodule and $\text{Cl}(\mathbb{U})_q$ is a $U_q\mathfrak{su}_{1,1}$ -module Cl_q -bimodule. As usual,

$$\begin{aligned}\Omega(\mathbb{U})_q^{(0,1)} &= \{f + dz^*g \mid f, g \in \mathcal{D}(\mathbb{U})_q\}, \\ \Omega(\mathbb{U})_q^{(0,1)} &= \text{Cl}(\mathbb{U})_q / \text{Cl}(\mathbb{U})_q \cdot dz.\end{aligned}\tag{4.5}$$

It follows from (4.5) that $\Omega(\mathbb{U})_q^{(0,1)}$ is a Cl_q -module. We intend to equip $\Omega(\mathbb{U})_q^{(0,1)}$ with an invariant Hermitian metric and an invariant scalar product.

Consider a sesquilinear map $\mathbf{h}_{\mathbb{U}} : \Omega(\mathbb{U})_q^{(0,*)} \times \Omega(\mathbb{U})_q^{(0,*)} \rightarrow \mathcal{D}(\mathbb{U})_q$,

$$\mathbf{h}_{\mathbb{U}}(\varphi_0 + dz^* \cdot \varphi_1, \psi_0 + dz^* \cdot \psi_1) = \psi_0^* \varphi_0 + q^{-2} \psi_1^* (1 - zz^*)^2 \varphi_1, \tag{4.6}$$

with $\varphi_0, \varphi_1, \psi_0, \psi_1 \in \mathcal{D}(\mathbb{U})_q$.

Proposition 4.2

- i) $\mathbf{h}_{\mathbb{U}}(\omega, \omega) \geq 0$ for all $\omega \in \Omega(\mathbb{U})_q^{(0,*)}$ and $\mathbf{h}_{\mathbb{U}}(\omega, \omega) = 0 \Rightarrow \omega = 0$,
- ii) $\xi \mathbf{h}_{\mathbb{U}}(\omega_1, \omega_2) = \mathbf{h}_{\mathbb{U}}(\xi_{(2)} \omega_1, (S(\xi_{(1)}))^* \omega_2)$ for all $\xi \in U_q\mathfrak{su}_{1,1}$, $\omega_1, \omega_2 \in \Omega(\mathbb{U})_q^{(0,*)}$.
- iii) $\mathbf{h}_{\mathbb{U}}(c\omega_1, \omega_2) = \mathbf{h}_{\mathbb{U}}(\omega_1, c^* \omega_2)$ for all $c \in \text{Cl}_q$, $\omega_1, \omega_2 \in \Omega(\mathbb{U})_q^{(0,*)}$.

This proposition can be proved in a similar manner as proposition 3.1.

Propositions 4.1 and 4.2 allow us to equip $\Omega(\mathbb{U})_q^{(0,*)}$ with a $U_q\mathfrak{su}_{1,1}$ -invariant scalar product: $(\omega_1, \omega_2) \stackrel{\text{def}}{=} \int_{\mathbb{U}_q} \mathbf{h}_{\mathbb{U}}(\omega_1, \omega_2) d\nu$.

Consider the Hilbert space $\mathcal{H}_q$ which is the completion of the pre-Hilbert space $\Omega(\mathbb{U})_q^{(0,*)}$.

The boundedness of the operators $T(f)$, $f \in \text{Pol}(\mathbb{C})_q$, and the explicit form of the invariant integral $\nu : \mathcal{D}(\mathbb{U})_q \rightarrow \mathbb{C}$ imply the following

Proposition 4.3 *The linear operators $\Omega(\mathbb{U})_q^{(0,*)} \rightarrow \Omega(\mathbb{U})_q^{(0,*)}$, $\omega \mapsto c\omega$, are bounded for all $c \in \text{Cl}_q$.*

Corollary 4.4 *There exists a unique $*$ -representation π of Cl_q (by bounded operators) in $\mathcal{H}_q$ such that for all $c \in \text{Cl}_q$, $\omega \in \Omega(\mathbb{U})_q^{(0,*)}$,*

$$\pi(c)\omega = c\omega.$$

We close this section by recalling that the structure of $U_q\mathfrak{su}_{1,1}$ -module (super)differential algebra can be canonically transferred from Ω_q onto $\Omega(\mathbb{U})_q^{(0,*)}$ (see [8]):

$$df_0 = -\frac{1}{1-q^2}(dz \cdot f_0 \cdot z^* + z \cdot f_0 \cdot dz^*).$$

5 The Dirac operator in the quantum disc

The Dirac operator is normally defined in terms of special affine connections. Nevertheless, in the case of Kähler manifold, this operator admits a definition by a simple explicit formula (see, for example, [6]). In the special case of the unit disc $\mathbb{U} \subset \mathbb{C}$ and the Clifford bundle as it is considered in this work, the Dirac operator in $\Omega(\mathbb{U})^{(0,*)} = \Omega(\mathbb{U})^{(0,0)} + \Omega(\mathbb{U})^{(0,1)}$ is defined by

$$\mathcal{D} = \begin{pmatrix} 0 & (\bar{\partial})^* \\ \bar{\partial} & 0 \end{pmatrix}, \quad (5.1)$$

where $(\bar{\partial})^*$ being a formal adjoint differential operator (with respect to the standard $SU_{1,1}$ -invariant scalar products in $\Omega(\mathbb{U})^{(0,0)}$, $\Omega(\mathbb{U})^{(0,1)}$). The operator $\mathcal{D}$ is essentially selfadjoint in the Hilbert space completion $\mathcal{H}$ of $\Omega(\mathbb{U})^{(0,*)}$.

It is worth to note that

$$\mathcal{D}^2 = \begin{pmatrix} \square_0 & 0 \\ 0 & \square_1 \end{pmatrix},$$

with $\square_0 = (\bar{\partial})^* \bar{\partial}$, $\square_1 = \bar{\partial}(\bar{\partial})^*$ being essentially selfadjoint $SU_{1,1}$ -invariant differential operators (invariant Laplace operators in the spaces of functions and differential type $(0, 1)$ forms, respectively).

Now we are going to pass from the classical disc to the quantum disc. Just as in the classical case, one has the decompositions

$$\Omega(\mathbb{U})_q = \bigoplus_{i,j=0}^1 \Omega(\mathbb{U})_q^{(i,j)}, \quad d = \partial + \bar{\partial},$$

with

$$\begin{aligned} \partial &: \Omega(\mathbb{U})_q^{(i,j)} \rightarrow \Omega(\mathbb{U})_q^{(i+1,j)}, \\ \bar{\partial} &: \Omega(\mathbb{U})_q^{(i,j)} \rightarrow \Omega(\mathbb{U})_q^{(i,j+1)} \end{aligned}$$

(these decompositions are related to the standard bigrading of the algebra of differential forms: $\deg(z) = \deg(z^*) = \deg(f_0) = (0, 0)$, $\deg(dz) = (1, 0)$, $\deg(dz^*) = (0, 1)$).

In some sense the quantum case turns out to be easier than the classical one, because we have

Proposition 5.1 *The linear operator $\bar{\partial} : \Omega(\mathbb{U})_q^{(0,0)} \rightarrow \Omega(\mathbb{U})_q^{(0,1)}$ is bounded.*

See a proof in [10].

Let $\mathcal{H}_q^{(0)}$ and $\mathcal{H}_q^{(1)}$ be the closures of the linear manifolds $\Omega(\mathbb{U})_q^{(0,0)} = \mathcal{D}(\mathbb{U})_q$ and $\Omega(\mathbb{U})_q^{(0,1)}$, respectively, in $\mathcal{H}_q$. From proposition 5.1 it follows that the linear map $\bar{\partial} : \Omega(\mathbb{U})_q^{(0,0)} \rightarrow \Omega(\mathbb{U})_q^{(0,1)}$ has an extension by a continuity up to a bounded operator $\bar{\partial} : \mathcal{H}_q^{(0)} \rightarrow \mathcal{H}_q^{(1)}$. Now the decomposition $\mathcal{H}_q = \mathcal{H}_q^{(0)} \oplus \mathcal{H}_q^{(1)}$ allows us to introduce the Dirac operator in the quantum disc by (5.1). As we have shown, in the quantum case the operator $\mathcal{D}$ is bounded in $\mathcal{H}_q$.

Note that $\mathcal{D} = \mathcal{D}^*$, and that the Dirac operator is an endomorphism of the $U_q \mathfrak{su}_{1,1}$ -module $\Omega(\mathbb{U})_q^{(0,*)}$, by this property of the operator $\bar{\partial}$ and the $U_q \mathfrak{su}_{1,1}$ -invariance of the scalar products.

Introduce the standard notation $\frac{\partial}{\partial z}, \frac{\partial}{\partial z^*}$ for the linear operators defined by

$$\partial\psi = dz \frac{\partial\psi}{\partial z}, \quad \bar{\partial}\psi = dz^* \frac{\partial\psi}{\partial z^*}.$$

Proposition 5.2 *For all $\psi \in \mathcal{D}(\mathbb{U})_q$*

$$(\bar{\partial})^* : dz^* \psi \mapsto q^2(1 - zz^*)^2 \frac{\partial\psi}{\partial z}.$$

Proof. It suffices to prove that the linear operators $q^2 \frac{\partial}{\partial z}, \frac{\partial}{\partial z^*}$ are formal adjoints with respect to the scalar product in $\mathcal{D}(\mathbb{U})_q$

$$\varphi \times \psi \mapsto \int_{\mathbb{U}_q} \psi^* \varphi (1 - zz^*)^2 d\nu,$$

determined by the 'q-Lebesgue measure' $(1 - zz^*)^2 d\nu$. The latter assertion can be proved in the same way as the corresponding statement in [11].

Corollary 5.3 *For all $\varphi, \psi \in \mathcal{D}(\mathbb{U})_q$*

$$\begin{aligned} \mathcal{P}(\varphi + dz^* \psi) &= q^2(1 - zz^*)^2 \frac{\partial\psi}{\partial z} + dz^* \frac{\partial\varphi}{\partial z^*}, \\ (\bar{\partial})^* \bar{\partial}\varphi &= q^2(1 - zz^*)^2 \frac{\partial}{\partial z} \frac{\partial}{\partial z^*}, \\ \bar{\partial}(\bar{\partial})^*(dz^* \psi) &= q^2 dz^* \frac{\partial}{\partial z^*} (1 - zz^*)^2 \frac{\partial}{\partial z^*}. \end{aligned}$$

6 Concluding remarks

In A. Connes' non-commutative Riemannian geometry [1, chapter VI] the differential d is closely related to the Dirac operator $\mathcal{D}$. Specifically, for any 'function' f , the elements f and df are in the Clifford algebra. They act in the same space as $\mathcal{D}$, and

$$df \cdot \psi = [\mathcal{D}, f] \cdot \psi. \quad (6.1)$$

A similar relation is also valid in the case of quantum disc as one can see from the following obvious

Proposition 6.1 *For all $f \in \text{Pol}(\mathbb{C})_q, \psi \in \mathcal{D}(\mathbb{U})_q$*

$$(\bar{\partial}f) \cdot \psi = [\mathcal{D}, f] \cdot \psi. \quad (6.2)$$

Of course, one can replace the Cl_q -module $\Omega_q^{(0,*)}$ by Cl_q treated as a Cl_q -module, in order to pass from $\bar{\partial}$ to d , i.e. for getting the relation (6.1) instead of (6.2). This is just what stands for the Dirac bundle in [6, chapter 5A].

There exists a unique Hermitian metric $\mathbf{h}$ which has all the properties stated in proposition 3.1 and satisfies the relation $\mathbf{h}(dz, dz) = \mathbf{h}(dz^*, dz^*)$:

$$\begin{aligned} & \mathbf{h}(\varphi_{00} + dz\varphi_{10} + dz^*\varphi_{01} + dzdz^*\varphi_{11}, \psi_{00} + dz\psi_{10} + dz^*\psi_{01} + dzdz^*\psi_{11}) = \\ & = \psi_{00}^*\varphi_{00} + \frac{1}{1+q^2} (\psi_{10}^*(1-zz^*)^2\varphi_{10} + \psi_{01}^*(1-zz^*)^2\varphi_{01}) + \frac{1}{q^2(1+q^2)} \psi_{11}^*(1-zz^*)^4\varphi_{11}, \end{aligned}$$

with $\varphi_{ij}, \psi_{ij} \in \text{Pol}(\mathbb{C})_q$.

REMARK. The $U_q\mathfrak{su}_{1,1}$ -invariance of the Hermitian metric $\mathbf{h}$ follows from the $U_q\mathfrak{su}_{1,1}$ -invariance of the following elements:

$$\begin{aligned} (1-zz^*)^{-2}dzdz^* & \in \widetilde{\Omega}_q,^3 \\ (1-zz^*)^{-2}dz \otimes dz^* & \in \widetilde{\Omega}_q^{(1,0)} \otimes_{\text{Pol}(\mathbb{C})_q} \widetilde{\Omega}_q^{(0,1)}, \\ (1-zz^*)^{-2}dz^* \otimes dz & \in \widetilde{\Omega}_q^{(0,1)} \otimes_{\text{Pol}(\mathbb{C})_q} \widetilde{\Omega}_q^{(1,0)}. \end{aligned}$$

In fact, one can use the two latter elements in order to define the following morphisms of $U_q\mathfrak{su}_{1,1}$ -modules:

$$\begin{aligned} \widetilde{\Omega}_q^{(0,1)} \otimes_{\widetilde{\text{Pol}(\mathbb{C})_q}} \widetilde{\Omega}_q^{(1,0)} & \rightarrow \widetilde{\text{Pol}(\mathbb{C})_q}, & \psi dz^* \otimes dz \varphi & \mapsto \psi(1-zz^*)^2\varphi; \\ \widetilde{\Omega}_q^{(1,0)} \otimes_{\widetilde{\text{Pol}(\mathbb{C})_q}} \widetilde{\Omega}_q^{(0,1)} & \rightarrow \widetilde{\text{Pol}(\mathbb{C})_q}, & \psi dz \otimes dz^* \varphi & \mapsto \psi(1-zz^*)^2\varphi; \\ \widetilde{\Omega}_q^{(1,0)} \otimes_{\widetilde{\text{Pol}(\mathbb{C})_q}} \widetilde{\Omega}_q^{(0,1)} \otimes_{\widetilde{\text{Pol}(\mathbb{C})_q}} \widetilde{\Omega}_q^{(1,0)} \otimes_{\widetilde{\text{Pol}(\mathbb{C})_q}} \widetilde{\Omega}_q^{(0,1)} & \rightarrow \widetilde{\text{Pol}(\mathbb{C})_q}, \\ \psi dz \otimes dz^* \otimes dz \otimes dz^* \varphi & \mapsto \psi(1-zz^*)^4\varphi. \end{aligned}$$

What remains to elaborate is the fact that the corresponding tensor algebra is a $U_q\mathfrak{su}_{1,1}$ -module algebra.

In sections 2 – 5 we have considered a more simple case ($\Omega_q^{(0,*)}$ instead of Ω_q) in order to reduce the relevant computations and to make the exposition more plausible.

It should be noted that A. Connes assumed in his theory that the spectrum of the Dirac operator is discrete. This fails both for classical and quantum discs because of the non-compactness of $\mathbb{U}$. Even more, the spectrum of $(\bar{\partial})^*\bar{\partial}$ in the space of square summable functions is absolutely continuous in the classical case [2] and in the quantum case [10] as well. In particular, the spectrum of our Dirac operator is not discrete.

References

- [1] A. Connes, Non-Commutative Geometry, Academic Press inc., London, 1994.
- [2] S. Helgason. Groups and Geometric Analysis, Academic Press inc, Orlando et al, 1984.
- [3] J. C. Jantzen. Lectures on Quantum Groups. Graduate studies in Mathematics, Vol. 6, 1996, Amer. Math. Soc.
- [4] S. Klimek, A. Lesniewski, *A two-parameter quantum deformation of the unit disc*, J. Funct. Anal. **115**, (1993), 1 – 23.

³Note that the tilde symbol stands for the localization with respect to the multiplicative closed system $(1-zz^*)^{2\mathbb{N}}$.

- [5] A. Klimyk, K. Schmüdgen, *Quantum Groups and their Representations*, Springer-Verlag, Berlin et al., 1997.
- [6] M. L. Michelson, *Clifford and spinor cohomology on Kähler manifolds*, Amer. J. Math. **102** (1980), 1083 – 1146.
- [7] K. Schmüdgen, A Schüler *Covariant differential calculi on quantum spaces and on quantum groups*, C. R. Acad. Sci., Paris, Ser I **316** (1993), 1155 – 1160.
- [8] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum disc: the basic structures* (this volume).
- [9] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *On function theory in quantum disc: covariance*, math.QA/9808037, 1998, 16 p.
- [10] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *On function theory in quantum disc: integral representations*, math.QA/9808015, 1998, 17 p.
- [11] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *On function theory in quantum disc: q -differential equations and Fourier transform*, math.QA/9809002, 1998, 17 p.
- [12] S. Sinel'shchikov, A. Stolin, L. Vaksman, *Spherical principal non-degenerate series of representations for the quantum group $SU_{2,2}$* (this volume).

ON UNIQUENESS OF COVARIANT DEFORMATION WITH SEPARATION OF VARIABLES OF THE QUANTUM DISC

D. Shklyarov

1 Introduction

The concept of deformation quantization was introduced around 1977 by Bayen, Flato, Fronsdal, Lichnerowicz, and Sternheimer [1]. Since then the concept has become very popular. In this approach quantization means a deformation of the usual product of smooth function on a 'phase space' into a noncommutative associative product $\star_t$ (star-product) with additional properties.

The problem of constructing star-products explicitly is still of importance. For some special symplectic manifolds star-products can be constructed by using the famous Berezin quantization method [2]. Specifically, the method can be explored for a wide class of Kahler manifolds. The star-product on a Kahler manifold constructed by means of the Berezin scheme possesses some remarkable properties. First, it respects the complex structure of the Kahler manifold. Second, it respects the action of the group of biholomorphic automorphisms of the Kahler manifold.

In general, it seems to be important to look for those star-products on a symplectic manifold which keep some additional geometric structures on this manifold. In this connection, we want to mention the result on complete classification of star-products with separation of variables on Kahler manifolds [8] and results concerning invariant star-products on homogeneous symplectic manifolds [4], [5].

A remarkable class of Kahler manifolds to which the Berezin method is applicable is the class of bounded symmetric domains. The simplest example is the unit disc within complex plane $\mathbb{C}$. The star-products on bounded symmetric domains arising from the Berezin method were studied in [10] in the simplest case of the disc and in [6] in general setting.

Recently, in the framework of the quantum group theory q -analogues of bounded symmetric domains have been constructed [13]. In [14] we used a q -analog of the Berezin method to produce a deformation of product in a noncommutative algebra of 'functions on the quantum unit disc' (see section 2 for definitions). This deformation respects both the complex structure and the quantum group symmetry action (precise definitions are to be found in section 3). The main result of this paper is Theorem 5.1 which states that the deformation constructed in [14] is essentially the unique deformation possessing the above properties (specifically, any deformation of that kind can be obtained from the Berezin one via change of parameter).

Acknowledgement. I thank L. Vaksman for posing the problem and fruitful discussions, and E. Karolinsky for many valuable remarks on improving the text. Unfortunately,

This research was supported in part by of the US Civilian Research & Development Foundation (Award No UM1-2091) and by Swedish Academy of Sciences (project No 11293562).

I didn't succeed to take into account all his remarks, and I hope to do it in forthcoming versions of this paper.

2 The algebra $\text{Pol}(\mathbb{C})_q$

Everywhere in the sequel we suppose that $q \in (0, 1)$ and the ground field is the field $\mathbb{C}$ of complex numbers.

Denote by $\text{Pol}(\mathbb{C})_q$ the involutive unital algebra given by its generator z and the unique commutation relation

$$z^* z = q^2 z z^* + 1 - q^2. \quad (2.1)$$

This algebra was studied in [11]. It serves as the initial object in constructing function theory in the quantum unit disc (see [12]).

The algebra $\text{Pol}(\mathbb{C})_q$ may be endowed with an important extra structure, namely, the structure of a $U_q \mathfrak{su}_{1,1}$ -module algebra. Let us recall one what it means.

To start with, remind the definition of the quantum universal enveloping algebra $U_q \mathfrak{sl}_2$ and its "real form" $U_q \mathfrak{su}_{1,1}$. $U_q \mathfrak{sl}_2$ is a Hopf algebra over $\mathbb{C}$ determined by the generators K, K^{-1}, E, F and the relations

$$K K^{-1} = K^{-1} K = 1, \quad K^{\pm 1} E = q^{\pm 2} E K^{\pm 1}, \quad K^{\pm 1} F = q^{\mp 2} F K^{\pm 1},$$

$$EF - FE = \frac{K - K^{-1}}{q - q^{-1}},$$

$$\Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}, \quad \Delta(E) = E \otimes 1 + K \otimes E, \quad \Delta(F) = F \otimes K^{-1} + 1 \otimes F.$$

Note that

$$\varepsilon(E) = \varepsilon(F) = \varepsilon(K^{\pm 1} - 1) = 0,$$

$$S(K^{\pm 1}) = K^{\mp 1}, \quad S(E) = -K^{-1} E, \quad S(F) = -F K,$$

with $\varepsilon : U_q \mathfrak{sl}_2 \rightarrow \mathbb{C}$ and $S : U_q \mathfrak{sl}_2 \rightarrow U_q \mathfrak{sl}_2$ being the counit and the antipode of $U_q \mathfrak{sl}_2$, respectively. Equip $U_q \mathfrak{sl}_2$ with the involution given on the generators by

$$E^* = -K F, \quad F^* = -E K^{-1}, \quad (K^{\pm 1})^* = K^{\pm 1}. \quad (2.2)$$

The pair $(U_q \mathfrak{sl}_2, *)$ is the $*$ -Hopf algebra. It is denoted by $U_q \mathfrak{su}_{1,1}$.

Let $\mathcal{F}$ stands for a unital algebra equipped also with a structure of $U_q \mathfrak{sl}_2$ -module. $\mathcal{F}$ is called a $U_q \mathfrak{sl}_2$ -module algebra if the multiplication $m : \mathcal{F} \otimes \mathcal{F} \rightarrow \mathcal{F}$ is a morphism of $U_q \mathfrak{sl}_2$ -modules and for any $\xi \in U_q \mathfrak{sl}_2$

$$\xi(1) = \varepsilon(\xi) \cdot 1 \quad (2.3)$$

(in other words, the unit 1 of the algebra $\mathcal{F}$ is $U_q \mathfrak{sl}_2$ -invariant).

An involutive algebra $\mathcal{F}$ is said to be a $U_q \mathfrak{su}_{1,1}$ -module algebra if it is a $U_q \mathfrak{sl}_2$ -module one and

$$(\xi f)^* = (S(\xi))^* f^* \quad (2.4)$$

for any $\xi \in U_q \mathfrak{su}_{1,1}$ and $f \in \mathcal{F}$ (the first asterisk in the right hand side of (2.4) means the involution (2.2)).

We have explained the meaning of the term ' $U_q \mathfrak{su}_{1,1}$ -module algebra'. The following statement is proved in [13].

Proposition 2.1 *There exists a unique structure of $U_q\mathfrak{su}_{1,1}$ -module algebra in $\text{Pol}(\mathbb{C})_q$ such that*

$$K^{\pm 1}z = q^{\pm 2}z, \quad Ez = -q^{1/2}z^2, \quad Fz = q^{1/2}. \quad (2.5)$$

Note that uniqueness of $U_q\mathfrak{su}_{1,1}$ -module algebra structure in $\text{Pol}(\mathbb{C})_q$ satisfying (2.5) is a simple consequence of definitions. Indeed, after application of the involution to the both hand sides of equalities (2.5) we find that the following relations hold (see (2.4))

$$K^{\pm 1}z^* = q^{\mp 2}z^*, \quad Ez^* = q^{-3/2}, \quad Fz^* = -q^{-5/2}z^{*2}. \quad (2.6)$$

Relations (2.3), (2.5), and (2.6) allow one to apply $E, F, K^{\pm 1}$ to any element of $\text{Pol}(\mathbb{C})_q$. This implies the uniqueness. Existence of $U_q\mathfrak{su}_{1,1}$ -module algebra structure in $\text{Pol}(\mathbb{C})_q$ given by (2.5) is a more difficult fact and we don't adduce its proof (an analogous result is proved in [13] in a much more general setting).

Let $\mathbb{C}[z]_q$ and $\mathbb{C}[z^*]_q$ stand for the unital subalgebras in $\text{Pol}(\mathbb{C})_q$ generated by z and z^* , respectively. In view of (2.5) and (2.6), these subalgebras in the $U_q\mathfrak{sl}_2$ -module algebra $\text{Pol}(\mathbb{C})_q$ are $U_q\mathfrak{sl}_2$ -module algebras themselves (however, these subalgebras are not $U_q\mathfrak{su}_{1,1}$ -module subalgebras).

The defining relation (2.1) allows one to rewrite any element f of the algebra $\text{Pol}(\mathbb{C})_q$ in the form of a linear combination of "normally ordered" monomials:

$$f = \sum_{i,j} a_{ij} z^i z^{*j}, \quad a_{ij} \in \mathbb{C}.$$

Thus, there is a natural isomorphism of vector spaces

$$\text{Pol}(\mathbb{C})_q \simeq \mathbb{C}[z]_q \otimes \mathbb{C}[z^*]_q. \quad (2.7)$$

Evidently, this isomorphism respects the action of $U_q\mathfrak{sl}_2$.

In conclusion of this section, let us comment formulas (2.5) and (2.6). As we mentioned in Introduction, the unit disc is a homogeneous space of the group $SU(1,1)$. Formulas (2.5) and (2.6) are just q -analogs of the corresponding 'infinitesimal' action of $U\mathfrak{su}_{1,1}$ in the space of polynomials.

3 Deformations of the algebra $\text{Pol}(\mathbb{C})_q$.

Real, and covariant deformations

First of all, let us explain what the term 'deformation of the algebra $\text{Pol}(\mathbb{C})_q$ ' means. Let $\mathbb{C}[[t]]$ and $\text{Pol}(\mathbb{C})_q[[t]]$ stand for the algebra of formal power series in t over $\mathbb{C}$ and the $\mathbb{C}[[t]]$ -module of formal series in t over $\text{Pol}(\mathbb{C})_q$, respectively. According to the traditional approach [7], a deformation of the algebra $\text{Pol}(\mathbb{C})_q$ is an associative $\mathbb{C}[[t]]$ -bilinear product (called star-product)

$$\star_t : \text{Pol}(\mathbb{C})_q[[t]] \times \text{Pol}(\mathbb{C})_q[[t]] \rightarrow \text{Pol}(\mathbb{C})_q[[t]]$$

given for $f, g \in \text{Pol}(\mathbb{C})_q \subset \text{Pol}(\mathbb{C})_q[[t]]$ by

$$f \star_t g = f \cdot g + t \cdot m_1(f, g) + t^2 \cdot m_2(f, g) + \dots \quad (3.1)$$

with $m_i : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q \rightarrow \text{Pol}(\mathbb{C})_q$ being some $\mathbb{C}$ -bilinear maps. The product of arbitrary elements $f, g \in \text{Pol}(\mathbb{C})_q[[t]]$ is defined via the conditions of $\mathbb{C}$ -bilinearity and t -adic continuity.

In subsequent sections we consider deformations of certain class only. In this section we present all necessary definitions.

A deformation $\star_t$ of the algebra $\text{Pol}(\mathbb{C})_q$ is said to be deformation with separation of variables (see [8, 3]) if

$$f \star_t g = f \cdot g$$

with $f \in \mathbb{C}[z]_q$ or $g \in \mathbb{C}[z^*]_q$.

Remark 1. Deformations with separation of variables possess the following property: any such deformation can be reconstructed from the series $z^* \star_t z$.

A deformation $\star_t$ of the algebra $\text{Pol}(\mathbb{C})_q$ is called real if for any $f, g \in \text{Pol}(\mathbb{C})_q$

$$(f \star_t g)^* = g^* \star_t f^*$$

(we suppose the formal parameter t is real: $t^* = t$). In this case the algebra $\text{Pol}(\mathbb{C})_q[[t]]$ is an involutive algebra.

Finally, let us define the notion of a covariant deformation of the algebra $\text{Pol}(\mathbb{C})_q$. Its classical counterpart is the notion of a $SU(1, 1)$ -invariant deformation of the unit disc. Endow $\text{Pol}(\mathbb{C})_q[[t]]$ with a structure of $U_q \mathfrak{sl}_2$ -module via $\mathbb{C}[[t]]$ -linearity (and t -adic continuity). A deformation $\star_t$ of the algebra $\text{Pol}(\mathbb{C})_q$ is said to be covariant if the algebra $\text{Pol}(\mathbb{C})_q[[t]]$ is a $U_q \mathfrak{sl}_2$ -module algebra with respect to the product $\star_t$ (see section 2 for definitions).¹

Further we restrict ourselves of studying real covariant deformations with separation of variables (abbreviated to RCW-deformations) of the algebra $\text{Pol}(\mathbb{C})_q$ only.

4 Examples of RCW-deformations of the algebra $\text{Pol}(\mathbb{C})_q$

The simplest example of a RCW-deformation is the trivial deformation (the trivial deformation is the deformation $\star_t$ such that $f \star_t g = f \cdot g$ for any $f, g \in \text{Pol}(\mathbb{C})_q$). The remaining part of this section is devoted to a more substantial example, namely, the Berezin deformation. This deformation is constructed in [14] by a q -analogue of the Berezin quantization method. This approach is applicable to all q -Cartan domains [13]. In the simplest case of quantum disc this method could be replaced by a plausible procedure, to be sketched below.

Let us begin with some non-rigorous arguments. Suppose $t \in [0, 1)$. Consider the unital involutive algebra given by its generator and the following relation [9]:

$$z^* z = q^2 z z^* + 1 - q^2 + \frac{(1 - q^2)t}{1 - q^2 t} (1 - z^* z)(1 - z z^*). \quad (4.1)$$

We denote it by $\text{Pol}(\mathbb{C})_{q,t}$. Note that application of the involution to the both hand sides of (4.1) leads to the relation

$$z^* z = q^2 z z^* + 1 - q^2 + \frac{(1 - q^2)t}{1 - q^2 t} (1 - z z^*)(1 - z^* z). \quad (4.2)$$

In particular, the elements $1 - z z^*$ and $1 - z^* z$ commute.

We describe the procedure of producing the Berezin deformation [14] starting from the family of algebras $\text{Pol}(\mathbb{C})_{q,t}$. Consider a polynomials f in (non-commuting) variables z ,

¹Note that if a deformation $\star_t$ is real and covariant then the algebra $\text{Pol}(\mathbb{C})_q[[t]]$ with the product $\star_t$ automatically become a $U_q \mathfrak{su}_{1,1}$ -module algebra.

z^* and try to put it in normal order using relation (4.1) ("to put in normal order" means to rewrite as a linear combination of the monomials $z^i z^{*j}$, $i, j \in \mathbb{Z}_+$). The procedure of normal ordering is just the iterative process: at every subsequent step of this process we look for all the monomials containing combination $z^* z$ at least once; then we replace precisely one such combination with the right hand side of (4.1) in each found monomial. In general, we can't put the polynomial f in normal order in a finite number of steps. But it is not difficult to observe that for an arbitrarily large N any polynomial f can be written in a finite number of steps in the form

$$f = f_0 + t f_1 + t^2 f_2 + \dots + t^N f_N + t^{N+1} f_{N+1} \quad (4.3)$$

with $f_0, f_1, \dots, f_N$ being normally ordered polynomials independent of t . Thus, after "infinite number" of steps we should have a formal series

$$f_0 + t f_1 + t^2 f_2 + \dots + t^N f_N + \dots \in \text{Pol}(\mathbb{C})_q[[t]]$$

whose coefficients are normally ordered polynomials. It can be proved that the series depends on the polynomial f only (i.e., it is independent of arbitrariness in choice of the combination $z^* z$ at every step of the process). We shall call it the asymptotic expansion of f .

Let us define a deformation $\star_t$ of the algebra $\text{Pol}(\mathbb{C})_q$ as follows. Because of isomorphism (2.7), it is sufficient to define $z^i z^{*j} \star_t z^k z^{*l}$. We set

$$z^i z^{*j} \star_t z^k z^{*l} = \text{asymptotic expansion of } z^i z^{*j} z^k z^{*l}.$$

It is almost explicit from the algorithm of normal ordering that this star-product is associative and defines a deformation with separation of variables. Realness of this deformation is a less evident fact. ²

The deformation constructed in this way turns out to be covariant. It is a consequence of the following statement [14] (see also [9]).

Proposition 4.1 *There exists a unique structure of $U_q \mathfrak{su}_{1,1}$ -module algebra in $\text{Pol}(\mathbb{C})_{q,t}$ satisfying (2.5).*

Evidently, formulas (2.6) hold in $\text{Pol}(\mathbb{C})_{q,t}$ as well. To explain covariance of the deformation it suffices to note that the $U_q \mathfrak{sl}_2$ -action in $\text{Pol}(\mathbb{C})_{q,t}$ is independent of t , commutes with multiplication by t , and respects the normal order.

We have explained roughly the idea of constructing the Berezin deformation of the algebra $\text{Pol}(\mathbb{C})_q$. Let us expound the above 'algorithm' rigorously.

To start with, let us agree about the following notation. If F is a $\mathbb{C}[[t]]$ -module then F_n stands for the $\mathbb{C}[[t]]$ -module $F/t^n F$ and $\overline{F}$ for the completion of F in the t -adic topology. Note that if F is an (involutive) $\mathbb{C}[[t]]$ -algebra then F_n and $\overline{F}$ are naturally endowed with structures of an (involutive) $\mathbb{C}[[t]]$ -algebra and a topological (involutive) $\mathbb{C}[[t]]$ -algebra, respectively.

Consider the unital involutive $\mathbb{C}[[t]]$ -algebra given by its generator and relation (4.1). Denote it by $\mathcal{P}$. Evidently, relation (4.2) holds in this algebra as well. Using proposition

²To prove the realness one actually should show that the procedure of normal ordering "commutes" with application of the involution. It is easy to observe that application of the involution to a polynomial f followed by normal ordering via relation (4.1) gives us the same series as normal ordering via relation (4.2) instead of (4.1) followed by application of the involution. Now realness of the deformation is due to the following fact: the procedures of normal ordering via (4.1) and (4.2) lead to the same result.

4.1, one shows that there exists a unique structure of $U_q\mathfrak{su}_{1,1}$ -module algebra in $\mathcal{P}$ given by (2.5) and (2.6). Denote the $\mathbb{C}[[t]]$ -algebra $\text{Pol}(\mathbb{C})_q \otimes \mathbb{C}[[t]]$ by P . Endow the latter algebra with a structure $U_q\mathfrak{su}_{1,1}$ -module algebra via $\mathbb{C}[[t]]$ -linearity.

Let I_n stands for the unique $\mathbb{C}[[t]]$ -linear map from P_n to $\mathcal{P}_n$ such that $I_n : z^i z^{*j} \mapsto z^i z^{*j}$. Certainly, I_n is an embedding of $\mathbb{C}[[t]]$ -modules and respects $U_q\mathfrak{sl}_2$ -actions. Moreover, $I_n(f^*) = (I_n(f))^*$ for any $f \in P_n$. I_n turns out to be an isomorphism of $\mathbb{C}[[t]]$ -modules. Let us prove this via induction on n .

The case $n = 1$ is evident (moreover, due to (2.1) and (4.1), I_1 is an isomorphism of $U_q\mathfrak{su}_{1,1}$ -module algebras). Suppose I_k is an isomorphism of $\mathbb{C}[[t]]$ -modules for any $k \leq n$. It suffices to show that the embedding $I_{n+1} : P_{n+1} \rightarrow \mathcal{P}_{n+1}$ is surjective.

Let $\varphi_{n+1} : P_{n+1} \rightarrow P_n$ and $\psi_{n+1} : \mathcal{P}_{n+1} \rightarrow \mathcal{P}_n$ stand for the natural surjective $\mathbb{C}[[t]]$ -linear maps induced by the inclusions $t^{n+1}P \subset t^n P$ and $t^{n+1}\mathcal{P} \subset t^n \mathcal{P}$. It is clear that

$$\psi_{n+1} \circ I_{n+1} = I_n \circ \varphi_{n+1}. \quad (4.4)$$

Let $f \in \mathcal{P}_{n+1}$. In view of the inductive hypothesis, there exists $g \in P_{n+1}$ such that $I_n \circ \varphi_{n+1}(g) = \psi_{n+1}(f)$. (4.4) implies $\psi_{n+1} \circ I_{n+1}(g) = \psi_{n+1}(f)$, i.e., $I_{n+1}(g) - f \in \text{Ker} \psi_{n+1}$. In other words, $I_{n+1}(g) - f = t^n g_0$ for some $g_0 \in \mathcal{P}_{n+1}$. Relation (4.1) and the definition of the maps I_n imply $t^n \mathcal{P}_{n+1} \subset \text{Im} I_{n+1}$. By (4.4) $f \in \text{Im} I_{n+1}$. So, $I_n : P_n \rightarrow \mathcal{P}_n$ is an isomorphism of $\mathbb{C}[[t]]$ -modules.

The family of isomorphisms I_n , $n = 1, 2, \dots$, satisfies (4.4). It means that there exists the limit isomorphism of topological $\mathbb{C}[[t]]$ -modules

$$I_\infty : \overline{P} \rightarrow \overline{\mathcal{P}}.$$

Evidently, $\overline{P}$ and $\overline{\mathcal{P}}$ inherit structures of $U_q\mathfrak{su}_{1,1}$ -module $\mathbb{C}[[t]]$ -algebras (moreover, $U_q\mathfrak{su}_{1,1}$ -module $\mathbb{C}[[t]]$ -algebra $\overline{P}$ is isomorphic to $\text{Pol}(\mathbb{C})_q[[t]]$). The map I_∞ is an isomorphism of $U_q\mathfrak{sl}_2$ -modules and intertwines actions of the involutions in $\overline{P}$ and $\overline{\mathcal{P}}$. But I_∞ is not an algebra isomorphism. Define in $\overline{P} \simeq \text{Pol}(\mathbb{C})_q[[t]]$ a new product by

$$f \star_t g = I_\infty^{-1} (I_\infty(f) \cdot I_\infty(g)).$$

Evidently, $\text{Pol}(\mathbb{C})_q[[t]]$ endowed with this new product become a $U_q\mathfrak{su}_{1,1}$ -module algebra.

This deformation is just the one described informally in the first part of this section. We shall call it the Berezin deformation of $\text{Pol}(\mathbb{C})_q$ ([14]) and denote the attached star-product by $\star_B$.

Remark 2. We present in [14] explicit formulas for $\star_B$, i.e., for the corresponding bilinear maps m_i (see (3.1)).

5 Uniqueness of the RCW-deformation of the algebra $\text{Pol}(\mathbb{C})_q$

First of all, let us consider the following construction. Suppose $\star_t$ is a deformation of the algebra $\text{Pol}(\mathbb{C})_q$ and $m_i : \text{Pol}(\mathbb{C})_q \times \text{Pol}(\mathbb{C})_q \rightarrow \text{Pol}(\mathbb{C})_q$ are the attached bilinear maps (see (3.1)). Let $c(t) \in \mathbb{C}[[t]]$, $c(0) = 0$. Using $c(t)$, one can define a new star-product $\star'_t$ given by

$$f \star'_t g = f \cdot g + c(t) \cdot m_1(f, g) + c(t)^2 \cdot m_2(f, g) + \dots \quad (5.1)$$

(The product of arbitrary elements $f, g \in \text{Pol}(\mathbb{C})_q[[t]]$ is defined via the conditions of $\mathbb{C}$ -bilinearity and t -adic continuity.) It is said that $\star'_t$ is obtained from $\star_t$ via change of

formal parameter. If the initial deformation $\star_t$ is a RCW-deformation and the series $c(t)$ is real then $\star'_t$ is a RCW-deformation as well.

The principal result of this paper is

Theorem 5.1 *Any RCW-deformation of the algebra $\text{Pol}(\mathbb{C})_q$ can be obtained from the Berezin deformation via change of formal parameter.*

Proof of the Theorem. Let us use the traditional notation $M[[t]]$ for the $\mathbb{C}[[t]]$ -module of formal series in t over a vector space M .

Let $\star_t$ denotes a RCW-deformation of the algebra $\text{Pol}(\mathbb{C})_q$. Endow $\text{Pol}(\mathbb{C})_q[[t]]$ with the algebra structure using the star-product $\star_t$. To prove the theorem we shall show that the elements z and z^* of the algebra $\text{Pol}(\mathbb{C})_q[[t]]$ satisfy a commutation relation which can be obtained from (4.1) via change of formal parameter.

To start with, let us construct an embedding of the algebra $\text{Pol}(\mathbb{C})_q[[t]]$ into the algebra of endomorphisms of a $\mathbb{C}[[t]]$ -module.

Let J stands for the left ideal in $\text{Pol}(\mathbb{C})_q[[t]]$ generated by z^* :

$$J = \text{Pol}(\mathbb{C})_q[[t]] \star_t z^*.$$

Since $\star_t$ is a product with separation of variables, we have $J = J_0[[t]]$ with $J_0 = \text{Pol}(\mathbb{C})_q z^*$. Thus, $\text{Pol}(\mathbb{C})_q[[t]]$ splits into the direct sum of $\mathbb{C}[[t]]$ -submodules

$$\text{Pol}(\mathbb{C})_q[[t]] = J + \mathbb{C}[z]_q[[t]]. \quad (5.2)$$

From (5.2), we have the natural isomorphism of $\mathbb{C}[[t]]$ -modules

$$\varphi : \text{Pol}(\mathbb{C})_q[[t]]/J \rightarrow \mathbb{C}[z]_q[[t]]. \quad (5.3)$$

The $\mathbb{C}[[t]]$ -module $\text{Pol}(\mathbb{C})_q[[t]]/J$ is endowed with the evident $\text{Pol}(\mathbb{C})_q[[t]]$ -module structure. Using isomorphism (5.3), one can "transfer" $\text{Pol}(\mathbb{C})_q[[t]]$ -module structure onto $\mathbb{C}[z]_q[[t]]$. As a result, we get the algebra homomorphism

$$T : \text{Pol}(\mathbb{C})_q[[t]] \rightarrow \text{End}_{\mathbb{C}[[t]]}(\mathbb{C}[z]_q[[t]]).$$

Lemma 5.2 *T is an embedding.*

Proof of Lemma 5.2. To prove injectivity of T , we shall describe more or less explicitly the endomorphisms $T(z)$ and $T(z^*)$ of the $\mathbb{C}[[t]]$ -module $\mathbb{C}[z]_q[[t]]$. It is easy to describe the former endomorphism. Indeed, each direct summand in (5.2) is invariant under multiplication from the left by elements of the subalgebra $\mathbb{C}[z]_q[[t]] \subset \text{Pol}(\mathbb{C})_q[[t]]$. This implies

$$T(z) : z^n \mapsto z^{n+1}.$$

To describe explicitly $T(z^*)$, one should use an extra structure in the $\text{Pol}(\mathbb{C})_q[[t]]$ -module $\mathbb{C}[z]_q[[t]]$ arising from the covariance of the deformation $\star_t$.

Let us consider the Hopf subalgebra $U_q \mathfrak{b}_- \subset U_q \mathfrak{sl}_2$ generated by $K^{\pm 1}$ and F . Certainly, J is a $U_q \mathfrak{b}_-$ -invariant subspace in $\text{Pol}(\mathbb{C})_q[[t]]$ (see the definition of J and relations (2.5),(2.6)). Thus, the $\mathbb{C}[[t]]$ -module $\text{Pol}(\mathbb{C})_q[[t]]/J$ is endowed with a natural structure of a $U_q \mathfrak{b}_-$ -module. Moreover, since $\text{Pol}(\mathbb{C})_q[[t]]$ is a $U_q \mathfrak{sl}_2$ -module algebra, $\text{Pol}(\mathbb{C})_q[[t]]/J$ is a $U_q \mathfrak{b}_-$ -module $\text{Pol}(\mathbb{C})_q[[t]]$ -module, i.e., the multiplication map

$$\text{Pol}(\mathbb{C})_q[[t]] \otimes (\text{Pol}(\mathbb{C})_q[[t]]/J) \rightarrow (\text{Pol}(\mathbb{C})_q[[t]]/J)$$

is a morphism of $U_q\mathfrak{b}_-$ -modules. Furthermore, the $U_q\mathfrak{b}_-$ -action respects decomposition (5.2) (this is a direct consequence of (2.5) and (2.6)). Therefore, the structure of $\text{Pol}(\mathbb{C})_q[[t]]$ -module in $\mathbb{C}[z]_q[[t]]$ (defined above via isomorphism (5.3)) is compatible with the natural $U_q\mathfrak{b}_-$ -module structure. In other words, $\mathbb{C}[z]_q[[t]]$ is a $U_q\mathfrak{b}_-$ -module $\text{Pol}(\mathbb{C})_q[[t]]$ -module. This observation allows to calculate $T(z^*)$.

Suppose that $T(z^*) : z^n \mapsto \phi_n(z)$ for some $\phi_n(z) \in \mathbb{C}[z]_q[[t]]$. Formulas (2.5), (2.6) and $U_q\mathfrak{b}_-$ -moduleness of the $\text{Pol}(\mathbb{C})_q[[t]]$ -module $\mathbb{C}[z]_q[[t]]$ -module imply $K(\phi_n(z)) = q^{2n-2}\phi_n(z)$. It is clear that the latter equality holds only when $\phi_n(z) = c_n(t)z^{n-1}$, $c_n(t) \in \mathbb{C}[[t]]$. It is possible to calculate $c(0)$: since

$$z^* \star_t z \equiv q^2 z z^* + 1 - q^2 \pmod{t}$$

(see (2.1)), we get

$$z^* \star_t z^n \equiv q^{2n} z^n z^* + (1 - q^{2n}) z^{n-1} \pmod{t}.$$

Thus, $c_n(t) \equiv 1 - q^{2n} \pmod{t}$. As a result, we have

$$T(z) : z^n \mapsto z^{n+1}, \quad (5.4)$$

$$T(z^*) : z^n \mapsto c_n(t) z^{n-1}, \quad (5.5)$$

with

$$c_n(t) \equiv 1 - q^{2n} \pmod{t}. \quad (5.6)$$

We are now in a position to prove injectivity of T . Let $f = \sum_{i,j} a_{ij} z^i z^{*j} \in \text{Pol}(\mathbb{C})_q[[t]]$ be a non-zero element such that $T(f) = 0$. Suppose j_0 is the least non-negative integer such that $a_{ij_0} \neq 0$ for some i . From (5.4) and (5.5),

$$T(f) : z^{j_0} \mapsto \sum_i a_{ij_0} z^i.$$

Thus, $a_{ij_0}(t)c_{j_0}(t)c_{j_0-1}(t)\dots c_1(t) = 0$ for any i . In view of (5.6), $a_{ij_0}(t) = 0$ for any i . It contradicts our choice of j_0 . Q.E.D.

Lemma 5.3 *There exists a formal series $c(t)$ with real coefficients such that $c(0) = 0$ and*

$$T(z^*)T(z) = q^2 T(z)T(z^*) + 1 - q^2 + \frac{(1 - q^2)t}{1 - q^2 t} (1 - T(z)T(z^*)) (1 - T(z^*)T(z)). \quad (5.7)$$

Proof of Lemma 5.3. Let us invoke $U_q\mathfrak{b}_-$ -moduleness of the $\text{Pol}(\mathbb{C})_q[[t]]$ -module $\mathbb{C}[z]_q$ once more. Apply the element $F \in U_q\mathfrak{b}_-$ to both hand sides of the equality $T(z^*)(z^n) = c_n(t)z^{n-1}$ (see (5.5)):

$$T(Fz^*)(K^{-1}z^n) + T(z^*)(Fz^n) = c_n(t)Fz^{n-1}. \quad (5.8)$$

Equalities (2.5) and easy induction on n give

$$K^{-1}z^n = q^{-2n}z^n, \quad Fz^n = q^{1/2} \frac{1 - q^{-2n}}{1 - q^{-2}} z^{n-1}. \quad (5.9)$$

According to (2.6) and (5.9), (5.8) can be rewritten as follows

$$-q^{5/2}T(z^{*2})(q^{-2n}z^n) + q^{1/2}\frac{1-q^{-2n}}{1-q^{-2}}T(z^*)(z^{n-1}) = q^{1/2}\frac{1-q^{2-2n}}{1-q^{-2}}c_n(t)z^{n-2}. \quad (5.10)$$

From (5.5),

$$-q^{5/2-2n}c_n(t)c_{n-1}(t)z^{n-2} + q^{1/2}\frac{1-q^{-2n}}{1-q^{-2}}c_{n-1}(t)z^{n-2} = q^{1/2}\frac{1-q^{2-2n}}{1-q^{-2}}c_n(t)z^{n-2}. \quad (5.11)$$

For $n = 0$ or $n = 1$, the latter equality holds trivially. For $n > 1$, it is equivalent to the following 'recurrence' relation

$$-q^{2-2n}(1-q^{-2})c_n(t)c_{n-1}(t) + (1-q^{-2n})c_{n-1}(t) = (1-q^{2-2n})c_n(t). \quad (5.12)$$

Let us define the series $c(t)$ by

$$c_1(t) = \frac{1-q^2}{1-q^2c(t)}. \quad (5.13)$$

From (5.6), we have $c(0) = 0$. In view of (5.12),

$$c_n(t) = \frac{1-q^{2n}}{1-q^{2n}c(t)} \quad (5.14)$$

(it suffices to proceed by induction on n). So, there exists a series $c(t)$ such that $c(0) = 0$ and

$$T(z)(z^n) = z^{n+1}, \quad (5.15)$$

$$T(z^*)(z^n) = \frac{1-q^{2n}}{1-q^{2n}c(t)}z^{n-1}. \quad (5.16)$$

The latter equalities imply (5.7). Now, to finish proof of Lemma 5.3, it remains to show that the series $c(t)$ given by (5.14) has real coefficients. Because of the definition of $c_1(t)$,

$$z^* \star_t z = c_1(t) + f \quad (5.17)$$

for some $f \in J$. Apply the element $K \in U_q \mathfrak{sl}_2$ to the both hand sides of (5.17). Since the deformation $\star_t$ is covariant, we have $Kf = f$, i.e., $f = \sum_j a_j(t)z^j z^{*j}$. Apply the involution to the both hand sides of the equality

$$z^* \star_t z = c_1(t) + \sum_j a_j(t)z^j z^{*j}.$$

Now realness of coefficients of $c_1(t)$ is due to realness of the deformation. It follows that coefficients of $c(t)$ are real as well (see (5.14)). Lemma 5.3 is proved.

Lemmas 5.2 and 5.3 imply existence of a series $c(t)$ with real coefficient such that $c(0) = 0$ and

$$z^* \star_t z = q^2 z \star_t z^* + 1 - q^2 + \frac{(1-q^2)c(t)}{1-q^2c(t)}(1 - z \star_t z^*) \star_t (1 - z^* \star_t z). \quad (5.18)$$

It is not difficult to finish the proof of Theorem 5.1. The Berezin deformation $\star_B$ certainly satisfies the condition

$$z^* \star_B z = q^2 z \star_B z^* + 1 - q^2 + \frac{(1 - q^2)t}{1 - q^2 t} (1 - z \star_B z^*) \star_B (1 - z^* \star_B z)$$

(see the previous section). Moreover, the series $z^* \star_B z$ can be reconstructed from the latter relation in a unique way. Similarly, there are no two different deformations with separation of variables satisfying (5.18) (see Remark 1). It is clear how to construct one such deformation: for $f, g \in \text{Pol}(\mathbb{C})_q$ one should set

$$f \star_t g = f \cdot g + c(t) \cdot m_1(f, g) + c(t)^2 \cdot m_2(f, g) + \dots,$$

with m_i given by

$$f \star_B g = f \cdot g + t \cdot m_1(f, g) + t^2 \cdot m_2(f, g) + \dots$$

It is straightforward now that the deformation $\star_t$ can be obtained from the Berezin one via change of formal parameter. Theorem 5.1 is proved.

Remark 3. At first sight, we didn't invoke the full "quantum symmetry" in the proof of Theorem 5.1 (i. e., $U_q \mathfrak{su}_{1,1}$ -covariance of the algebra $\text{Pol}(\mathbb{C})_q[[t]]$). What we really used was the $U_q \mathfrak{b}_-$ -covariance. But it is simple to show that realness and $U_q \mathfrak{b}_-$ -covariance of a deformation with separation of variables imply its $U_q \mathfrak{su}_{1,1}$ -covariance.

References

- [1] *F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer*, Deformation theory and quantization, Part I. Lett. Math. Phys., **1** (1977), 521 – 530.
- [2] *F. A. Berezin*, Quantisation of Kahler manifold. Comm. Math. Phys., **40** (1975), 153.
- [3] *M. Bordemann and S. Waldmann*, A Fedosov Star Product of Wick Type for Kahler Manifolds. q-alg/9605012.
- [4] *M. Cahen and S. Gutt*, Produits- $\star$ sur un espace riemannien symétrique. C. R. Acad. Sci., Paris, **291A** (1980), 543 – 545.
- [5] *M. Cahen and S. Gutt*, Quantization of Kahler Manifolds. IV Lett. Math. Phys., **34** (1995), 159 – 168.
- [6] *M. Cahen, S. Gutt, and J. Rawnsley*, Invariant $\star$ -products of holomorphic functions on the hyperbolic Hermitian spaces. Lett. Math. Phys., **5** (1981), 219 – 228.
- [7] *M. Gerstenhaber*, On the deformation of rings and algebras. Ann. Math., **79** (1964), 59 – 103.
- [8] *A. Karabegov*, Deformation quantizations with separation of variables on a Kahler Manifold. Comm. Math. Phys., **180** (1996), 745 – 755.
- [9] *S. Klimek and A. Lesniewski*, A Two-Parameter Quantum Deformation of the Unit Disc. J. Funct. A., **115** (1993), 1 – 23.
- [10] *C. Moreno and P. Ortega-Navarro*, $\star$ -Products on $D^1(\mathbb{C})$, S^2 and Related Spectral Analysis. Lett. Math. Phys., **7** (1983), 181 – 193.

- [11] *G. Nagy and A. Nica*, On the 'quantum disc' and 'non-commutative circle'. in Algebraic Methods in Operator Theory, R. E. Curto, P. E. T. Jorgensen (eds.), Birkhauser, Boston (1994), 276 – 290.
- [12] *D. Shklyarov, S. Sinel'shchikov, and L. Vaksman*, Quantum disc: the basic structures. This volume.
- [13] *S. Sinel'shchikov and L. Vaksman*, Harish-Chandra Embedding and q-Analogues of Bounded Symmetric Domains. This volume.
- [14] *D. Shklyarov, S. Sinel'shchikov, and L. Vaksman*, On function theory in quantum disc: a q-analogue of the Berezin transform. E-print: math.QA/9809018.

PART II
UNITARY REPRESENTATIONS AND
NON-COMMUTATIVE HARMONIC ANALYSIS

HARISH-CHANDRA EMBEDDING AND q-ANALOGUES OF BOUNDED SYMMETRIC DOMAINS

S. Sinel'shchikov ^{*} and *L. Vaksman* [†]

Institute for Low Temperature Physics & Engineering
National Academy of Sciences of Ukraine

1. This work is devoted to study of a very restricted class of homogeneous spaces associated to quantum groups [1, 4]. We follow [8] in describing here the construction of algebras of functions and differential forms on these quantum homogeneous spaces.

We hope to extend to the above context a great deal of the results of function theory and harmonic analysis in bounded symmetric domains [3]. This is shown here to be available for the simplest one among such domains, the quantum disc [9].

Our subsequent constructions are q-analogues of the corresponding Harish-Chandra's constructions which allow one to embed a Hermitian symmetric space of non-compact type into $\mathbb{C}^N$ [2].

Let A be a Hopf algebra, ε its counit, and S its antipode. Consider an algebra F equipped also with a structure of A -module. F is said to be an A -module (covariant) algebra if

- i) the multiplication $m : F \otimes F \rightarrow F$, $m : f_1 \otimes f_2 \mapsto f_1 \cdot f_2$; $f_1, f_2 \in F$ is a morphism of A -modules;
- ii) the unit $1 \in F$ is an invariant: $\xi 1 = \varepsilon(\xi) 1$, $\xi \in A$.

If A is a Hopf $*$ -algebra, and F is also equipped with an involution, then the definition of covariance should include the following compatibility condition for involutions:

$$\forall \xi \in A, f \in F \quad (\xi f)^* = (S(\xi))^* f^*.$$

In the sequel all the algebras of "functions" ($\mathbb{C}[\mathfrak{g}_{-1}]_q$, $\mathbb{C}[\bar{\mathfrak{g}}_{-1}]_q$, $\text{Pol}(\mathfrak{g}_{-1})_q$) and "differential forms" are covariant algebras.

2. Let $\mathfrak{g}$ be a simple complex Lie algebra, $\mathfrak{h} \subset \mathfrak{g}$ a Cartan subalgebra, $\alpha_j \in \mathfrak{h}^*$, $j = 1, \dots, l$, a system of simple roots with α_{j_0} being one of those roots. Consider the $\mathbb{Z}$ -grading $\mathfrak{g} = \bigoplus_m \mathfrak{g}_m$ given by

$$\mathfrak{g}_m = \{\xi \in \mathfrak{g} \mid [H_0, \xi] = 2m\xi\},$$

with $H_0 \in \mathfrak{h}$ such that

$$\alpha_j(H_0) = 0, j \neq j_0; \quad \alpha_{j_0}(H_0) = 2.$$

This lecture has been delivered at D. Volkov Memorial Seminar Held in Kharkov, January 1997 and published in *Supersymmetry and Quantum Field Theory*, J. Wess and P. Akulov (eds). Springer, 1998, 312–316.

^{*}Partially supported by ISF grant U2B200 and grant DKNT-1.4/12

[†]Partially supported by the grant INTAS-94-4720, ISF grant U21200 and grant DKNT-1.4/12

If this grading terminates,

$$\mathfrak{g} = \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_{+1},$$

then clearly $\mathfrak{g}_{\pm 1}$ are Abelian Lie subalgebras. This is just the case when Harish-Chandra's construction presents a bounded symmetric domain U in the vector space $\mathfrak{g}_{-1}$.

Note also that in the case $\mathfrak{g} = \mathfrak{sl}_{m+n}$, $\alpha_{j_0} = \alpha_m$, U is the matrix ball in the space of $m \times n$ matrices:

$$\mathfrak{g}_{-1} \simeq \text{Mat}(m, n); \quad U = \{T \in \text{Mat}(m, n) \mid \|T\| < 1\}.$$

3. Turn to the quantum case and fix $q \in (0, 1)$.

Remind that the Hopf algebra $U_q \mathfrak{sl}_2$ is determined by its generators $K^{\pm 1}, E, F$ and the relations

$$\begin{aligned} K \cdot K^{-1} &= K^{-1} \cdot K = 1, & K^{\pm 1} \cdot E &= q^{\pm 2} E K^{\pm 1}, \\ K^{\pm 1} \cdot F &= q^{\mp 2} F K^{\pm 1}, & EF - FE &= (K - K^{-1})/(q - q^{-1}). \end{aligned}$$

Comultiplication $\Delta : U_q \mathfrak{sl}_2 \rightarrow U_q \mathfrak{sl}_2 \otimes U_q \mathfrak{sl}_2$ is given by $\Delta(E) = E \otimes 1 + K \otimes E$, $\Delta(F) = F \otimes K^{-1} + 1 \otimes F$, $\Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}$.

This Hopf algebra was introduced by E. Sklyanin, and its generalization $U_q \mathfrak{g}$ to the case of an arbitrary simple Lie algebra $\mathfrak{g}$ in the works of V. Drinfeld and M. Jimbo [1]. $U_q \mathfrak{g}$ is determined by the generators $\{K_j^{\pm 1}, E_j, F_j\}_{j=1, \dots, l}$ and the well known relations [4]. In this setting, every simple root $\alpha_j, j = 1, \dots, l$, generates an embedding $\varphi_j : U_q \mathfrak{sl}_2 \rightarrow U_q \mathfrak{g}$ given by

$$\varphi_j : K^{\pm 1} \mapsto K_j^{\pm 1}, \quad \varphi_j : E \mapsto E_j, \quad \varphi_j : F \mapsto F_j.$$

Here $q_j = q^{d_j}$ with $d_j > 0$ such that $d_i a_{ij} = a_{ji} d_j$ for all i, j .

4. Equip $U_q \mathfrak{g}$ with a structure of graded algebra:

$$\begin{aligned} \deg K_j &= \deg E_j = \deg F_j = 0, & j &\neq j_0 \\ \deg K_{j_0} &= 0, & \deg E_{j_0} &= 1, \quad \deg F_{j_0} = -1. \end{aligned}$$

The embedding $\mathfrak{g}_{-1} \subset U \mathfrak{g}$ has no good q -analog. This forces us to use the generalized Verma modules instead of $U \mathfrak{g}$.

Let V be a graded $U_q \mathfrak{g}$ -module determined by its generator $v \in V$ and the relations

$$\begin{aligned} E_i v &= 0, & K_i^{\pm 1} v &= v, & i &= 1, \dots, l, \\ F_j v &= 0, & j &\neq j_0. \end{aligned}$$

In the classic limit $q \rightarrow 1$ there is an embedding $\mathfrak{g}_{-1} \hookrightarrow V$, $\xi \mapsto \xi v$. This allows one to treat the homogeneous component $V_{-1} = \{v \in V \mid \deg v = -1\}$ as a q -analog of the vector space $\mathfrak{g}_{-1}$.

We need also the graded $U_q \mathfrak{g}$ -module V' given by its generator v' and the relations

$$\begin{aligned} E_i v' &= 0, & K_i^{\pm 1} v' &= q^{\mp a_{ij_0}} v', & i &= 1, \dots, l, \\ F_j^{-a_{ij_0}+1} v' &= 0, & j &\neq j_0; & \deg v' &= -1. \end{aligned}$$

5. Introduce the notation $\mathbb{C}[\mathfrak{g}_{-1}]_q = \bigoplus_m (V_m)^*$, $\bigwedge^1(\mathfrak{g}_{-1})_q = \bigoplus_m (V'_m)^*$ for the dual to the $U_q \mathfrak{g}$ -modules V and V' respectively graded $U_q \mathfrak{g}$ -modules. The elements $f \in \mathbb{C}[\mathfrak{g}_{-1}]_q$ are be

called holomorphic polynomials, and the elements $\omega \in \bigwedge^1(\mathfrak{g}_{-1})_q$ differential 1-forms. The linear operator $d : \mathbb{C}[\mathfrak{g}_{-1}]_q \rightarrow \bigwedge^1(\mathfrak{g}_{-1})_q$ is defined via the adjoint operator $d^* : V' \rightarrow V$. In turn, d^* is defined as the unique $U_q\mathfrak{g}$ -module morphism with $d^* : v' \mapsto F_{j_0}v$. Evidently, the differential d is a morphism of $U_q\mathfrak{g}^{op}$ -modules.

6. Besides the comultiplication $\Delta : U_q\mathfrak{g} \rightarrow U_q\mathfrak{g} \otimes U_q\mathfrak{g}^{op}$ we need also the opposite comultiplication Δ^{op} . It is used to equip the vector spaces $V \otimes V$, $V \otimes V'$, $V' \otimes V$ with a structure of $U_q\mathfrak{g}$ -modules.

The maps $v \mapsto v \otimes v$, $v' \mapsto v \otimes v'$, $v' \mapsto v' \otimes v$ admit the unique extensions to morphisms of $U_q\mathfrak{g}$ -modules

$$V \rightarrow V \otimes V, \quad V' \rightarrow V \otimes V', \quad V' \rightarrow V' \otimes V.$$

The adjoint operator to the comultiplication $V \rightarrow V \otimes V$ equips $\mathbb{C}[\mathfrak{g}_{-1}]_q$ with a structure of associative algebra. Similarly, the operators dual to the above morphisms $V' \rightarrow V \otimes V'$, $V' \rightarrow V' \otimes V$ equip $\bigwedge^1(\mathfrak{g}_{-1})_q$ with a structure of a bimodule over $\mathbb{C}[\mathfrak{g}_{-1}]_q$.

It is easy to show that $d(f_1 f_2) = df_1 \cdot f_2 + f_1 \cdot df_2$ for all $f_1, f_2 \in \mathbb{C}[\mathfrak{g}_{-1}]_q$. This allows one to pass from the 1-forms to the higher differential forms (see, for instance, the construction of G. Maltiniotis [5]).

7. It is possible to describe the above algebras by their generators and relations. Even in the simplest case $\mathfrak{g} = \mathfrak{sl}_N$ our approach yields the profound results [7].

It should be noted that our approach to the construction of algebras of differential forms is completely analogous to that of V. G. Drinfeld to the construction of the algebra of functions on a formal quantum group [1].

8. If we replace in the above construction the $U_q\mathfrak{g}$ -module V with a highest weight vector by the $U_q\mathfrak{g}$ -module with a lowest weight vector, we obtain the algebra $\mathbb{C}[\bar{\mathfrak{g}}_{-1}]_q$ of antiholomorphic polynomials on the quantum vector space $\mathfrak{g}_{-1}$.

The tensor product

$$\text{Pol}(\mathfrak{g}_{-1})_q = \mathbb{C}[\mathfrak{g}_{-1}]_q \otimes \mathbb{C}[\bar{\mathfrak{g}}_{-1}]_q$$

is equipped with a structure of algebra by means of the universal R-matrix together with the corresponding "commutativity morphism" [1]:

$$\check{R} : \mathbb{C}[\bar{\mathfrak{g}}_{-1}]_q \otimes \mathbb{C}[\mathfrak{g}_{-1}]_q \rightarrow \mathbb{C}[\mathfrak{g}_{-1}]_q \otimes \mathbb{C}[\bar{\mathfrak{g}}_{-1}]_q.$$

9. For the sake of passage from 'complex quantum Lie groups to real ones' equip the Hopf algebra $U_q\mathfrak{g}$ with an involution:

$$E_j^* = \begin{cases} K_j F_j & j \neq j_0 \\ -K_j F_j & j = j_0 \end{cases}, \quad F_j^* = \begin{cases} E_j K_j^{-1} & j \neq j_0 \\ -E_j K_j^{-1} & j = j_0 \end{cases},$$

$$(K_j^{\pm 1})^* = K_j^{\pm 1}, \quad i, j \in \{1, \dots, l\}.$$

The involution in $\text{Pol}(\mathfrak{g}_{-1})_q$ presented in [8] possess the property $\forall \xi \in U_q\mathfrak{g}, f \in \text{Pol}(\mathfrak{g}_{-1})_q$ $(\xi f)^* = (S(\xi))^* f^*$.

In the simplest case $\mathfrak{g} = \mathfrak{sl}_2$ one obtains the well known $*$ -algebra given by the generators z, z^* and the relation $z^* z - q^2 z z^* = 1 - q^2$ [8].

The passage from the polynomial algebra to the algebra of continuous functions in the closure of a bounded symmetric domain is made by means of a C^* -completion. In the special case of quantum disc this argument was used, in particular, in [6].

In [9] the q -analogs for the basic integral representations of the function theory in the quantum disc were obtained. Besides, there are several results for the quantum ball [10].

10. Finally, we express our gratitude to V. P. Akulov for helpful discussions of the results of this work.

References

- [1] V. G. Drinfeld, *Quantum groups*, in Proceedings of the International Congress of Mathematicians, Berkeley, 1986, A. M. Gleason (ed.), 1987, American Mathematical Society, Providence RI, 798 - 820.
- [2] S. Helgason. *Differential Geometry and Symmetric Spaces*, Acad. Press, N.-Y. – London, 1962.
- [3] L.-K. Hua. *Harmonic analysis of functions of several complex variables in the classical domains*, Transl. Math. Mono, Vol. 6, Amer. Math. Soc., Providence RI, 1963.
- [4] J. C. Jantzen. *Lectures on Quantum Groups*. Graduate studies in Mathematics, Vol. 6, 1996. Amer. Math. Soc.
- [5] G. Maltsiniotis, *Le langage des espaces et des groupes quantiques*, Commun. Math. Phys., **151**, (1993), 275 - 302.
- [6] G. Nagy, A. Nica. *On the "quantum disc" and a "non-commutative circle"*, in: Algebraic Methods on Operator Theory, R. E. Curto, P. E. T. Jorgensen (eds.), Birkhauser, Boston, 1994, p. 276 - 290.
- [7] S. Sinel'shchikov and L. Vaksman, *Hidden symmetry of the differential calculus on the quantum matrix space*, J. Phys. A., Math. Gen., **30** (1997), L23 – L26 (see also this volume).
- [8] S. Sinel'shchikov and L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, Mathematical Physics, Analysis, and Geometry, **1**, No 1, 1998, 75 – 100; q-alg/9703005.
- [9] S. Sinel'shchikov, D. Shklyarov, and L. Vaksman. *On function theory in quantum disc: integral representations*, math.QA/9808015.
- [10] L. L. Vaksman, *Integral intertwining operators and quantum homogeneous spaces*, Theoretical and Mathematical Physics, **105** (1995), No 3, 355 – 363.

q-ANALOGUES OF SOME BOUNDED SYMMETRIC DOMAINS

D. L. Shklyarov S. D. Sinel'shchikov L. L. Vaksman

Institute for Low Temperature Physics & Engineering
National Academy of Sciences of Ukraine,
Kharkov 61103, Ukraine

Abstract. We study q -analogues of matrix balls. A description of algebras of finite functions in the quantum matrix balls, an explicit form for the invariant integral in the space of finite functions, q -analogues for the weighted Bergman spaces, together with an explicit formula for the corresponding Bergman kernel, are presented.

1 Introduction

Hermitian symmetric spaces of non-compact type constitute one of the most important classes of homogeneous symmetric spaces. A well known result by Harish-Chandra claims that any such space can be realized as a bounded symmetric domain in a complex vector space V (via the so-called Harish-Chandra embedding).

Irreducible bounded symmetric domains were classified by E. Cartan. They are among the important subjects in Lie theory, geometry and function theory.

The first step in studying q -analogues of irreducible bounded symmetric domains was made in [2]. This work provides a q -analogue for the Harish-Chandra embedding and, in particular, a construction for q -analogues of the polynomial algebra and the differential calculus on V .

Our subject is the simplest class among those q -analogues, the quantum matrix balls. In the classical case $q = 1$ the corresponding vector space V is the space $\text{Mat}_{m,n}$ of rectangle complex matrices, and the matrix ball is defined as

$$\mathbb{U} = \{\mathbf{z} \in \text{Mat}_{m,n} | \mathbf{z}\mathbf{z}^* < 1\}.$$

This ball is a homogeneous space of the group $SU_{n,m}$.

Proofs of all presented results and also some results concerning the special case of quantum disc ($n = m = 1$) can be found in our electronic preprints (<http://xxx.lanl.gov/>).

2 Polynomials and finite functions in the quantum matrix ball

Everywhere in the sequel $q \in (0, 1)$, $m, n \in \mathbb{N}$, $m \leq n$, $N = m + n$. We use the standard notation $su_{n,m}$ for the Lie algebra of the group $SU_{n,m}$.

The Hopf algebra $U_q \mathfrak{sl}_N$ is determined by its generators $\{E_i, F_i, K_i^{\pm 1}\}_{i=1, \dots, N-1}$ and the well known Drinfeld-Jimbo relations [1]. Equip $U_q \mathfrak{sl}_N$ with the involution defined on the

This lecture has been delivered at the 8-th International Colloquium 'Quantum groups and Integrable Systems' held in Prague, June 1999; published in Czechoslovak Journal of Physics **50** (2000), No 1, 175 – 180

generators $K_j^{\pm 1}$, E_j , F_j , $j = 1, \dots, N-1$ by

$$(K_j^{\pm 1})^* = K_j^{\pm 1}, \quad E_j^* = \begin{cases} K_j F_j, & j \neq n \\ -K_j F_j, & j = n \end{cases}, \quad F_j^* = \begin{cases} E_j K_j^{-1}, & j \neq n \\ -E_j K_j^{-1}, & j = n \end{cases}.$$

The Hopf $*$ -algebra $U_q \mathfrak{su}_{n,m} = (U_q \mathfrak{sl}_N, *)$ arising this way is a q -analogue of the Hopf algebra $U \mathfrak{su}_{n,m}$.

Remind some well known definitions. An algebra F is said to be an A -module algebra if it is a module over a Hopf algebra A , the unit of F is an invariant and the multiplication $F \otimes F \rightarrow F$, $f_1 \otimes f_2 \mapsto f_1 \cdot f_2$, is a morphism of A -modules. In the case of a $*$ -algebra F and a Hopf- $*$ -algebra A , there is an additional requirement that the involutions agree as follows:

$$(af)^* = (S(a))^* f^*, \quad a \in A, f \in F, \quad (2.1)$$

with $S : A \rightarrow A$ being the antipode of A .

In [2] a $U_q \mathfrak{su}_{n,m}$ -module algebra $\text{Pol}(\text{Mat}_{m,n})_q$ and its $U_q \mathfrak{sl}_N$ -module subalgebra $\mathbb{C}[\text{Mat}_{m,n}]_q$ were introduced (the notation $\mathfrak{g}_{-1}$ was used in [2] instead of $\text{Mat}_{m,n}$). These algebras are q -analogues of polynomial algebras in the vector spaces $\text{Mat}_{m,n}$. We present below a description of these algebras in terms of generators and relations, together with explicit formulae for the $U_q \mathfrak{sl}_N$ -action in $\mathbb{C}[\text{Mat}_{m,n}]_q$ (corresponding explicit formulae for $U_q \mathfrak{su}_{n,m}$ -action in $\text{Pol}(\text{Mat}_{m,n})_q$ can be produced via (2.1)).

With the definitions of [2] as a background, one can prove the following two propositions.

Proposition 2.1 *There exists a unique family $\{z_a^\alpha\}_{a=1,\dots,n;\alpha=1,\dots,m}$ of elements of the $U_q \mathfrak{sl}_N$ -module algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$ such that for all $a = 1, \dots, n; \alpha = 1, \dots, m$*

$$H_n z_a^\alpha = \begin{cases} 2z_a^\alpha, & a = n \ \& \ \alpha = m \\ z_a^\alpha, & a = n \ \& \ \alpha \neq m \quad \text{or} \quad a \neq n \ \& \ \alpha = m \\ 0, & \text{otherwise} \end{cases}, \quad (2.2)$$

$$F_n z_a^\alpha = q^{1/2} \cdot \begin{cases} 1, & a = n \ \& \ \alpha = m \\ 0, & \text{otherwise} \end{cases}, \quad (2.3)$$

$$E_n z_a^\alpha = -q^{1/2} \cdot \begin{cases} q^{-1} z_a^m z_n^\alpha, & a \neq n \ \& \ \alpha \neq m \\ (z_n^m)^2, & a = n \ \& \ \alpha = m \\ z_n^m z_a^\alpha, & \text{otherwise} \end{cases}, \quad (2.4)$$

and with $k \neq n$

$$H_k z_a^\alpha = \begin{cases} z_a^\alpha, & k < n \ \& \ a = k \quad \text{or} \quad k > n \ \& \ \alpha = N - k \\ -z_a^\alpha, & k < n \ \& \ a = k + 1 \quad \text{or} \quad k > n \ \& \ \alpha = N - k + 1 \\ 0, & \text{otherwise} \end{cases}, \quad (2.5)$$

$$F_k z_a^\alpha = q^{1/2} \cdot \begin{cases} z_{a+1}^\alpha, & k < n \ \& \ a = k \\ z_{a+1}^\alpha, & k > n \ \& \ \alpha = N - k \\ 0, & \text{otherwise} \end{cases}, \quad (2.6)$$

$$E_k z_a^\alpha = q^{-1/2} \cdot \begin{cases} z_{a-1}^\alpha, & k < n \ \& \ a = k + 1 \\ z_{a-1}^\alpha, & k > n \ \& \ \alpha = N - k + 1 \\ 0, & \text{otherwise} \end{cases}. \quad (2.7)$$

REMARK. The elements $\{H_j\}_{j=1,\dots,N-1}$ and $\{K_j\}_{j=1,\dots,N-1}$ are related as follows:

$$K_j = q^{H_j}$$

(the exact definition of $\{H_j\}$ one can find in [2]).

Proposition 2.2 $\{z_a^\alpha\}_{a=1,\dots,n;\alpha=1,\dots,m}$ generate $\mathbb{C}[\text{Mat}_{m,n}]_q$ as an algebra and $\text{Pol}(\text{Mat}_{m,n})_q$ as a $*$ -algebra. The complete list of relations is as follows:

$$z_a^\alpha z_b^\beta = \begin{cases} q z_b^\beta z_a^\alpha & , \quad a = b \ \& \ \alpha < \beta \quad \text{or} \quad a < b \ \& \ \alpha = \beta \\ z_b^\beta z_a^\alpha & , \quad a < b \ \& \ \alpha > \beta \\ z_b^\beta z_a^\alpha + (q - q^{-1}) z_a^\beta z_b^\alpha & , \quad a < b \ \& \ \alpha < \beta \end{cases}, \quad (2.8)$$

$$(z_b^\beta)^* z_a^\alpha = q^2 \cdot \sum_{a',b'=1}^n \sum_{\alpha',\beta'=1}^m R(b, a, b', a') R(\beta, \alpha, \beta', \alpha) \cdot z_{a'}^{\alpha'} (z_{b'}^{\beta'})^* + (1 - q^2) \delta_{ab} \delta^{\alpha\beta}, \quad (2.9)$$

with δ_{ab} , $\delta^{\alpha\beta}$ being the Kronecker symbols and

$$R(b, a, b', a') = \begin{cases} q^{-1} & , \quad a \neq b \ \& \ b = b' \ \& \ a = a' \\ 1 & , \quad a = b = a' = b' \\ -(q^{-2} - 1) & , \quad a = b \ \& \ a' = b' \ \& \ a' > a \\ 0 & , \quad \text{otherwise} \end{cases}. \quad (2.10)$$

EXAMPLE. In the simplest case $m = n = 1$ the relations presented above describe a very well known $U_q \mathfrak{su}_{1,1}$ -module algebra

$$z^* z = q^2 z z^* + 1 - q^2. \quad (2.11)$$

Consider the $*$ -algebra $\text{Fun}(\mathbb{U})_q \supset \text{Pol}(\text{Mat}_{m,n})_q$ derived from $\text{Pol}(\text{Mat}_{m,n})_q$ by adding a generator f_0 such that

$$f_0 = f_0^2 = f_0^*, \quad (z_a^\alpha)^* f_0 = f_0 z_a^\alpha = 0, \quad a = 1, \dots, n; \quad \alpha = 1, \dots, m. \quad (2.12)$$

(Relations (2.12) allow one to treat f_0 as a q -analogue of the function that equal to 1 in the center of the ball and equal to 0 in other points.)

Proposition 2.3 *There exists a unique extension of the structure of a $U_q \mathfrak{su}_{n,m}$ -module algebra from $\text{Pol}(\text{Mat}_{m,n})_q$ onto $\text{Fun}(\mathbb{U})_q$ such that*

$$H_n f_0 = 0, \quad F_n f_0 = -\frac{q^{1/2}}{q^{-2} - 1} f_0 \cdot (z_n^m)^*, \quad E_n f_0 = -\frac{q^{1/2}}{1 - q^2} z_n^m \cdot f_0 \quad (2.13)$$

and with $k \neq n$

$$H_k f_0 = F_k f_0 = E_k f_0 = 0. \quad (2.14)$$

The two-sided ideal $D(\mathbb{U})_q \stackrel{\text{def}}{=} \text{Fun}(\mathbb{U})_q f_0 \text{Fun}(\mathbb{U})_q$ is a $U_q \mathfrak{su}_{n,m}$ -module algebra. Its elements will be called the finite functions in the quantum matrix ball.

3 Invariant integral

It is well known that in the classical case $q = 1$ the positive $SU_{n,m}$ -invariant integral could not be defined on the polynomial algebra. However, it is well defined on the space of finite smooth functions. These observations are still applicable in the quantum case.

Consider the representation T of $\text{Fun}(\mathbb{U})_q$ in the space $\mathcal{H} = \text{Fun}(\mathbb{U})_q f_0 = \text{Pol}(\text{Mat}_{m,n})_q f_0$:

$$T(f)\psi = f\psi, \quad f \in \text{Fun}(\mathbb{U})_q, \quad \psi \in \mathcal{H}. \quad (3.1)$$

REMARK. It can be shown that there exists a unique positive scalar product in $\mathcal{H}$ such that $(f_0, f_0) = 1$, and

$$(T(f)\psi_1, \psi_2) = (\psi_1, T(f^*)\psi_2), \quad f \in \text{Fun}(\mathbb{U})_q, \quad \psi_1, \psi_2 \in \mathcal{H}. \quad (3.2)$$

Moreover one can prove that the $*$ -algebra $\text{Pol}(\text{Mat}_{m,n})_q$ admits a unique up to unitary equivalence *faithful* irreducible $*$ -representation by *bounded* operators in a Hilbert space. This $*$ -representation can be produced via extending the operators $T(f)$, $f \in \text{Pol}(\text{Mat}_{m,n})_q$, onto the completion of the pre-Hilbert space $\mathcal{H}$.

Remind the notation $U_q \mathfrak{b}_+$ for the subalgebra of $U_q \mathfrak{sl}_N$ generated by the elements $\{E_i, K_i^{\pm 1}\}_{i=1, \dots, N-1}$. Obviously,

$$U_q \mathfrak{b}_+ \mathcal{H} \subset \mathcal{H},$$

and thus we obtain the representation Γ of the algebra $U_q \mathfrak{b}_+$ in $\mathcal{H}$. Let also

$$\check{\rho} = \frac{1}{2} \sum_{j=1}^{N-1} j(N-j) H_j. \quad (3.3)$$

Proposition 3.1 *The linear functional*

$$\int_{\mathbb{U}_q} f d\nu = \text{tr}(T(f)\Gamma(q^{-2\check{\rho}})), \quad f \in D(\mathbb{U})_q, \quad (3.4)$$

is well defined, $U_q \mathfrak{su}_{n,m}$ -invariant and positive (i.e. $\int_{\mathbb{U}_q} f^ f d\nu > 0$ for $f \neq 0$).*

4 Weighted Bergman spaces and Bergman kernels

Our intention is to produce q -analogues of weighted Bergman spaces. In the case $q = 1$ one has

$$\det(1 - \mathbf{z}\mathbf{z}^*) = 1 + \sum_{k=1}^m (-1)^k \mathbf{z}^{\wedge k} \mathbf{z}^{*\wedge k}, \quad (4.1)$$

with $\mathbf{z}^{\wedge k}$, $\mathbf{z}^{*\wedge k}$ being the "exterior powers" of the matrices $\mathbf{z}$, $\mathbf{z}^*$, that is, matrices formed by the minors of order k .

Let $1 \leq \alpha_1 < \alpha_2 < \dots < \alpha_k \leq m$, $1 \leq a_1 < a_2 < \dots < a_k \leq n$. Introduce q -analogues of minors for the matrix $\mathbf{z}$:

$$\mathbf{z}^{\wedge k}_{\{a_1, a_2, \dots, a_k\}}^{\{\alpha_1, \alpha_2, \dots, \alpha_k\}} = \sum_{s \in S_k} (-q)^{l(s)} z_{a_1}^{\alpha_{s(1)}} z_{a_2}^{\alpha_{s(2)}} \dots z_{a_k}^{\alpha_{s(k)}}, \quad (4.2)$$

with $l(s) = \text{card}\{(i, j) \mid i < j \ \& \ s(i) > s(j)\}$ being the length of the permutation s .

The q-analogue $y \in \text{Pol}(\text{Mat}_{m,n})_q$ for the polynomial $\det(1 - \mathbf{z}\mathbf{z}^*)$ is defined by

$$y = 1 + \sum_{k=1}^m (-1)^k \sum_{\{J' \mid \text{card}(J')=k\}} \sum_{\{J'' \mid \text{card}(J'')=k\}} \mathbf{z}^{\wedge^k_{J''}} \cdot \left(\mathbf{z}^{\wedge^k_{J'}} \right)^*. \quad (4.3)$$

Let $\lambda > m + n - 1$. Now one can define the integral with weight y^λ as follows:

$$\int_{\mathbb{U}_q} f d\nu_\lambda \stackrel{\text{def}}{=} C(\lambda) \int_{\mathbb{U}_q} f y^\lambda d\nu, \quad f \in D(\mathbb{U})_q, \quad (4.4)$$

where $C(\lambda) = \prod_{j=0}^{n-1} \prod_{k=0}^{m-1} (1 - q^{2(\lambda+1-N)} q^{2(j+k)})$ provides $\int_{\mathbb{U}_q} 1 d\nu_\lambda = 1$.

The Hilbert space $L^2(d\nu_\lambda)_q$ is defined as a completion of the space $D(\mathbb{U})_q$ of finite functions with respect to the norm $\|f\|_\lambda = \left(\int_{\mathbb{U}_q} f^* f d\nu_\lambda \right)^{1/2}$. The closure $L^2_a(d\nu_\lambda)_q$ in $L^2(d\nu_\lambda)_q$ of the algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$ will be called a weighted Bergman space.

Consider the orthogonal projection P_λ in $L^2(d\nu_\lambda)_q$ onto the weighted Bergman space $L^2_a(d\nu_\lambda)_q$. Our goal here is to show that P_λ could be written as an integral operator

$$P_\lambda f = \int_{\mathbb{U}_q} K_\lambda(\mathbf{z}, \boldsymbol{\zeta}^*) f(\boldsymbol{\zeta}) d\nu_\lambda(\boldsymbol{\zeta}), \quad f \in D(\mathbb{U})_q. \quad (4.5)$$

The main intention of this section is to introduce the algebra $\mathbb{C}[[\text{Mat}_{m,n} \times \overline{\text{Mat}}_{m,n}]]_q$ of kernels of integral operators and to determine an explicit form of the Bergman kernel $K_\lambda \in \mathbb{C}[[\text{Mat}_{m,n} \times \overline{\text{Mat}}_{m,n}]]_q$.

Introduce the notation

$$h_i = \sum_{\substack{J' \subset \{1,2,\dots,m\} \\ \text{card}(J')=i}} \sum_{\substack{J'' \subset \{1,2,\dots,n\} \\ \text{card}(J'')=i}} \mathbf{z}^{\wedge^i_{J''}} \otimes \left(\mathbf{z}^{\wedge^i_{J'}} \right)^*. \quad (4.6)$$

Let $\mathbb{C}[\overline{\text{Mat}}_{m,n}]_q \subset \text{Pol}(\text{Mat}_{m,n})_q$ be the unital subalgebra generated by $(z_a^\alpha)^*$, $a = 1, 2, \dots, n$, $\alpha = 1, 2, \dots, m$, and $\mathbb{C}[\text{Mat}_{m,n}]_q^{\text{op}}$ the algebra which **differs from** $\mathbb{C}[\text{Mat}_{m,n}]_q$ **by a replacement of its multiplication law to the opposite one**. The tensor product algebra $\mathbb{C}[\text{Mat}_{m,n}]_q^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{m,n}]_q$ will be called an algebra of polynomial kernels. It is possible to show that in this algebra $h_i h_j = h_j h_i$ for all $i, j = 1, 2, \dots, m$.

We follow [2] in equipping $\text{Pol}(\text{Mat}_{m,n})_q$ with a $\mathbb{Z}$ -gradation: $\deg(z_a^\alpha) = 1$, $\deg((z_a^\alpha)^*) = -1$, $a = 1, 2, \dots, n$, $\alpha = 1, 2, \dots, m$. In this context one has:

$$\mathbb{C}[\text{Mat}_{m,n}]_q^{\text{op}} = \bigoplus_{i=0}^{\infty} \mathbb{C}[\text{Mat}_{m,n}]_{q,i}^{\text{op}}, \quad \mathbb{C}[\overline{\text{Mat}}_{m,n}]_q = \bigoplus_{j=0}^{\infty} \mathbb{C}[\overline{\text{Mat}}_{m,n}]_{q,-j}, \quad (4.7)$$

$$\mathbb{C}[\text{Mat}_{m,n}]_q^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{m,n}]_q = \bigoplus_{i,j=0}^{\infty} \mathbb{C}[\text{Mat}_{m,n}]_{q,i}^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{m,n}]_{q,-j}. \quad (4.8)$$

The kernel algebra $\mathbb{C}[[\text{Mat}_{m,n} \times \overline{\text{Mat}}_{m,n}]]_q$ will stand for a completion of $\mathbb{C}[\text{Mat}_{m,n}]_q^{\text{op}} \otimes \mathbb{C}[\overline{\text{Mat}}_{m,n}]_q$ in the topology associated to the gradation in (4.8).

Proposition 4.1 *Let K_λ be an element of the algebra $\mathbb{C}[[\text{Mat}_{m,n} \times \overline{\text{Mat}}_{m,n}]]_q$ defined by*

$$K_\lambda = \prod_{j=0}^{\infty} \left(1 + \sum_{i=1}^m (-q^{2(\lambda+j)})^i h_i \right) \cdot \prod_{j=0}^{\infty} \left(1 + \sum_{i=1}^m (-q^{2j})^i h_i \right)^{-1} \quad (4.9)$$

Then (4.5) holds.

REMARK. A q -analogue of the ordinary Bergman kernel for the matrix ball is derivable from (4.9) by a substitution $\lambda = m + n$:

$$K = \prod_{j=0}^{m+n-1} \left(1 + \sum_{i=1}^m (-q^{2j})^i h_i \right)^{-1} \xrightarrow{q \rightarrow 1} (\det(1 - \mathbf{z} \cdot \boldsymbol{\zeta}^*))^{-(m+n)}.$$

References

- [1] M. Rosso: in Séminaire BOURBAKI, **744** (1991) 443 – 483.
- [2] S. Sinel'shchikov, L. Vaksman: Mathematical Physics, Analysis and Geometry **1** (1998) 75 – 100; q-alg/9703005.

Notes of the Editor

The proofs of the results announced in this work can be found in [1, 2]. These results acquired a further development in [3].

References

- [1] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035.
- [2] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum matrix ball: The Bergman kernel*, math.QA/9909036.
- [3] L. Vaksman, *Quantum matrix ball: the Cauchy-Szegö kernel and the Shilov boundary*, Matematicheskaya Fizika. Analiz. Geometriya, **8** (2001), No 4, 366 – 384; math.QA/0101179.

ON A q -ANALOGUE OF THE FOCK INNER PRODUCT

D.Shklyarov

Institute for Low Temperature Physics & Engineering
National Academy of Sciences of Ukraine,
Kharkov 61103, Ukraine

1 Introduction

Let $\text{Mat}_{m,n}$ be the space of complex $n \times m$ matrices. We denote by z_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$ the standard coordinate functions on $\text{Mat}_{m,n}$ given by the matrix entries. Let $\mathbb{C}[\text{Mat}_{m,n}]$ stands for the space of polynomials on $\text{Mat}_{m,n}$ (i. e. polynomials in z_a^α 's).

We recall that the Fock inner product in the space $\mathbb{C}[\text{Mat}_{m,n}]$ is defined by

$$(P, Q)_F = \int_{\text{Mat}_{m,n}} P(\mathbb{Z}) \overline{Q(\mathbb{Z})} e^{-\text{tr}(\mathbb{Z}\mathbb{Z}^*)} d\mathbb{Z} \quad (1.1)$$

with $\mathbb{Z} = (z_a^\alpha)$ and $d\mathbb{Z}$ being the Lebesgue measure on $\text{Mat}_{m,n}$ such that

$$(1, 1)_F = 1.$$

The inner product possesses the following remarkable property

$$\left(\frac{\partial P}{\partial z_a^\alpha}, Q \right)_F = (P, z_a^\alpha Q)_F \quad \forall a, \alpha, \quad (1.2)$$

which allows us to rewrite the product in the differential form

$$(P, Q)_F = \partial_P(\overline{Q})(0) \quad (1.3)$$

where ∂_P stands for the differential operator with constant coefficients derived from the polynomial P by substituting $z_a^\alpha \rightarrow \frac{\partial}{\partial z_a^\alpha}$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$, and $\overline{Q}$ stands for the polynomial derived from Q by changing the coefficients in the monomial basis to the complex conjugate ones.

At first glance, there is no necessity to involve the matrix space $\text{Mat}_{m,n}$ into the definition of the Fock inner product since the definition depends on the dimension $m \cdot n$ of $\text{Mat}_{m,n}$ only (indeed, $\text{tr}(\mathbb{Z}\mathbb{Z}^*)$ coincides with the usual Euclidean norm under the natural identification $\text{Mat}_{m,n} \cong \mathbb{C}^{mn}$). Our motivation comes from the theory of bounded symmetric domains (see, for instance, [1]). Suppose $D \subset \mathbb{C}^d$ is a bounded symmetric domain. One associates to D a Euclidean norm $\|\cdot\|$ in $\mathbb{C}^d$ and then the Fock inner product in the space of polynomials on $\mathbb{C}^d$:

$$(P, Q)_F = \int_{\mathbb{C}^d} P(\mathbf{z}) \overline{Q(\mathbf{z})} e^{-\|\mathbf{z}\|^2} d\mathbf{z}$$

with $\mathbf{z} = (z_1, z_2, \dots, z_d)$ and $d\mathbf{z}$ being the Lebesgue measure on $\mathbb{C}^d$ normalized by the same condition as in (1.1). Let G be the group of biholomorphic automorphisms of D and $K \subset G$ the stationary subgroup of the center $0 \in D$. Then K is a compact subgroup in $GL(d)$, and the Fock inner product is invariant under the induced action of K in the space of polynomials on $\mathbb{C}^d$. The same $\mathbb{C}^d$ may contain different bounded symmetric domains, and even if the associated Fock inner products coincide, the choice of a particular bounded symmetric domain indicates the symmetry group of the inner product. For example, by using the notation $\text{Mat}_{m,n}$ we indicate the connection of the inner product (1.1) to the matrix unit ball $D_{m,n} = \{\mathbf{Z} | \mathbf{Z}\mathbf{Z}^* < I\} \subset \text{Mat}_{m,n}$. The ball, we recall, is a homogeneous space of the group $SU(n, m)$, and the stationary subgroup of the zero matrix 0 is $S(U(n) \times U(m))$. The action of the latter group in $\mathbb{C}[\text{Mat}_{m,n}]$ is described explicitly by

$$P^g(\mathbf{Z}) = P(\bar{g}_2^t \mathbf{Z} g_1), \quad g = (g_1, g_2) \in S(U(n) \times U(m)),$$

and the invariance of the Fock inner product is written as follows

$$(P^g, Q^g)_F = (P, Q)_F, \quad \forall g \in S(U(n) \times U(m)). \quad (1.4)$$

It worth noting that easy computability and the invariance of the Fock inner product make it extremely useful in function theory on bounded symmetric domains.

We turn now to the subject of the present paper. It is known that the algebra $\mathbb{C}[\text{Mat}_{m,n}]$ has a non-commutative counterpart $\mathbb{C}[\text{Mat}_{m,n}]_q$ studied in quantum group theory. The latter algebra is the unital algebra given by its generators z_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, and the following relations

$$z_a^\alpha z_b^\beta = \begin{cases} q z_b^\beta z_a^\alpha & , \quad a = b \ \& \ \alpha < \beta \quad \text{or} \quad a < b \ \& \ \alpha = \beta \\ z_b^\beta z_a^\alpha & , \quad a < b \ \& \ \alpha > \beta \\ z_b^\beta z_a^\alpha + (q - q^{-1}) z_a^\beta z_b^\alpha & , \quad a < b \ \& \ \alpha < \beta \end{cases}, \quad (1.5)$$

Throughout the paper q is supposed to be a number from the interval $(0; 1)$ and the ground field is $\mathbb{C}$.

Around five years ago L.Vaksman and S.Sincl'shchikov [9] constructed certain analogs of bounded symmetric domains in framework of quantum group theory. Namely, they associated to a bounded symmetric domain D in a complex vector space V certain non-commutative algebras $\mathbb{C}[V]_q$, $\text{Pol}(V)_q$, which they treated as the algebras of holomorphic resp. arbitrary polynomials on the *quantum* vector space V . The algebras of continuous or finite functions on the *quantum* bounded symmetric domain D are derived then from $\text{Pol}(V)_q$ via some completion procedure. Further investigation has shown [4, 6] that in the case of the matrix unit ball (i.e. in the case $V = \text{Mat}_{m,n}$) the corresponding algebra of holomorphic polynomials from [9] is just the algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$.

It turned out [4, 5] that many constructions and problems of classical function theory in bounded symmetric domains admit natural generalization to the quantum setting. Thus, it is reasonable to expect that there should be an appropriate q -analog of the Fock inner product and that it will serve the same purposes as its classical counterpart does.

The aim of the present paper is to present a candidate for that q -analog in the case of the quantum matrix space $\text{Mat}_{m,n}$.

The structure of the paper is as follows. In the next section we formulate our main results. Sections 3, 4, and 5 contain an auxiliary material concerning quantum group symmetry. Specifically, in section 3 we discuss the notions of module algebras and modules over module algebras. In sections 4,5 we describe an action of a quantum group in algebras and spaces we deal with. The proof of the main results is given in section 6.

The author is grateful to L. Vaksman for numerous interesting discussions of the results.

2 Statement of main results

The equality (1.2) served for us as a guide when we were looking for a q -analog of the Fock inner product. The point is that there are natural quantum analogs of the partial derivatives $\frac{\partial}{\partial z_a^\alpha}$. They are constructed by using a first order differential calculus over the algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$. Let us recall a definition of that differential calculus. Let $\Lambda^1(\text{Mat}_{m,n})_q$ be the $\mathbb{C}[\text{Mat}_{m,n}]_q$ -bimodule given by its generators dz_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, and the relations

$$z_b^\beta dz_a^\alpha = \sum_{\alpha', \beta'=1}^m \sum_{a', b'=1}^n R_{\beta\alpha}^{\beta'\alpha'} R_{ba}^{b'a'} dz_{a'}^{\alpha'} \cdot z_{b'}^{\beta'}, \quad (2.1)$$

with

$$R_{ba}^{b'a'} = \begin{cases} q^{-1} & , \quad a = b = a' = b' \\ 1 & , \quad a \neq b \quad \& \quad a = a' \quad \& \quad b = b' \\ q^{-1} - q & , \quad a < b \quad \& \quad a = b' \quad \& \quad b = a' \\ 0 & , \quad \text{otherwise} \end{cases} \quad (2.2)$$

The map $d : z_a^\alpha \mapsto dz_a^\alpha$ can be extended up to a linear operator $d : \mathbb{C}[\text{Mat}_{m,n}]_q \rightarrow \Lambda^1(\text{Mat}_{m,n})_q$ satisfying the Leibnitz rule. The pair $(\Lambda^1(\text{Mat}_{m,n})_q, d)$ is the first order differential calculus over $\mathbb{C}[\text{Mat}_{m,n}]_q$. It worth noting that this first order differential calculus coincides with certain 'canonical' one, defined in [9].

The q -analogs of the partial derivatives may be defined now via the differential d as follows:

$$df = \sum_{a=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial z_a^\alpha} dz_a^\alpha, \quad f \in \mathbb{C}[\text{Mat}_{m,n}]_q.$$

It is reasonable to rise the question about existence of an inner product in $\mathbb{C}[\text{Mat}_{m,n}]_q$ with the property (1.2) where the classical partial derivatives are replaced by the quantum ones. If such an inner product existed we would have the algebra *antihomomorphism*

$$\mathbb{C}[\text{Mat}_{m,n}]_q \rightarrow \mathbb{C}[\partial]_q, \quad z_a^\alpha \mapsto \frac{\partial}{\partial z_a^\alpha}, \quad \forall a, \alpha \quad (2.3)$$

where $\mathbb{C}[\partial]_q$ stands for the algebra of quantum differential operators with constant coefficients (i.e., the unital algebra of linear operators in $\mathbb{C}[\text{Mat}_{m,n}]_q$ generated by the quantum partial derivatives). However, in reality (2.3) is the algebra *homomorphism*, that is, the partial derivatives satisfy the same relations as the generators of $\mathbb{C}[\text{Mat}_{m,n}]_q$ do. To prove this, one should use the higher order differential calculus over $\mathbb{C}[\text{Mat}_{m,n}]_q$ associated to

the first order one [8, 9, 4]. Specifically, let $\Lambda(\text{Mat}_{m,n})_q$ be the unital algebra given by its generators z_a^α , dz_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, satisfying (1.5), (2.1), and the relations

$$dz_b^\beta dz_a^\alpha = - \sum_{\alpha', \beta'=1}^m \sum_{a', b'=1}^n R_{\beta\alpha}^{\beta'\alpha'} R_{ba}^{b'a'} dz_{a'}^{\alpha'} \cdot dz_{b'}^{\beta'}, \quad (2.4)$$

with R defined by (2.2). The algebra admits the natural grading by degrees of differential forms. There exists a unique extension of the differential $d : \mathbb{C}[\text{Mat}_{m,n}]_q \rightarrow \Lambda^1(\text{Mat}_{m,n})_q$ to a linear operator $d : \Lambda(\text{Mat}_{m,n})_q \rightarrow \Lambda(\text{Mat}_{m,n})_q$ which satisfies the (graded) Leibnitz rule and the property $d^2 = 0$, which implies, in particular, $d : dz_a^\alpha \mapsto 0$ (for all a, α). The property means

$$\sum_{a, a'=1}^n \sum_{\alpha, \alpha'=1}^m \frac{\partial}{\partial z_a^\alpha} \frac{\partial}{\partial z_{a'}^{\alpha'}} (f) dz_a^\alpha dz_{a'}^{\alpha'} = 0, \quad \forall f \in \mathbb{C}[\text{Mat}_{m,n}]_q.$$

This equality and the relations (2.4) imply the desired commutation relations between the quantum partial derivatives:

$$\frac{\partial}{\partial z_a^\alpha} \frac{\partial}{\partial z_b^\beta} = \begin{cases} q \frac{\partial}{\partial z_b^\beta} \frac{\partial}{\partial z_a^\alpha} & , \quad a = b \text{ \& } \alpha < \beta \quad \text{or} \quad a < b \text{ \& } \alpha = \beta \\ \frac{\partial}{\partial z_b^\beta} \frac{\partial}{\partial z_a^\alpha} & , \quad a < b \text{ \& } \alpha > \beta \\ \frac{\partial}{\partial z_b^\beta} \frac{\partial}{\partial z_a^\alpha} + (q - q^{-1}) \frac{\partial}{\partial z_a^\alpha} \frac{\partial}{\partial z_b^\beta} & , \quad a < b \text{ \& } \alpha < \beta \end{cases}. \quad (2.5)$$

Thus we can't expect existence of an inner product in $\mathbb{C}[\text{Mat}_{m,n}]_q$ satisfying the property (1.2). However, the relations (2.5) suggest a way to overcome the problem: one can try to look for such an inner product which makes the operator $\frac{\partial}{\partial z_a^\alpha}$ conjugate to the operator of *right multiplication* by z_a^α in $\mathbb{C}[\text{Mat}_{m,n}]_q$. Such an inner product turned out to exist, and this observation is one of the main results of the paper:

Theorem 2.1 *There exists a unique inner product $(\cdot, \cdot)_F$ in $\mathbb{C}[\text{Mat}_{m,n}]_q$ satisfying the properties*

$$(1, 1)_F = 1, \quad (2.6)$$

$$\left(\frac{\partial P}{\partial z_a^\alpha}, Q \right)_F = (P, Q \cdot z_a^\alpha)_F \quad \forall a, \alpha. \quad (2.7)$$

Note that uniqueness follows immediately from the two properties since any partial derivative is an operator of degree -1 with respect to the natural $\mathbb{Z}_+$ -grading in $\mathbb{C}[\text{Mat}_{m,n}]_q$ by powers of monomials.

Our next result may be formulated as follows: the inner product $(\cdot, \cdot)_F$ is invariant with respect to an action of the *quantum group* $S(U(n) \times U(m))$ in $\mathbb{C}[\text{Mat}_{m,n}]_q$. To present a precise formulation, we need some preparation.

Let us recall the notion of the Drinfeld-Jimbo quantized universal enveloping algebra of $\mathfrak{sl}(k)$. Let (a_{ij}) be the Cartan matrix for $\mathfrak{sl}(k)$. The Hopf algebra $U_q \mathfrak{sl}(k)$ is determined by the generators E_i, F_i, K_i, K_i^{-1} , $i = 1, \dots, k-1$, and the relations

$$K_i K_j = K_j K_i, \quad K_i K_i^{-1} = K_i^{-1} K_i = 1, \quad K_i E_j = q^{a_{ij}} E_j K_i,$$

$$\begin{aligned}
K_i F_j &= q^{-aj} F_j K_i, & E_i F_j - F_j E_i &= \delta_{ij} (K_i - K_i^{-1}) / (q - q^{-1}) \\
E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0, & |i - j| &= 1 \\
F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0, & |i - j| &= 1 \\
[E_i, E_j] &= [F_i, F_j] = 0, & |i - j| &\neq 1.
\end{aligned} \tag{2.8}$$

The comultiplication Δ , the antipode S , and the counit ε are determined by

$$\Delta(E_i) = E_i \otimes 1 + K_i \otimes E_i, \quad \Delta(F_i) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \quad \Delta(K_i) = K_i \otimes K_i, \tag{2.9}$$

$$\begin{aligned}
S(E_i) &= -K_i^{-1} E_i, & S(F_i) &= -F_i K_i, & S(K_i) &= K_i^{-1}, \\
\varepsilon(E_i) &= \varepsilon(F_i) = 0, & \varepsilon(K_i) &= 1.
\end{aligned} \tag{2.10}$$

Let $U_q(\mathfrak{sl}(n) \oplus \mathfrak{sl}(m))$ stands for the Hopf algebra $U_q \mathfrak{sl}(n) \otimes U_q \mathfrak{sl}(m)$.

Recall the standard terminology. Let A be a Hopf algebra. An algebra F is said to be an A -module algebra if F carries a structure of A -module and multiplication in F agrees with the A -action (i. e. the multiplication $F \otimes F \rightarrow F$ is a morphism of A -modules).

The algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$ possesses the well known structure of $U_q(\mathfrak{sl}(n) \oplus \mathfrak{sl}(m))$ -module algebra:

$$K_i \otimes 1(z_a^\alpha) = \begin{cases} q z_a^\alpha, & a = i \\ q^{-1} z_a^\alpha, & a = i + 1 \\ z_a^\alpha, & \text{otherwise} \end{cases}, \quad 1 \otimes K_j(z_a^\alpha) = \begin{cases} q z_a^\alpha, & \alpha = m - j \\ q^{-1} z_a^\alpha, & \alpha = m - j + 1 \\ z_a^\alpha, & \text{otherwise} \end{cases}, \tag{2.11}$$

$$F_i \otimes 1(z_a^\alpha) = \begin{cases} q^{1/2} z_{a+1}^\alpha, & a = i \\ 0, & \text{otherwise} \end{cases}, \quad 1 \otimes F_j(z_a^\alpha) = \begin{cases} q^{1/2} z_a^{\alpha+1}, & \alpha = m - j \\ 0, & \text{otherwise} \end{cases}, \tag{2.12}$$

$$E_i \otimes 1(z_a^\alpha) = \begin{cases} q^{-1/2} z_{a-1}^\alpha, & a = i + 1 \\ 0, & \text{otherwise} \end{cases}, \quad 1 \otimes E_j(z_a^\alpha) = \begin{cases} q^{-1/2} z_a^{\alpha-1}, & \alpha = m - j + 1 \\ 0, & \text{otherwise} \end{cases}. \tag{2.13}$$

The Hopf algebra $U_q \mathfrak{sl}(n) \oplus \mathfrak{sl}(m)$ is derived from $U_q(\mathfrak{sl}(n) \oplus \mathfrak{sl}(m))$ by adding the generator K_0 , commuting with the other generators and satisfies the properties

$$\Delta(K_0) = K_0 \otimes K_0, \quad S(K_0) = K_0^{-1}, \quad \varepsilon(K_0) = 1. \tag{2.14}$$

Let us extend the $U_q(\mathfrak{sl}(n) \oplus \mathfrak{sl}(m))$ -module algebra structure in $\mathbb{C}[\text{Mat}_{m,n}]_q$ to a $U_q \mathfrak{sl}(n) \oplus \mathfrak{sl}(m)$ -module algebra structure as follows:

$$K_0(z_a^\alpha) = q^{n+m} z_a^\alpha \quad \forall a, \alpha. \tag{2.15}$$

The $*$ -Hopf algebra $U_q(\mathfrak{su}(n) \oplus \mathfrak{su}(m))$ is the pair $(U_q(\mathfrak{sl}(n) \oplus \mathfrak{sl}(m)), *)$ with $*$ being the involution in $U_q(\mathfrak{sl}(n) \oplus \mathfrak{sl}(m))$ given by

$$\begin{aligned}
(E_i \otimes 1)^* &= K_i F_i \otimes 1, & (F_i \otimes 1)^* &= E_i K_i^{-1} \otimes 1, & (K_i^{\pm 1} \otimes 1)^* &= K_i^{\pm 1} \otimes 1, \\
(1 \otimes E_j)^* &= 1 \otimes K_j F_j, & (1 \otimes F_j)^* &= 1 \otimes E_j K_j^{-1}, & (1 \otimes K_j^{\pm 1})^* &= 1 \otimes K_j^{\pm 1}.
\end{aligned}$$

This involution is extended to an involution in $U_q \mathfrak{sl}(n) \oplus \mathfrak{sl}(m)$ by setting $K_0^* = K_0$. The resulting $*$ -Hopf algebra $(U_q \mathfrak{sl}(n) \oplus \mathfrak{sl}(m), *)$ is denoted by $U_q \mathfrak{su}(n) \oplus \mathfrak{su}(m)$.

Now we are ready to formulate the invariance property of the q -Fock inner product.

Theorem 2.2 *The inner product $(\cdot, \cdot)_F$ is $U_q\mathfrak{sl}(\mathfrak{u}(n) \oplus \mathfrak{u}(m))$ -invariant, i.e.*

$$(\xi(P), Q)_F = (P, \xi^*(Q))_F \quad (2.16)$$

for all $P, Q \in \mathbb{C}[\text{Mat}_{m,n}]_q$ and $\xi \in U_q\mathfrak{sl}(\mathfrak{gl}(n) \oplus \mathfrak{gl}(m))$.

Our proof of the above two theorems is based on the following more or less explicit description of the q -Fock inner product.

Consider the unital involutive algebra $\mathcal{P}(m, n)_q$ with the generators z_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, satisfying (1.5) and the relations

$$(z_b^\beta)^* z_a^\alpha = \sum_{a', b'=1}^n \sum_{\alpha', \beta'=1}^m \widehat{R}_{ba}^{b'a'} \widehat{R}_{\beta\alpha}^{\beta'\alpha'} z_{a'}^{\alpha'} (z_{b'}^{\beta'})^* + \delta_{ab} \delta^{\alpha\beta}, \quad (2.17)$$

with δ_{ab} , $\delta^{\alpha\beta}$ being the Kronecker symbols and

$$\widehat{R}_{ba}^{b'a'} = \begin{cases} 1 & , \quad a \neq b \text{ \& } b = b' \text{ \& } a = a' \\ q & , \quad a = b = a' = b' \\ q - q^{-1} & , \quad a = b \text{ \& } a' = b' \text{ \& } a' > a \\ 0 & , \quad \text{otherwise} \end{cases} \quad (2.18)$$

Clearly, the algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$ is embedded into $\mathcal{P}(m, n)_q$.

It worth noting that the algebra $\mathcal{P}(1, n)_q$ is isomorphic to the well known twisted CCR-algebra introduced by W.Pusz and S.Woronowicz [3]. For arbitrary m and n the algebra is isomorphic to the algebra $\text{Pol}(\text{Mat}_{m,n})_q$ of polynomials on the quantum matrix space [4]. A precise definition of $\text{Pol}(\text{Mat}_{m,n})_q$ and a description of the isomorphism are to be found in Section 4.

Let us consider the $\mathcal{P}(m, n)_q$ -module H given by its unique generator e_{vac} and the relations $(z_a^\alpha)^* e_{vac} = 0$ for all a and α . The aforementioned isomorphism $\mathcal{P}(m, n)_q \cong \text{Pol}(\text{Mat}_{m,n})_q$ allows us to use results from [4, 6] to derive the following statements concerning $\mathcal{P}(m, n)_q$. First of all, the multiplication map $m : \mathcal{P}(m, n)_q \otimes \mathcal{P}(m, n)_q \rightarrow \mathcal{P}(m, n)_q$ induces the isomorphism of vector spaces $\mathbb{C}[\text{Mat}_{m,n}]_q \otimes \mathbb{C}[\overline{\text{Mat}}_{m,n}]_q \rightarrow \mathcal{P}(m, n)_q$ with $\mathbb{C}[\overline{\text{Mat}}_{m,n}]_q = \{f^* \mid f \in \mathbb{C}[\text{Mat}_{m,n}]_q\}$ (note that surjectivity is a simple consequence of the relations (2.17)). Thus

$$H = \mathbb{C}[\text{Mat}_{m,n}]_q e_{vac}. \quad (2.19)$$

Further, there exists a unique inner product in H so that

$$(1, 1) = 1, \quad (f e_1, e_2) = (e_1, f^* e_2) \quad (2.20)$$

for any $f \in \mathcal{P}(m, n)_q$ and $e_1, e_2 \in H$. The equality (2.19) allows us to regard the inner product as the one on $\mathbb{C}[\text{Mat}_{m,n}]_q$.

Theorem 2.3 *The inner product $(\cdot, \cdot)$ in $\mathbb{C}[\text{Mat}_{m,n}]_q$ coincides with $(\cdot, \cdot)_F$, that is*

$$\left(\frac{\partial P}{\partial z_a^\alpha}, Q \right) = (P, Q \cdot z_a^\alpha) \quad \forall a, \alpha. \quad (2.21)$$

Though we'll present a complete proof of the theorem, it still seems to be somewhat mysterious since it relates two, at first glance, different quantum analogs of the Weyl algebra: $\mathcal{P}(m, n)_q$ and the algebra generated by quantum partial derivatives and the operators of right multiplication by z_a^α 's (we'll denote the latter algebra by $A_{m,n}(\mathbb{C})_q$).

Let us explain very briefly a logic of the proof of the three theorems. Obviously, theorem 3 implies theorem 1. In section 6 we express the inner product $(\cdot, \cdot)$ through another one denoted by $\langle \cdot, \cdot \rangle$. The latter inner product is obviously invariant and this gives us theorem 2. Thus, the relation (2.21) is, in a sense, central in the paper, and most of section 6 is devoted to its proof.

3 Quantum symmetry

The aim of this section is to remind some general notions from quantum group theory.

In this section A denotes a Hopf algebra with the comultiplication Δ , the antipode S , and the counit ε . An algebra F is said to be an A -module algebra if F carries a structure of A -module and multiplication in F agrees with the A -action:

$$\xi(f_1 \cdot f_2) = \sum_j \xi'_j(f_1) \cdot \xi''_j(f_2), \quad f_1, f_2 \in F, \quad \xi \in A, \quad \sum_j \xi'_j \otimes \xi''_j = \Delta(\xi) \quad (3.1)$$

(i. e. the multiplication $F \otimes F \rightarrow F$ is a morphism of A -modules). If A or F have some additional structures, this definition includes some extra requirements. For example, if F is unital, one requires A -invariance of the unit:

$$\xi(1) = \varepsilon(\xi) \cdot 1, \quad \xi \in A.$$

In the case of a $*$ -algebra F and a $*$ -Hopf algebra A one imposes the requirement of agreement of the involutions:

$$(\xi(f))^* = S(\xi)^*(f^*), \quad \xi \in A, f \in F. \quad (3.2)$$

Some examples of module algebras naturally appear in representation theory and harmonic analysis.

Suppose a smooth manifold X is acted by a Lie group G . This induces an action of the Lie algebra $\mathfrak{g}$ of G in the space $C^\infty(X)$ by means of vector fields. In turn, the $\mathfrak{g}$ -action induces an action of the universal enveloping algebra $U\mathfrak{g}$ in $C^\infty(X)$ by means of differential operators. The usual Leibnitz rule means that $C^\infty(X)$ is a $U\mathfrak{g}$ -module algebra.

Another important example of a module algebra is the algebra of linear endomorphisms of a vector space, acted by a Hopf algebra. Let V be an A -module. Endow the space $\text{End}_{\mathbb{C}}(V)$ with a structure of A -module as follows:

$$\text{ad}\xi(T) = \sum_j \xi'_j \cdot T \cdot S(\xi''_j), \quad (3.3)$$

where $\xi \in A$, $T \in \text{End}_{\mathbb{C}}(V)$, $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$, S is the antipode of A , and the elements in the right-hand side are multiplied within the algebra $\text{End}_{\mathbb{C}}(V)$. It follows from elementary properties of Hopf algebras that this action of A makes $\text{End}_{\mathbb{C}}(V)$ into an A -module algebra.

Now turn to another important notion, namely, that of modules over module algebra. Let F be an A -module algebra and M be a left (right) F -module. Then M is said to be an A -module left (resp. right) F -module, if M carries a structure of A -module and

the multiplication $F \otimes M \rightarrow M$ (resp. $M \otimes F \rightarrow M$) agrees with the A -action, that is, $F \otimes M \rightarrow M$ (resp. $M \otimes F \rightarrow M$) is a morphism of A -modules.

Let again X be a smooth manifold acted by a Lie group G , and $T^*X \rightarrow X$ be the cotangent bundle. T^*X inherits a natural G -action. Then the space of smooth sections of the bundle (i.e. the space of 1-forms) is acted by the universal enveloping algebra $U\mathfrak{g}$, and the action agrees with multiplication by smooth functions. Thus, 1-forms constitute a $U\mathfrak{g}$ -module left (as well as right) $C^\infty(X)$ -module.

Let V be an A -module, and $\text{End}_{\mathbb{C}}(V)$ is given with a structure of A -module as in (3.3). It is straightforward that the natural action $T \otimes v \mapsto T(v)$ makes V into a A -module left $\text{End}_{\mathbb{C}}(V)$ -module.

Observe that the latter example is quite general. Indeed, let F be an A -module algebra and M an A -module left F -module. The multiplication $F \otimes M \rightarrow M$ induces the natural algebra homomorphism $F \rightarrow \text{End}_{\mathbb{C}}(M)$. It is easy to verify that this is a morphism of A -module algebras (here $\text{End}_{\mathbb{C}}(M)$ is viewed with the A -module structure given by (3.3)).

Finally, let us agree about the following useful notation. Suppose F is an algebra and an A -module. If, instead of (3.1), we have

$$\xi(f_1 \cdot f_2) = \sum_j \xi_j''(f_1) \cdot \xi_j'(f_2) \quad \forall f_1, f_2 \in F, \quad \xi \in A \quad (3.4)$$

(with $\sum_j \xi_j' \otimes \xi_j'' = \Delta(\xi)$) then we call F an A^{op} -module algebra. Similarly, suppose F is an A^{op} -module algebra and M is a left (right) F -module and an A -module. Then M is said to be an A^{op} -module left (resp. right) F -module, if the multiplication $F \otimes M \rightarrow M$ (resp. $M \otimes F \rightarrow M$) satisfies the property

$$\xi(f \cdot m) = \sum_j \xi_j''(f) \cdot \xi_j'(m) \quad \forall f \in F, \quad m \in M, \quad \xi \in A$$

(resp. $\xi(m \cdot f) = \sum_j \xi_j''(m) \cdot \xi_j'(f)$).

An important example of A^{op} -module algebras is constructed as follows. For an A -module V the space $\text{End}_{\mathbb{C}}(V)$ admits the following alternative structure of A -module:

$$\text{ad}'\xi(T) = \sum_j \xi_j'' \cdot T \cdot S^{-1}(\xi_j'). \quad (3.5)$$

This A -action makes $\text{End}_{\mathbb{C}}(V)$ into an A^{op} -module algebra, and the natural action $T \otimes v \mapsto T(v)$ makes V into an A^{op} -module left $\text{End}_{\mathbb{C}}(V)$ -module.

Some important examples to the aforementioned notions appear in the next sections.

4 Examples of quantum symmetry: functions in the quantum matrix ball

In this section we describe some non-trivial examples of a quantum group action in algebras. The algebras we deal with in this section are treated as the algebras of functions on certain quantum G -spaces.

The first example of a module algebra already appeared in section 2. There we described a well known structure $U_q\mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -module algebra structure in $\mathbb{C}[\text{Mat}_{m,n}]_q$. It is observed in [4] that this $U_q\mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -module algebra structure may be extended to a $U_q\mathfrak{sl}(n+m)$ -module one. First, we have to explain in what sense the latter structure extends the former one. The point is that there is a natural embedding of Hopf algebras $U_q\mathfrak{sl}(n) \oplus \mathfrak{gl}(m) \hookrightarrow U_q\mathfrak{sl}(n+m)$ determined by

$$\begin{aligned}
E_i \otimes 1 &\mapsto E_i, & F_i \otimes 1 &\mapsto F_i, & K_i^{\pm 1} \otimes 1 &\mapsto K_i^{\pm 1}, \\
1 \otimes E_j &\mapsto E_{n+j}, & 1 \otimes F_j &\mapsto F_{n+j}, & 1 \otimes K_j^{\pm 1} &\mapsto K_{n+j}^{\pm 1},
\end{aligned}$$

$$K_0 \mapsto K_n^{nm} \cdot \prod_{i=1}^{n-1} K_i^{mi} \cdot \prod_{j=1}^{m-1} K_{n+j}^{n(m-j)}.$$

We also recall one the notation $U_q \mathfrak{su}(n, m)$ for the $*$ -Hopf algebra $(U_q \mathfrak{sl}(n+m), *)$ where the involution is defined as follows

$$(K_j^{\pm 1})^* = K_j^{\pm 1}, \quad E_j^* = \begin{cases} K_j F_j, & j \neq n \\ -K_j F_j, & j = n \end{cases}, \quad F_j^* = \begin{cases} E_j K_j^{-1}, & j \neq n \\ -E_j K_j^{-1}, & j = n \end{cases}.$$

One verifies easily that the above embedding respects the involutions in $U_q \mathfrak{su}(n) \oplus \mathfrak{u}(m)$ and $U_q \mathfrak{su}(n, m)$. Let us agree to use the notation $U_q \mathfrak{sl}(\mathfrak{gl}(n) \oplus \mathfrak{gl}(m))$ ($U_q \mathfrak{su}(\mathfrak{u}(n) \oplus \mathfrak{u}(m))$) to denote the image of the above embedding, i.e. the corresponding Hopf (resp. $*$ -Hopf) subalgebra in $U_q \mathfrak{sl}(n+m)$ (resp. $U_q \mathfrak{su}(n, m)$).

It follows from (2.11) and (2.15) that

$$K_n z_a^\alpha = \begin{cases} q^2 z_a^\alpha, & a = n \text{ \& } \alpha = m \\ q z_a^\alpha, & a = n \text{ \& } \alpha \neq m \\ z_a^\alpha, & \text{otherwise} \end{cases} \quad \text{or} \quad a \neq n \text{ \& } \alpha = m. \quad (4.1)$$

To describe the $U_q \mathfrak{sl}(n+m)$ -module algebra structure in $\mathbb{C}[\text{Mat}_{m,n}]_q$ completely, we have to add to (2.11), (2.12), (2.13), and (4.1) formulae for the action of the generators E_n and F_n :

$$F_n z_a^\alpha = \begin{cases} q^{1/2}, & a = n \text{ \& } \alpha = m \\ 0, & \text{otherwise} \end{cases}, \quad (4.2)$$

$$E_n z_a^\alpha = \begin{cases} -q^{-1/2} z_a^m z_n^\alpha, & a \neq n \text{ \& } \alpha \neq m \\ -q^{1/2} (z_n^m)^2, & a = n \text{ \& } \alpha = m \\ -q^{1/2} z_n^m z_a^\alpha, & \text{otherwise} \end{cases}. \quad (4.3)$$

Let us say few words about the nature of the above $U_q \mathfrak{sl}(n+m)$ -action in $\mathbb{C}[\text{Mat}_{m,n}]_q$. Its classical counterpart admits the following description. As it is noted in Introduction, the vector space $\text{Mat}_{m,n}$ contains the so called matrix ball $D_{m,n} = \{\mathbf{Z} \mid \mathbf{Z}\mathbf{Z}^* < \mathbf{I}\}$. The group $SU(n, m)$ acts in $D_{m,n}$ via biholomorphic automorphisms. Thus, elements of the universal enveloping algebra $U \mathfrak{su}(n, m)$ (and hence of its complexification $U \mathfrak{sl}(n+m)$) act in the space of holomorphic functions in $D_{m,n}$ by means of differential operators. The differential operators have polynomial coefficients and, therefore, preserve the subspace of polynomials. This $U \mathfrak{sl}(n+m)$ -action is the 'classical limit' of the above $U_q \mathfrak{sl}(n+m)$ -action in $\mathbb{C}[\text{Mat}_{m,n}]_q$.

The algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$ is a particular case of some algebras constructed in [9] (see Introduction). Let us recall here the description of $\mathbb{C}[\text{Mat}_{m,n}]_q$ given in [9] (see also [4], [6]).

Let us consider the generalized Verma module $V(0)$ over $U_q \mathfrak{sl}(n+m)$, given by its generator $v(0)$ and the relations

$$E_i v(0) = 0, \quad K_i v(0) = v(0), \quad \forall i, \quad (4.4)$$

$$F_i v(0) = 0, \quad i \neq n. \quad (4.5)$$

Viewed as a $U_q \mathfrak{sl}(n) \oplus \mathfrak{sl}(m)$ -module, the space $V(0)$ splits into direct sum of its finite dimensional $U_q \mathfrak{sl}(n) \oplus \mathfrak{sl}(m)$ -submodules $V(0)_{-k}$, $k \in \mathbb{Z}_+$, with

$$V(0)_{-k} = \{v \in V(0) \mid K_0 v = q^{(n+m)k} v\} \quad (4.6)$$

(here, we recall, $K_0 = K_n^{nm} \cdot \prod_{i=1}^{n-1} K_i^{mi} \cdot \prod_{j=1}^{m-1} K_{n+j}^{n(m-j)}$). Denote by $V(0)^*$ the *graded* dual $U_q \mathfrak{sl}(n+m)$ -module: $V(0)^* = \bigoplus_{k \in \mathbb{Z}_+} (V(0)_{-k})^*$.

Let us equip the tensor product $V(0) \otimes V(0)$ with a $U_q \mathfrak{sl}(n+m)$ -module structure via the opposite comultiplication

$$\xi(v_1 \otimes v_2) = \sum_j \xi_j''(v_1) \otimes \xi_j'(v_2), \quad \xi \in U_q \mathfrak{sl}(n+m), \quad v_1, v_2 \in V(0). \quad (4.7)$$

The relations (4.4), (4.5) imply that the maps $v(0) \mapsto v(0) \otimes v(0)$, $v(0) \mapsto 1$ are extendable up to morphisms of $U_q \mathfrak{sl}(n+m)$ -modules $\Delta_- : V(0) \rightarrow V(0) \otimes V(0)$, $\varepsilon_- : V(0) \rightarrow \mathbb{C}$. It can be shown that Δ_- and ε_- make $V(0)$ into a coassociative coalgebra with a counit. Thus, the dual maps $m = (\Delta_-)^* : V(0)^* \otimes V(0)^* \rightarrow V(0)^*$, $1 = (\varepsilon_-)^* : \mathbb{C} \rightarrow V(0)^*$ make $V(0)^*$ into an associative unital algebra. Moreover, the product map m is a morphism of $U_q \mathfrak{sl}(n+m)$ -modules and the unit 1 is $U_q \mathfrak{sl}(n+m)$ -invariant, i.e. $V(0)^*$ is a $U_q \mathfrak{sl}(n+m)$ -module algebra. It turns out to be isomorphic to $\mathbb{C}[\text{Mat}_{m,n}]_q$ [4].

Remind the notation $\text{Pol}(\text{Mat}_{m,n})_q$ for the algebra of (not necessary holomorphic) polynomials on the quantum matrix space (see section 2). This algebra is the unital involutive algebra with the generators z_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, satisfying (1.5) and the relations

$$(z_b^\beta)^* z_a^\alpha = \sum_{a', b'=1}^n \sum_{\alpha', \beta'=1}^m \widehat{R}_{b,a}^{\beta', a'} \widehat{R}_{\beta, \alpha}^{\beta', \alpha'} z_{a'}^{\alpha'} (z_{b'}^{\beta'})^* + \delta_{ab} \delta^{\alpha\beta} \cdot (1 - q^2) \quad (4.8)$$

(the matrix R is given by (2.18)). As we stated in section 2, $\text{Pol}(\text{Mat}_{m,n})_q$ is isomorphic to the $*$ -algebra $\mathcal{P}(m, n)_q$: the isomorphism $J : \text{Pol}(\text{Mat}_{m,n})_q \rightarrow \mathcal{P}(m, n)_q$ is determined by $J : z_a^\alpha \mapsto (1 - q^2)^{\frac{1}{2}} z_a^\alpha$ for all a and α .

The algebra $\text{Pol}(\text{Mat}_{m,n})_q$ is a particular case of involutive algebras introduced in [9]. It follows from a general result of [9] that there exists a unique structure of $U_q \mathfrak{su}(n, m)$ -module algebra in $\text{Pol}(\text{Mat}_{m,n})_q$ so that (2.11), (2.12), (2.13), (4.1), (4.2), (4.3) hold. This means that we may use the relation (3.2) to 'transfer' the $U_q \mathfrak{sl}(n+m)$ -action from the subalgebra $\mathbb{C}[\text{Mat}_{m,n}]_q \subset \text{Pol}(\text{Mat}_{m,n})_q$ to the subalgebra $\mathbb{C}[\overline{\text{Mat}}_{m,n}]_q = \{f^* \mid f \in \mathbb{C}[\text{Mat}_{m,n}]_q\} \subset \text{Pol}(\text{Mat}_{m,n})_q$ of 'antiholomorphic' polynomials, and the resulting $U_q \mathfrak{sl}(n+m)$ -action respects the commutation relations (4.8).

The following example of a $U_q \mathfrak{su}(n, m)$ -module algebra appeared for the first time in [4]. Let us add to $\text{Pol}(\text{Mat}_{m,n})_q$ one more generator f_0 such that

$$f_0 = f_0^2 = f_0^*, \quad (z_a^\alpha)^* f_0 = f_0 z_a^\alpha = 0, \quad \forall a, \alpha. \quad (4.9)$$

The relations allow us to treat f_0 as a q -analogue of the function, which is equal to 1 in the center of the matrix ball $D_{m,n}$ and to 0 in other points. We denote the $*$ -algebra

by $\text{Fun}(\mathcal{D}_{m,n})_q$. There exists a unique extension of the structure of a $U_q\mathfrak{su}(n, m)$ -module algebra from $\text{Pol}(\text{Mat}_{m,n})_q$ to $\text{Fun}(\mathcal{D}_{m,n})_q$ such that

$$K_n f_0 = f_0, \quad F_n f_0 = -\frac{q^{1/2}}{q^{-2} - 1} f_0 \cdot (z_n^m)^*, \quad E_n f_0 = -\frac{q^{1/2}}{1 - q^2} z_n^m \cdot f_0, \quad (4.10)$$

$$(K_j^{\pm 1} - 1)f_0 = F_j f_0 = E_j f_0 = 0, \quad j \neq n. \quad (4.11)$$

Obviously, the two-sided ideal $\mathcal{D}(\mathcal{D}_{m,n})_q = \text{Fun}(\mathcal{D}_{m,n})_q \cdot f_0 \cdot \text{Fun}(\mathcal{D}_{m,n})_q$ is a $U_q\mathfrak{su}(n, m)$ -module algebra. We call its elements finite functions in the quantum matrix ball $\mathcal{D}_{m,n}$.

In this section we dealt with module algebras of functions on quantum G -spaces. The next section is devoted to a module algebra of a completely different nature.

5 Example of quantum symmetry: q -Weyl algebra

In this section we present a remarkable structure of $U_q\mathfrak{sl}(n + m)^{op}$ -module algebra in the q -Weyl algebra $A_{m,n}(\mathbb{C})_q$ (see section 2). The contents of this section is closely related to that of paper [7]. However, the q -analog of the Weyl algebra, treated here, differs from that considered in [7].

Remind that $A_{m,n}(\mathbb{C})_q$ is the unital subalgebra in $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ generated by the q -partial derivatives $\frac{\partial}{\partial z_a^\alpha}$ and the operators $\hat{z}_a^\alpha$ of right multiplication by z_a^α .

Consider the structure of a $U_q\mathfrak{sl}(n + m)$ -module in $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ given by (3.5). The following observation plays an important role in our proof of the main results.

Proposition 5.1 *$A_{m,n}(\mathbb{C})_q$ is a $U_q\mathfrak{sl}(n + m)$ -submodule in $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$. The induced $U_q\mathfrak{sl}(n + m)$ -action makes $A_{m,n}(\mathbb{C})_q$ into a $U_q\mathfrak{sl}(n + m)^{op}$ -module algebra.*

Proof. The second statement is a straightforward consequence of the first one since any invariant subalgebra in a module algebra is automatically a module algebra. Let us proof the first statement.

We have to explain that for arbitrary $\xi \in U_q\mathfrak{sl}(n + m)$ and a, α

$$\text{ad}'\xi(\hat{z}_a^\alpha) \in A_{m,n}(\mathbb{C})_q, \quad (5.1)$$

$$\text{ad}'\xi\left(\frac{\partial}{\partial z_a^\alpha}\right) \in A_{m,n}(\mathbb{C})_q. \quad (5.2)$$

Note that (5.1) is a simple consequence of $U_q\mathfrak{sl}(n + m)$ -moduleness of the algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$. Let us prove (5.2).

The crucial role in the proof plays $U_q\mathfrak{sl}(n + m)$ -covariance of the first order differential calculus $(\Lambda^1(\text{Mat}_{m,n})_q, d)$ (see section 2) observed for the first time in [8]. The covariance means that there exists a unique structure of $U_q\mathfrak{sl}(n + m)$ -module in $\Lambda^1(\text{Mat}_{m,n})_q$ such that the differential d is a morphism of the $U_q\mathfrak{sl}(n + m)$ -modules and $\Lambda^1(\text{Mat}_{m,n})_q$ is a $U_q\mathfrak{sl}(n + m)$ -module $\mathbb{C}[\text{Mat}_{m,n}]_q$ -bimodule.

That $d : \mathbb{C}[\text{Mat}_{m,n}]_q \rightarrow \Lambda^1(\text{Mat}_{m,n})_q$ intertwines the $U_q\mathfrak{sl}(n + m)$ -actions may be written as follows

$$\xi \left(\sum_{b,\beta} \frac{\partial f}{\partial z_b^\beta} dz_b^\beta \right) = \sum_{c,\gamma} \frac{\partial \xi(f)}{\partial z_c^\gamma} dz_c^\gamma, \quad f \in \mathbb{C}[\text{Mat}_{m,n}]_q, \quad \xi \in U_q\mathfrak{sl}(n + m).$$

Since $\Lambda^1(\text{Mat}_{m,n})_q$ is, in particular, a $U_q\mathfrak{sl}(n+m)$ -module *left* $\mathbb{C}[\text{Mat}_{m,n}]_q$ -module, we may rewrite the latter equality:

$$\sum_{b,\beta} \xi_{(1)} \left(\frac{\partial f}{\partial z_b^\beta} \right) \xi_{(2)}(dz_b^\beta) = \sum_{c,\gamma} \frac{\partial \xi(f)}{\partial z_c^\gamma} dz_c^\gamma \quad (5.3)$$

(here and further in the proof $\xi_{(1)} \otimes \xi_{(2)}$ stands for $\Delta(\xi)$; similarly, $\xi_{(1)} \otimes \xi_{(2)} \otimes \xi_{(3)}$ stands for $1 \otimes \Delta(\Delta(\xi)) = \Delta \otimes 1(\Delta(\xi))$ and so on).

It is noted in [4] that $\Lambda^1(\text{Mat}_{m,n})_q$ is a free left $\mathbb{C}[\text{Mat}_{m,n}]_q$ -module with the generators dz_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$. Thus, for any $\eta \in U_q\mathfrak{sl}(n+m)$ there exist elements $f_{\beta,a}^{b,\alpha}(\eta) \in \mathbb{C}[\text{Mat}_{m,n}]_q$, $a, b = 1, \dots, n$, $\alpha, \beta = 1, \dots, m$, such that

$$\eta(dz_b^\beta) = \sum_{a,\alpha} f_{\beta,a}^{b,\alpha}(\eta) dz_a^\alpha.$$

Thus, we may rewrite (5.3) as follows:

$$\sum_{b,\beta} \xi_{(1)} \left(\frac{\partial f}{\partial z_b^\beta} \right) f_{\beta,a}^{b,\alpha}(\xi_{(2)}) dz_a^\alpha = \sum_{c,\gamma} \frac{\partial \xi(f)}{\partial z_c^\gamma} dz_c^\gamma,$$

or, in terms of operators in $\mathbb{C}[\text{Mat}_{m,n}]_q$,

$$\sum_{b,\beta} \mathcal{R}_{f_{\beta,a}^{b,\alpha}(\xi_{(2)})} \cdot \xi_{(1)} \cdot \frac{\partial}{\partial z_b^\beta} = \frac{\partial}{\partial z_a^\alpha} \cdot \xi \quad (5.4)$$

where $\mathcal{R}_f$ stands for the operator in $\mathbb{C}[\text{Mat}_{m,n}]_q$ of right multiplication by $f \in \mathbb{C}[\text{Mat}_{m,n}]_q$.

Now we are ready to prove (5.2). By the definition

$$\text{ad}' \xi \left(\frac{\partial}{\partial z_a^\alpha} \right) = \xi_{(2)} \cdot \frac{\partial}{\partial z_a^\alpha} \cdot S^{-1}(\xi_{(1)}).$$

(5.4) implies

$$\xi_{(2)} \cdot \frac{\partial}{\partial z_a^\alpha} \cdot S^{-1}(\xi_{(1)}) = \sum_{b,\beta} \xi_{(3)} \cdot \mathcal{R}_{f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)}))} \cdot S^{-1}(\xi_{(2)}) \cdot \frac{\partial}{\partial z_b^\beta}.$$

Observe that for any $f \in \mathbb{C}[\text{Mat}_{m,n}]_q$ and any $\eta \in U_q\mathfrak{sl}(n+m)$

$$\eta \cdot \mathcal{R}_f = \mathcal{R}_{\eta_{(2)}(f)} \cdot \eta_{(1)}.$$

Thus

$$\sum_{b,\beta} \xi_{(3)} \cdot \mathcal{R}_{f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)}))} \cdot S^{-1}(\xi_{(2)}) \cdot \frac{\partial}{\partial z_b^\beta} = \sum_{b,\beta} \mathcal{R}_{\xi_{(4)}(f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)})))} \cdot \xi_{(3)} \cdot S^{-1}(\xi_{(2)}) \cdot \frac{\partial}{\partial z_b^\beta}.$$

What remains is to take into account the equality

$$\xi_{(1)} \otimes \xi_{(3)} S^{-1}(\xi_{(2)}) \otimes \xi_{(4)} = \xi_{(1)} \otimes \xi_{(2)}.$$

We finally get

$$\sum_{b,\beta} \mathcal{R}_{\xi_{(4)}(f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)})))} \cdot \xi_{(3)} \cdot S^{-1}(\xi_{(2)}) \cdot \frac{\partial}{\partial z_b^\beta} = \sum_{b,\beta} \mathcal{R}_{\xi_{(2)}(f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)})))} \cdot \frac{\partial}{\partial z_b^\beta}.$$

This finishes the proof of the proposition. ■

It is not very difficult to describe the $U_q \mathfrak{sl}(n+m)$ -action in $A_{m,n}(\mathbb{C})_q$ explicitly. But, for the purposes of the present paper, we need only the following partial result.

Impose the notation

$$\begin{aligned} L &= \text{linear span of } \{z_a^\alpha\}_{a,\alpha} \subset \mathbb{C}[\text{Mat}_{m,n}]_q, \\ L' &= \text{linear span of } \{\hat{z}_a^\alpha\}_{a,\alpha} \subset \text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q), \\ L'' &= \text{linear span of } \left\{ \frac{\partial}{\partial z_a^\alpha} \right\}_{a,\alpha} \subset \text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q). \end{aligned}$$

Note that L is a (finite dimensional) $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -submodule in $\mathbb{C}[\text{Mat}_{m,n}]_q$ due to (2.11), (2.12), (2.13), (4.1).

Proposition 5.2

i) The map $z_a^\alpha \mapsto \hat{z}_a^\alpha$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$ is extended to an isomorphism of the $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -modules $j' : L \rightarrow L'$;

ii) the map $z_a^\alpha \mapsto \frac{\partial}{\partial z_a^\alpha}$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$ is extended to an isomorphism of the vector spaces $j'' : L \rightarrow L''$, satisfying the following intertwining property:

$$j''(\xi(f)) = \text{ad}' \omega(\xi)(j''(f)), \quad \xi \in U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m), f \in \mathbb{C}[\text{Mat}_{m,n}]_q. \quad (5.5)$$

Here ω is the automorphism of $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ (the Chevalley involution) given by

$$\omega(E_i) = -F_i, \quad \omega(F_i) = -E_i, \quad \omega(K_i^{\pm 1}) = K_i^{\mp 1};$$

iii) $\frac{\partial}{\partial z_1^1}$ is a primitive vector in the $U_q \mathfrak{sl}(n+m)$ -module $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$:

$$\begin{aligned} \text{ad}' F_j \left(\frac{\partial}{\partial z_1^1} \right) &= 0 \quad \forall j, \\ \text{ad}' K_j \left(\frac{\partial}{\partial z_1^1} \right) &= \begin{cases} q^{-1} \frac{\partial}{\partial z_1^1} & , \quad j = 1 \text{ or } j = n+m-1 \\ \frac{\partial}{\partial z_1^1} & , \quad \text{otherwise} \end{cases}. \end{aligned}$$

Proof. Statement i) follows immediately from the $U_q \mathfrak{sl}(n+m)$ -moduleness of the algebra $\mathbb{C}[\text{Mat}_{m,n}]_q$.

Let us prove statement ii). First of all, let us prove that L'' is a (finite dimensional) $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -submodule in $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$.

We have derived the following formula for the action of $U_q \mathfrak{sl}(n+m)$ on q -partial derivatives (see the proof of the previous proposition):

$$\text{ad}' \xi \left(\frac{\partial}{\partial z_a^\alpha} \right) = \sum_{b,\beta} \mathcal{R}_{\xi_{(2)}(f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)})))} \cdot \frac{\partial}{\partial z_b^\beta} \quad (5.6)$$

where $\mathcal{R}_f$ stands for the operator in $\mathbb{C}[\text{Mat}_{m,n}]_q$ of right multiplication by f , and the functions $f_{\beta,a}^{b,\alpha}(\eta)$ are defined by means of the equality

$$\eta(dz_a^\alpha) = \sum_{b,\beta} f_{\beta,a}^{b,\alpha}(\eta) dz_b^\beta, \quad \eta \in U_q \mathfrak{sl}(n+m).$$

Observe that the elements $f_{\beta,a}^{b,\alpha}(\eta) \in \mathbb{C}[\text{Mat}_{m,n}]_q$ are constants for $\eta \in U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$. This is a consequence of (2.11), (2.12), (2.13), and (4.1). Thus, if $\xi \in U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$, the elements $\mathcal{R}_{\xi_{(2)}(f_{\beta,a}^{b,\alpha}(S^{-1}(\xi_{(1)})))}$ are constants as well. The formula (5.6) then implies that L'' is a $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -invariant subspace in $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$.

Remind (see section 3) that the map

$$\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q) \otimes \mathbb{C}[\text{Mat}_{m,n}]_q \rightarrow \mathbb{C}[\text{Mat}_{m,n}]_q, \quad T \otimes f \mapsto T(f) \quad (5.7)$$

makes $\mathbb{C}[\text{Mat}_{m,n}]_q$ into a $U_q \mathfrak{sl}(n+m)^{op}$ -module left $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ -module. Consequently, the restriction of the map onto the subspace $L'' \otimes L$ defines the map $\tau : L'' \otimes L \rightarrow \mathbb{C}[\text{Mat}_{m,n}]_q$, satisfying

$$\xi(\tau(T \otimes f)) = \tau(\text{ad}' \xi_{(2)}(T) \otimes \xi_{(1)}(f)), \quad \xi \in U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m).$$

Clearly, the image of τ is the subspace of constants $\mathbb{C} \subset \mathbb{C}[\text{Mat}_{m,n}]_q$. Thus, τ defines a $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -invariant pairing $\tau^{op} = \tau \circ \sigma : L \otimes L'' \rightarrow \mathbb{C}$ with σ being the flip of tensor multipliers. Note that the basis $\{z_a^\alpha\}_{a,\alpha}$ is dual to the basis $\left\{\frac{\partial}{\partial z_a^\alpha}\right\}_{a,\alpha}$ with respect to the pairing. This observation, together with the invariance of the pairing, implies, in particular, that the matrix of the operator $\text{ad}' E_j$ in the basis $\left\{\frac{\partial}{\partial z_a^\alpha}\right\}_{a,\alpha}$ coincides with the transposed to the matrix of the operator $-E_j K_j^{-1}$ in the basis $\{z_a^\alpha\}_{a,\alpha}$. But the latter matrix is easily seen to be the transposed to the matrix of the operator $-F_i$ in the basis $\{z_a^\alpha\}_{a,\alpha}$. This proves (5.5) in the case $\xi = E_j$, $j \neq n$. The cases $\xi = F_j$, $\xi = K_j^{\pm 1}$ may be proved in the same way.

Turn to statement iii) of the proposition. Since K_i for any i and F_j for $j \neq n$ belong to $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$, a part of the statement follows from statement ii) and formulae (2.11), (2.13), and (4.1). What we have to prove is the equality

$$\text{ad}' F_n \left(\frac{\partial}{\partial z_1^1} \right) = 0.$$

Use the formula (5.6):

$$\text{ad}' F_n \left(\frac{\partial}{\partial z_1^1} \right) = \sum_{b,\beta} \mathcal{R}_{F_n(f_{1,a}^{1,\alpha}(1))} \cdot \frac{\partial}{\partial z_b^\beta} - \sum_{b,\beta} \mathcal{R}_{K_n^{-1}(f_{1,a}^{1,\alpha}(K_n F_n))} \cdot \frac{\partial}{\partial z_b^\beta}.$$

Clearly, $f_{1,a}^{1,\alpha}(1) = \delta_{a,1} \delta_{\alpha,1}$, and $F_n(f_{1,a}^{1,\alpha}(1)) = 0$ for any a and α . By using (4.2), we also get $\mathcal{R}_{K_n^{-1}(f_{1,a}^{1,\alpha}(K_n F_n))} = 0$. That is

$$\text{ad}' F_n \left(\frac{\partial}{\partial z_1^1} \right) = 0.$$

The proposition is proved. ■

6 Proof of the main results

In this section we prove the theorems formulated in section 2.

Remind the notation $(\cdot, \cdot)$ for the inner product in $\mathbb{C}[\text{Mat}_{m,n}]_q$ introduced in section 2 (see theorem 2.3). First of all, we are going to express this inner product via another one, which is constructed by means of the algebra $\mathcal{D}(\text{D}_{m,n})_q$ of finite functions in the quantum matrix ball (see section 4).

Let P, Q be elements of $\mathbb{C}[\text{Mat}_{m,n}]_q$. Consider the finite function $f_0 \cdot Q^* \cdot P \cdot f_0 \in \mathcal{D}(\text{D}_{m,n})_q$ (here f_0 is the 'distinguished' finite function which, along with $\text{Pol}(\text{Mat}_{m,n})_q$, generates the algebra $\text{Fun}(\text{D}_{m,n})_q$; see section 4). The properties (4.9) imply that the finite function $f_0 \cdot Q^* \cdot P \cdot f_0$ differs from f_0 by a constant. Obviously, the constant depends linearly on P and conjugate linearly on Q . Thus, we get the sesquilinear form $\langle \cdot, \cdot \rangle$ on $\mathbb{C}[\text{Mat}_{m,n}]_q$:

$$f_0 \cdot Q^* \cdot P \cdot f_0 = \langle P, Q \rangle \cdot f_0. \quad (6.1)$$

It turns out to be closely related to the inner product $(\cdot, \cdot)$. Namely, recall one the notation J for the algebra isomorphism $\text{Pol}(\text{Mat}_{m,n})_q \rightarrow \mathcal{P}(m, n)_q$ (see section 4). It is explicitly given by $J : z_a^\alpha \mapsto (1 - q^2)^{\frac{1}{2}} \cdot z_a^\alpha$ for all a and α . It follows almost immediately from the definition of the inner product $(\cdot, \cdot)$ that

$$\langle P, Q \rangle = (J(P), J(Q)), \quad P, Q \in \mathbb{C}[\text{Mat}_{m,n}]_q. \quad (6.2)$$

Indeed, the subspace $\mathbb{C}[\text{Mat}_{m,n}]_q f_0 \subset \mathcal{D}(\text{D}_{m,n})_q$ is invariant under left multiplication by elements of $\text{Pol}(\text{Mat}_{m,n})_q$ (see (4.9)). This allows us to define a structure of $\mathcal{P}(m, n)_q$ -module in the subspace $\mathbb{C}[\text{Mat}_{m,n}]_q f_0 \subset \mathcal{D}(\text{D}_{m,n})_q$ by means of the isomorphism J :

$$F \otimes (P \cdot f_0) \mapsto J^{-1}(F) \cdot P \cdot f_0, \quad F \in \mathcal{P}(m, n)_q, P \in \mathbb{C}[\text{Mat}_{m,n}]_q.$$

Obviously, the $\mathcal{P}(m, n)_q$ -module $\mathbb{C}[\text{Mat}_{m,n}]_q f_0$ is isomorphic to H (see section 2), and the inner product $\langle \cdot, \cdot \rangle$, regarded as an inner product on $\mathbb{C}[\text{Mat}_{m,n}]_q f_0$, satisfies the properties (2.20). Since such an inner product is unique, we have (6.2).

In what follows, we denote by J the automorphism of $\mathbb{C}[\text{Mat}_{m,n}]_q$ given by $J : z_a^\alpha \mapsto (1 - q^2)^{\frac{1}{2}} \cdot z_a^\alpha$ for all a and α . Suppose we have proved theorem 2.3. Then theorem 2.2 is a straightforward consequence of (6.2). Indeed, J commutes with the $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -action since J is just a constant operator on any homogeneous component of $\mathbb{C}[\text{Mat}_{m,n}]_q$, and the homogeneous components are $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -submodules in $\mathbb{C}[\text{Mat}_{m,n}]_q$. So we have to prove $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -invariance of the inner product $\langle \cdot, \cdot \rangle$. This may be derived from the $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -moduleness of the algebra $\text{Fun}(\text{D}_{m,n})_q$ (see section 4) and the relations (4.11) as follows. The relations (4.11) mean $U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -invariance of the element f_0 :

$$\xi(f_0) = \varepsilon(\xi) \cdot f_0, \quad \xi \in U_q \mathfrak{sl}(n) \oplus \mathfrak{gl}(m).$$

This and the equality (3.2) imply

$$\xi(f_0 \cdot Q^* \cdot P \cdot f_0) = f_0 \cdot \xi_{(1)}(Q^*) \cdot \xi_{(2)}(P) \cdot f_0 = f_0 \cdot (S(\xi_{(1)})^*(Q))^* \cdot \xi_{(2)}(P) \cdot f_0, \quad P, Q \in \mathbb{C}[\text{Mat}_{m,n}]_q$$

(here $\xi_{(1)} \otimes \xi_{(2)}$ stands for $\Delta(\xi)$). On the other hand, by (6.1) $f_0 \cdot Q^* \cdot P \cdot f_0 = \langle P, Q \rangle \cdot f_0$, i.e.

$$f_0 \cdot (S(\xi_{(1)})^*(Q))^* \cdot \xi_{(2)}(P) \cdot f_0 = \xi(\langle P, Q \rangle \cdot f_0) = \varepsilon(\xi) \cdot \langle P, Q \rangle \cdot f_0.$$

Take into account (6.1) once again:

$$\langle \xi_{(2)}(P), S(\xi_{(1)})^*(Q) \rangle = \varepsilon(\xi) \cdot \langle P, Q \rangle, \quad \xi \in U_q \mathfrak{s}(\mathfrak{u}(n) \oplus \mathfrak{u}(m)).$$

The latter property is obviously equivalent to $U_q \mathfrak{s}(\mathfrak{u}(n) \oplus \mathfrak{u}(m))$ -invariance of the inner product $\langle \cdot, \cdot \rangle$.

In view of the above arguments, the first two theorems, stated in section 2, follow from the third one. The remaining part of this section is devoted to a proof of theorem 2.3. It will be convenient for us to prove the following statement instead of theorem 2.3 itself. It is equivalent to the statement of the theorem due to (6.2).

Theorem 6.1

$$\left\langle \frac{\partial P}{\partial z_a^\alpha}, Q \right\rangle = \frac{1}{1 - q^2} \cdot \langle P, Q \cdot z_a^\alpha \rangle \quad \forall a, \alpha. \quad (6.3)$$

Proof. Let us agree about the following notation: if $T \in \text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ then $T^\dagger \in \text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ stands for the conjugate operator to T with respect to the inner product $\langle \cdot, \cdot \rangle$. In this notation (6.3) says

$$(\hat{z}_a^\alpha)^\dagger = (1 - q^2) \cdot \frac{\partial}{\partial z_a^\alpha} \quad \forall a, \alpha. \quad (6.4)$$

Let us explain how $U_q \mathfrak{s}(\mathfrak{u}(n) \oplus \mathfrak{u}(m))$ -invariance of the inner product $\langle \cdot, \cdot \rangle$ allows one to reduce the general case to the case $a = 1, \alpha = 1$. Observe that the invariance of the inner product $\langle \cdot, \cdot \rangle$ means

$$\xi^\dagger = \xi^*, \quad \xi \in U_q \mathfrak{s}(\mathfrak{u}(n) \oplus \mathfrak{u}(m)). \quad (6.5)$$

Suppose we have already proved (6.4) in the case $a = 1, \alpha = 1$. Proposition 5.2 (i) and the formulae (2.12) imply

$$\hat{z}_2^1 = q^{-1/2} \cdot \text{ad}' F_1(\hat{z}_1^1).$$

Thus

$$\begin{aligned} (\hat{z}_2^1)^\dagger &= q^{-1/2} \cdot (\text{ad}' F_1(\hat{z}_1^1))^\dagger = q^{-1/2} \cdot (F_1 \cdot \hat{z}_1^1 - K_1^{-1} \cdot \hat{z}_1^1 \cdot K_1 \cdot F_1)^\dagger = \\ &= q^{-1/2} ((\hat{z}_1^1)^\dagger \cdot F_1^\dagger - F_1^\dagger \cdot K_1^\dagger \cdot (\hat{z}_1^1)^\dagger \cdot (K_1^{-1})^\dagger) = q^{-1/2} ((\hat{z}_1^1)^\dagger \cdot E_1 K_1^{-1} - E_1 K_1^{-1} \cdot K_1 \cdot (\hat{z}_1^1)^\dagger \cdot K_1^{-1}) = \\ &= -q^{-1/2} (1 - q^2) (E_1 \cdot \frac{\partial}{\partial z_1^1} \cdot K_1^{-1} - \frac{\partial}{\partial z_1^1} \cdot E_1 K_1^{-1}) = -q^{-1/2} (1 - q^2) \text{ad}' E_1 \left(\frac{\partial}{\partial z_1^1} \right) = (1 - q^2) \frac{\partial}{\partial z_2^1} \end{aligned}$$

(the latter equality follows from proposition 5.2 (ii)). The other cases may be proved in a completely analogous way. Thus, it remains to prove (6.4) in the case $a = 1, \alpha = 1$.

Lemma 6.2 Suppose $T \in \text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ satisfies the properties

$$i) \quad \text{ad}' K_j(T) = \begin{cases} q^{-1} T & , \quad j = 1 \text{ or } j = n + m - 1 \\ T & , \quad \text{otherwise} \end{cases}; \quad (6.6)$$

$$ii) \quad \text{ad}' F_j(T) = 0 \quad \forall j; \quad (6.7)$$

$$iii) \quad T(z_a^\alpha) = 0 \quad \forall a, \alpha.$$

Then $T \equiv 0$.

Proof of the lemma. Let us denote by $\mathbb{C}[\text{Mat}_{m,n}]_{q,k}$ the k -th homogeneous component in $\mathbb{C}[\text{Mat}_{m,n}]_q$ (remind that the latter algebra admits the natural $\mathbb{Z}_+$ -grading by powers of monomials). As we have already noted, each subspace $\mathbb{C}[\text{Mat}_{m,n}]_{q,k}$ is a (finite dimensional) $U_q\mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -submodule in $\mathbb{C}[\text{Mat}_{m,n}]_q$ ($U_q\mathfrak{sl}(n) \oplus \mathfrak{gl}(m)$ -invariance of the subspaces is a straightforward consequence of the formulae (2.11), (2.12), (2.13), (4.1)). The subspaces $\mathbb{C}[\text{Mat}_{m,n}]_{q,k}$ admit the following 'coordinateless' description (see (2.15))

$$\mathbb{C}[\text{Mat}_{m,n}]_{q,k} = \{f \in \mathbb{C}[\text{Mat}_{m,n}]_q \mid K_0(f) = q^{(n+m)k} f\} \quad (6.8)$$

(remind that K_0 corresponds to the element $K_n^{nm} \cdot \prod_{i=1}^{n-1} K_i^{mi} \cdot \prod_{j=1}^{m-1} K_{n+j}^{n(m-j)}$ under the embedding $U_q\mathfrak{sl}(n) \oplus \mathfrak{gl}(m) \subset U_q\mathfrak{sl}(n+m)$; see section 4). This description, together with (6.6), implies

$$T(\mathbb{C}[\text{Mat}_{m,n}]_{q,k}) \subset \mathbb{C}[\text{Mat}_{m,n}]_{q,k-1}.$$

We are going to prove the equality $T|_{\mathbb{C}[\text{Mat}_{m,n}]_{q,k}} = 0$ by induction in k . The property iii) of T is the induction base.

Suppose we have already proved that $T|_{\mathbb{C}[\text{Mat}_{m,n}]_{q,k}} = 0$ for any $k \leq M$ ($M \geq 1$), and let $T|_{\mathbb{C}[\text{Mat}_{m,n}]_{q,M+1}} \neq 0$. Let $f \in \mathbb{C}[\text{Mat}_{m,n}]_{q,M+1}$ be such an element that $T(f) \neq 0$. We may assume that f is a weight vector of the $U_q\mathfrak{sl}(n+m)$ -module $\mathbb{C}[\text{Mat}_{m,n}]_q$ (i.e. an eigenvector of each K_j , $j = 1, \dots, n+m-1$). In this case the element $T(f)$ is a weight vector as well (this is a consequence of (6.6) and $U_q\mathfrak{sl}(n+m)^{op}$ -moduleness of the left $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ -module $\mathbb{C}[\text{Mat}_{m,n}]_q$). We intend to construct an element $\tilde{f} \in \mathbb{C}[\text{Mat}_{m,n}]_{q,M+1}$ so that

- a) $T(\tilde{f}) \neq 0$;
- b) $F_j(T(\tilde{f})) = 0 \quad \forall j$;
- c) $T(\tilde{f})$ is a weight vector.

To start with, let us note that (6.7) means

$$F_j T = \begin{cases} q T F_j & , \quad j = 1 \text{ or } j = n+m-1 \\ T F_j & , \quad \text{otherwise} \end{cases} \quad (6.9)$$

Let some $j_1 \neq n$ satisfies $F_{j_1}(T(f)) \neq 0$. If there is no such j_1 then $\tilde{f} = f$. Indeed, f satisfies the properties a) and c), and the property b) for $j \neq n$. $F_n(T(f)) = T(F_n(f))$ due to (6.9). But the formula (6.8) implies $F_n(f) \in \mathbb{C}[\text{Mat}_{m,n}]_{q,M}$, and thus $T(F_n(f)) = 0$ (the induction assumption). So $\tilde{f}$ has been built.

If j_1 with the above property exists, we set $f_1 = F_{j_1}(f)$. Let $j_2 \neq n$ satisfies $F_{j_2}(T(f_1)) \neq 0$. If there is no such j_2 then we set $\tilde{f} = f_1$ and so on. Since $\mathbb{C}[\text{Mat}_{m,n}]_{q,M}$ is finite dimensional, this process will give us an element $\tilde{f}$ satisfying the properties a), b), and c). Thus we get a non-zero primitive weight vector (namely, $T(\tilde{f})$) in the $U_q\mathfrak{sl}(n+m)$ -module $\mathbb{C}[\text{Mat}_{m,n}]_q$, which belongs to $\mathbb{C}[\text{Mat}_{m,n}]_{q,M}$ with $M \geq 1$. Let us show that this is a contradiction.

Remind that the $U_q\mathfrak{sl}(n+m)$ -module $\mathbb{C}[\text{Mat}_{m,n}]_q$ is the graded dual to a generalized Verma module $V(0)$ (see section 4). Let us denote by $(\cdot, \cdot)_0$ the pairing $\mathbb{C}[\text{Mat}_{m,n}]_q \times V(0) \rightarrow \mathbb{C}$. The term 'dual $U_q\mathfrak{sl}(n+m)$ -module' means

$$(\xi(f), v)_0 = (f, S(\xi)(v))_0, \quad \forall \xi \in U_q\mathfrak{sl}(n+m), f \in \mathbb{C}[\text{Mat}_{m,n}]_q, v \in V(0). \quad (6.10)$$

Any element from $V(0)$ has the form $\xi(v(0))$, where $v(0)$ is the generator of $V(0)$ (see section 4) and ξ is an element from the unital subalgebra $U_q \mathfrak{n}_- \subset U_q \mathfrak{sl}(n+m)$ generated by all F_j 's. This observation is a consequence of the equalities (4.4) and a PBW-type theorem for the quantized universal enveloping algebra $U_q \mathfrak{sl}(n+m)$ [2].

Let us return now to the non-zero element $T(\tilde{f})$. It satisfies (6.7). One then has

$$(T(\tilde{f}), \xi(v(0)))_0 = (S^{-1}(\xi)(T(\tilde{f})), v(0))_0 = \varepsilon(\xi) \cdot (T(\tilde{f}), v(0))_0$$

for any $\xi \in U_q \mathfrak{n}_-$. This means that the functional $T(\tilde{f}) \in V(0)^*$ is equal to 0 on any graded component $V(0)_{-k}$ (see (4.6)). That is $T(\tilde{f}) \in \mathbb{C}[\text{Mat}_{m,n}]_{q,0} = \mathbb{C}$. Since $T(\tilde{f}) \in \mathbb{C}[\text{Mat}_{m,n}]_{q,M}$ ($M \geq 1$), we get $T(\tilde{f}) = 0$.

We see that the assumption $T|_{\mathbb{C}[\text{Mat}_{m,n}]_{q,M+1}} \neq 0$ leads to a contradiction. The lemma is proved. $\blacksquare$

Corollary 6.3 *The subspace in $\text{End}_{\mathbb{C}}(\mathbb{C}[\text{Mat}_{m,n}]_q)$ of operators, satisfying the properties i) and ii) of the previous lemma, is one dimensional.*

Proof. Let T' and T'' be two non-zero linear operators, satisfying the properties i) and ii). One has

$$T'(z_a^\alpha) = T''(z_a^\alpha) = 0, \quad a \neq 1 \quad \text{or} \quad \alpha \neq 1. \quad (6.11)$$

Indeed, by the formulae (2.12)

$$z_a^\alpha = c_a^\alpha \cdot F_{a-1} F_{a-2} \dots F_1 F_{n+m+1-\alpha} \dots F_{n+m-2} F_{n+m-1}(z_1^1)$$

for a non-zero constant c_a^α . Then

$$\begin{aligned} T'(z_a^\alpha) &= c_a^\alpha \cdot T'(F_{a-1} F_{a-2} \dots F_1 F_{n+m+1-\alpha} \dots F_{n+m-2} F_{n+m-1}(z_1^1)) = \\ &= c_a^\alpha \cdot F_{a-1} F_{a-2} \dots F_1 F_{n+m+1-\alpha} \dots F_{n+m-2} F_{n+m-1}(T'(z_1^1)) \end{aligned}$$

(the same is true for T''). What remains is to use the fact that $T'(z_1^1), T''(z_1^1) \in \mathbb{C}$ (we pointed out in the proof of the above lemma that any linear operator T in $\mathbb{C}[\text{Mat}_{m,n}]_q$, satisfying i) and ii), possesses the property $T(\mathbb{C}[\text{Mat}_{m,n}]_{q,k}) \subset \mathbb{C}[\text{Mat}_{m,n}]_{q,k-1}$.)

Suppose

$$T'(z_1^1) = c_1, \quad T''(z_1^1) = c_2$$

for certain constants c_1 and c_2 . If, for example, $c_1 = 0$ then $T' = 0$ due to the above lemma. Thus both constants are non-zero. Then, by using the lemma once again, we get

$$c_2 \cdot T' - c_1 \cdot T'' = 0.$$

$\blacksquare$

To complete the proof of theorem 6.1, it suffices to establish that

1. $(\hat{z}_1^1)^\dagger$ satisfies the conditions i), ii) of lemma 6.2;
2. $\frac{\partial}{\partial z_1^1}$ satisfies the conditions i), ii) of lemma 6.2;
3. $(\hat{z}_1^1)^\dagger(z_1^1) = (1 - q^2) \frac{\partial}{\partial z_1^1}(z_1^1)$.

Note that point 2 is just the statement iii) of proposition 5.2. Point 3 is a simple consequence of definitions: clearly, $(1 - q^2) \cdot \frac{\partial}{\partial z_1^1}(z_1^1) = 1 - q^2$; on the other hand, $(\hat{z}_1^1)^\dagger(z_1^1)$ is easily seen to be a constant, so

$$(\hat{z}_1^1)^\dagger(z_1^1) = \langle (\hat{z}_1^1)^\dagger(z_1^1), 1 \rangle = \langle z_1^1, z_1^1 \rangle = 1 - q^2.$$

What remains is to establish point 1. Observe, that the equalities

$$\begin{aligned} \text{ad}' K_j((\hat{z}_1^1)^\dagger) &= \begin{cases} q^{-1}(\hat{z}_1^1)^\dagger & , \quad j = 1 \text{ or } j = n + m - 1 \\ (\hat{z}_1^1)^\dagger & , \quad \text{otherwise} \end{cases} , \\ \text{ad}' F_j((\hat{z}_1^1)^\dagger) &= 0 \quad j \neq n \end{aligned} \quad (6.12)$$

may be derived from (6.5) and the statement i) of proposition 5.2. For example, we prove the equality $\text{ad}' F_1((\hat{z}_1^1)^\dagger) = 0$:

$$\begin{aligned} \text{ad}' F_1((\hat{z}_1^1)^\dagger) &= F_1 \cdot (\hat{z}_1^1)^\dagger - K_1^{-1} \cdot (\hat{z}_1^1)^\dagger \cdot K_1 \cdot F_1 = \\ &= (\hat{z}_1^1 \cdot F_1^\dagger - F_1^\dagger \cdot K_1^\dagger \cdot \hat{z}_1^1 \cdot (K_1^{-1})^\dagger)^\dagger = (\hat{z}_1^1 \cdot E_1 K_1^{-1} - E_1 K_1^{-1} \cdot K_1 \cdot \hat{z}_1^1 \cdot K_1^{-1})^\dagger = \\ &= -(E_1 \cdot \hat{z}_1^1 \cdot K_1^{-1} - \hat{z}_1^1 \cdot E_1 K_1^{-1}) = -(\text{ad}' E_1(\hat{z}_1^1))^\dagger = 0. \end{aligned}$$

The other cases in (6.12) are proved just as this one.

Finally, we have to show that $\text{ad}' F_n((\hat{z}_1^1)^\dagger) = 0$ or, equivalently, $F_n \cdot (\hat{z}_1^1)^\dagger = (\hat{z}_1^1)^\dagger \cdot F_n$ (see (6.9)). The latter equality, in turn, is equivalent to

$$F_n^\dagger \cdot \hat{z}_1^1 = \hat{z}_1^1 \cdot F_n^\dagger. \quad (6.13)$$

Lemma 6.4 $F_n^\dagger = -E_n K_n^{-1} + \frac{q^{1/2}}{1-q^2} \hat{z}_n^m.$

Proof of the lemma to be found in Appendix. ■

Now we are ready to prove (6.13). Let $P \in \mathbb{C}[\text{Mat}_{m,n}]_q$. Then

$$F_n^\dagger \hat{z}_1^1(P) - \hat{z}_1^1 F_n^\dagger(P) = F_n^\dagger(P \cdot z_1^1) - F_n^\dagger(P) \cdot z_1^1.$$

By the previous lemma

$$\begin{aligned} &F_n^\dagger(P \cdot z_1^1) - F_n^\dagger(P) \cdot z_1^1 = \\ &= -E_n K_n^{-1}(P \cdot z_1^1) + \frac{q^{1/2}}{1-q^2} P \cdot z_1^1 \cdot z_n^m + E_n K_n^{-1}(P) \cdot z_1^1 - \frac{q^{1/2}}{1-q^2} P \cdot z_n^m \cdot z_1^1 = \\ &= -E_n(K_n^{-1}(P) \cdot K_n^{-1}(z_1^1)) + E_n K_n^{-1}(P) \cdot z_1^1 + \frac{q^{1/2}}{1-q^2} P \cdot (z_1^1 \cdot z_n^m - z_n^m \cdot z_1^1) \stackrel{\text{see(4.1)}}{=} \\ &= -E_n K_n^{-1}(f) \cdot z_1^1 - P \cdot E_n(z_1^1) + E_n K_n^{-1}(P) \cdot z_1^1 + \frac{q^{1/2}}{1-q^2} P \cdot (z_1^1 \cdot z_n^m - z_n^m \cdot z_1^1) = \\ &= -P \cdot E_n(z_1^1) + \frac{q^{1/2}}{1-q^2} P \cdot (z_1^1 \cdot z_n^m - z_n^m \cdot z_1^1) \stackrel{\text{see(4.3)}}{=} \\ &= q^{-1/2} P \cdot z_1^m \cdot z_n^1 + \frac{q^{1/2}}{1-q^2} P \cdot (z_1^1 \cdot z_n^m - z_n^m \cdot z_1^1) \stackrel{\text{see(1.5)}}{=} 0. \end{aligned}$$

Theorem 6.1 is proved completely. ■

7 Appendix. Proof of lemma 6.4

Let $P, Q \in \mathbb{C}[\text{Mat}_{m,n}]_q$. By the definition

$$\langle F_n(P), Q \rangle = f_0 \cdot Q^* \cdot F_n(P) \cdot f_0.$$

The equalities (4.10) and $U_q \mathfrak{su}(n, m)$ -moduleness of the algebra $\text{Fun}(\text{D}_{m,n})_q$ (see section 4) imply

$$F_n(P) \cdot f_0 = F_n(P) \cdot K_n^{-1}(f_0) = F_n(P \cdot f_0) - P \cdot F_n(f_0) = F_n(P \cdot f_0) + \frac{q^{1/2}}{q^{-2} - 1} \cdot P \cdot f_0 \cdot (z_n^m)^*.$$

Thus

$$\begin{aligned} \langle F_n(P), Q \rangle &= f_0 \cdot Q^* \cdot F_n(P \cdot f_0) + \frac{q^{1/2}}{q^{-2} - 1} \cdot f_0 \cdot Q^* \cdot P \cdot f_0 \cdot (z_n^m)^* = \\ &= F_n(f_0 \cdot Q^* \cdot P \cdot f_0) - F_n(f_0 \cdot Q^*) \cdot K_n^{-1}(P \cdot f_0) + \frac{q^{1/2}}{q^{-2} - 1} \cdot f_0 \cdot Q^* \cdot P \cdot f_0 \cdot (z_n^m)^* = \\ &= F_n(\langle P, Q \rangle \cdot f_0) - F_n(f_0 \cdot Q^*) \cdot K_n^{-1}(P) \cdot f_0 + \frac{q^{1/2}}{q^{-2} - 1} \cdot \langle P, Q \rangle \cdot f_0 \cdot (z_n^m)^* \stackrel{\text{see(4.10)}}{=} \\ &= \frac{-q^{1/2}}{q^{-2} - 1} \cdot \langle P, Q \rangle \cdot f_0 \cdot (z_n^m)^* - F_n(f_0 \cdot Q^*) \cdot K_n^{-1}(P) \cdot f_0 + \frac{q^{1/2}}{q^{-2} - 1} \cdot \langle P, Q \rangle \cdot f_0 \cdot (z_n^m)^* = \\ &= -F_n(f_0 \cdot Q^*) \cdot K_n^{-1}(P) \cdot f_0. \end{aligned}$$

Evidently,

$$\begin{aligned} F_n(f_0 \cdot Q^*) &= F_n(f_0) \cdot K_n^{-1}(Q^*) + f_0 \cdot F_n(Q^*) \stackrel{\text{see(4.10)}}{=} \\ &= -\frac{q^{1/2}}{q^{-2} - 1} \cdot f_0 \cdot (z_n^m)^* \cdot K_n^{-1}(Q^*) + f_0 \cdot F_n(Q^*) \stackrel{\text{see(3.2)}}{=} \\ &= -\frac{q^{1/2}}{q^{-2} - 1} \cdot f_0 \cdot (z_n^m)^* \cdot (K_n(Q))^* + q^2 \cdot f_0 \cdot (E_n(Q))^*. \end{aligned}$$

Finally we get

$$\begin{aligned} \langle F_n(P), Q \rangle &= \frac{q^{1/2}}{q^{-2} - 1} \cdot f_0 \cdot (z_n^m)^* \cdot (K_n(Q))^* \cdot K_n^{-1}(P) \cdot f_0 - q^2 \cdot f_0 \cdot (E_n(Q))^* \cdot K_n^{-1}(P) \cdot f_0 = \\ &= \frac{q^{1/2}}{q^{-2} - 1} \cdot f_0 \cdot (K_n(Q) \cdot z_n^m)^* \cdot K_n^{-1}(P) \cdot f_0 - q^2 \cdot f_0 \cdot (E_n(Q))^* \cdot K_n^{-1}(P) \cdot f_0 = \\ &= \frac{q^{1/2}}{q^{-2} - 1} \cdot \langle K_n^{-1}(P), K_n(Q) \cdot z_n^m \rangle - q^2 \cdot \langle K_n^{-1}(P), E_n(Q) \rangle = \\ &= \frac{q^{1/2}}{q^{-2} - 1} \cdot \langle P, K_n^{-1}(K_n(Q) \cdot z_n^m) \rangle - q^2 \cdot \langle P, K_n^{-1}E_n(Q) \rangle \stackrel{\text{see(4.1)}}{=} \\ &= \frac{q^{-3/2}}{q^{-2} - 1} \cdot \langle P, Q \cdot z_n^m \rangle - \langle P, E_n K_n^{-1}(Q) \rangle = \\ &= \langle P, -E_n K_n^{-1}(Q) + \frac{q^{-3/2}}{q^{-2} - 1} \cdot Q \cdot z_n^m \rangle. \end{aligned}$$

References

- [1] J. Arazy, *A Survey of Invariant Hilbert Spaces of Analytic Functions on Bounded Symmetric Domains*, Contemp. Math., **185** (1995), 7 – 64.
- [2] J. C. Jantzen. Lectures in Quantum Groups. Graduate Studies in Mathematics, AMS.
- [3] W. Pusz, S. Woronowicz, *Twisted second quantization*, Reports Math. Phys., **27** (1989), 231 – 257.
- [4] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035, 1999.
- [5] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: The Bergman kernel*, math.QA/9909036.
- [6] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Fock representations and quantum matrices*, in preparation.
- [7] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Hidden Symmetry of Some Algebras of q -Bidifferential Operators*, in Noncommutative Structures in Mathematics and Physics, S. Duplij and J. Wess (eds.), Kluwer Academic Publishers, 2001, 309 – 320 (see also this volume).
- [8] S. Sinel'shchikov and L. Vaksman, *Hidden symmetry of the differential calculus on the quantum matrix space*, J. Phys. A. **30** (1997), 23 – 26 (see also this volume).
- [9] S. Sinel'shchikov and L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, Math. Phys., Anal., and Geom., **1** (1998), 75 – 100; q-alg/9703005, 1997.

GEOMETRIC REALIZATIONS FOR SOME SERIES OF REPRESENTATIONS OF THE QUANTUM GROUP $SU_{2,2}$

D. Shklyarov S. Sinel'shchikov L. Vaksman

Institute for Low Temperature Physics & Engineering
47 Lenin Avenue, 61103 Kharkov, Ukraine

1 Introduction

We consider some series of modules over the quantum universal enveloping Drinfeld-Jimbo algebra $U_q\mathfrak{g}$ in the special case $\dim \mathfrak{g} < \infty$, $0 < q < 1$. The finite dimensional $U_q\mathfrak{g}$ -modules are closely related to compact quantum groups; those were investigated well enough [4, 13]. Infinite dimensional $U_q\mathfrak{g}$ -modules we deal with in this work originate from our earlier paper [19], together with some applications therein to the theory of q-Cartan domains. To make the exposition more transparent, we restrict ourselves to a q-analogue of the ball in the space of all complex 2×2 matrices $\mathbb{U} = \{z \in \text{Mat}_2 \mid zz^* < 1\}$, which is among the simplest Cartan domains.

The classes of infinite dimensional $U_q\mathfrak{g}$ -modules in question differ from those considered by Letzter [11]. The problem of producing and investigating of the principal series of quantum Harish-Chandra modules in our case appears to be essentially more complicated.

It is worthwhile to note that some properties of the ladder representation of the quantum $SU_{2,2}$ described below are already well known [2].

Everywhere in the sequel $0 < q < 1$, the ground field is $\mathbb{C}$, and all the algebras are assumed to be unital, unless the contrary is stated explicitly.

Consider the Hopf algebra $U_q\mathfrak{g} = U_q\mathfrak{sl}_4$ determined by the standard lists of generators $E_j, F_j, K_j^{\pm 1}$, $j = 1, 2, 3$, and relations [4, 13]. The coproduct Δ , the counit ε , and the antipode S are given as follows:

$$\begin{aligned} \Delta E_j &= E_j \otimes 1 + K_j \otimes E_j, & \varepsilon(E_j) &= 0, & S(E_j) &= -K_j^{-1}E_j, \\ \Delta F_j &= F_j \otimes K_j^{-1} + 1 \otimes F_j, & \varepsilon(F_j) &= 0, & S(F_j) &= -F_jK_j, \\ \Delta K_j &= K_j \otimes K_j, & \varepsilon(K_j) &= 1, & S(K_j) &= K_j^{-1}. \end{aligned}$$

We call a $U_q\mathfrak{g}$ -module V $\mathbb{R}^3$ -weight module if $V = \bigoplus_{\mu} V_{\mu}$ with $\mu = (\mu_1, \mu_2, \mu_3) \in \mathbb{R}^3$, $V_{\mu} = \{v \in V \mid K_j^{\pm 1}v = q^{\pm \mu_j}v, j = 1, 2, 3\}$. Let $U_q\mathfrak{k} \subset U_q\mathfrak{g}$ be the Hopf subalgebra generated by $K_2^{\pm 1}, E_j, F_j, K_j^{\pm 1}, j = 1, 3$. Every $U_q\mathfrak{g}$ -module inherits a structure of $U_q\mathfrak{k}$ -module. We are interested in quantum $(\mathfrak{g}, \mathfrak{k})$ -modules, i.e. $\mathbb{R}^3$ -weight $U_q\mathfrak{g}$ -modules which are direct sums of finite dimensional $U_q\mathfrak{k}$ -modules.

This research was partially supported by Award No UM1-2091 of the Civilian Research & Development Foundation

This lecture has been delivered at the 13-th International Hutsulian Workshop, Kiev, September 2000; published in *Matematicheskaya Fizika. Analiz. Geometriya*, **8** (2001), No 1, 90 – 110

Equip the Hopf algebra $U_q\mathfrak{g}$ with an involution:

$$\begin{aligned} E_2^* &= -K_2 F_2, & F_2^* &= -E_2 K_2^{-1}, & K_2^* &= K_2, \\ E_j^* &= K_j F_j, & F_j^* &= E_j K_j^{-1}, & K_j^* &= K_j, & j &= 1, 3. \end{aligned}$$

We thus get a $*$ -Hopf algebra $(U_q\mathfrak{g}, *)$ which is a q -analogue of $U\mathfrak{su}_{2,2}$ and its subalgebra $(U_q\mathfrak{k}, *)$ is a q -analogue of $U\mathfrak{s}(\mathfrak{u}_2 \times \mathfrak{u}_2)$.

A quantum $(\mathfrak{g}, \mathfrak{k})$ -module V is said to be unitarizable if $(\xi v_1, v_2) = (v_1, \xi^* v_2)$ for some Hermitian scalar product in V and all $v_1, v_2 \in V$, $\xi \in U_q\mathfrak{g}$. Our purpose here is to produce some series of unitarizable quantum $(\mathfrak{g}, \mathfrak{k})$ -modules by means of non-commutative geometry and non-commutative function theory in q -Cartan domains [19, 15, 16, 17, 18].

The third named author would like to express his gratitude to H. P. Jakobsen, A. Klimyk, A. Stolin and L. Turowska for helpful discussions.

2 The $U_q\mathfrak{su}_{2,2}$ -module algebra $\text{Pol}(\text{Pl}_{2,4})_{q,x}$

Let e_1, e_2, e_3, e_4 be the standard basis in $\mathbb{C}^4$. Associate to every linear operator in $\mathbb{C}^2$ its graph, a two-dimensional subspace in $\mathbb{C}^4 = \mathbb{C}^2 \times \mathbb{C}^2$, which has trivial intersection with the linear span of e_1, e_2 . We are interested in the pairs (L, ω) , with L a subspace as above and ω its non-zero volume form (an skew-symmetric bilinear form) in L . We need a q -analogue of this algebraic variety which we call the Plücker manifold $\text{Pl}_{2,4}$. The matrix elements $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ of the linear operator L , together with $t^{\pm 1}$ related to the volume element ω , work as 'coordinates' on $\text{Pl}_{2,4}$.

An algebra F is called a $U_q\mathfrak{g}$ -module algebra if the multiplication $m : F \otimes F \rightarrow F$ is a morphism of $U_q\mathfrak{g}$ -modules, and the unit $1 \in F$ is a $U_q\mathfrak{g}$ -invariant. To rephrase, one can say that for all $f_1, f_2 \in F$, $j = 1, 2, 3$,

$$\begin{aligned} E_j(f_1 f_2) &= E_j(f_1) f_2 + (K_j f_1)(E_j f_2), & E_j 1 &= 0, \\ F_j(f_1 f_2) &= (F_j f_1)(K_j^{-1} f_2) + f_1(F_j f_2), & F_j 1 &= 0, \\ K_j^{\pm 1}(f_1 f_2) &= (K_j^{\pm 1} f_1)(K_j^{\pm 1} f_2), & K_j^{\pm 1} 1 &= 1. \end{aligned}$$

In the case of a $*$ -algebra F one should impose an additional compatibility requirement for involutions:

$$(\xi f)^* = (S(\xi))^* f^*, \quad \xi \in U_q\mathfrak{g}, \quad f \in F.$$

Once the $*$ -algebra F is given by the list of its generators and relations, the $U_q\mathfrak{g}$ -module structure in F is determined unambiguously by the action of the generators $E_j, F_j, K_j^{\pm 1}$, $j = 1, 2, 3$, on the generators of F .

Consider the $*$ -algebra $\text{Pol}(\text{Mat}_2)_q$ given by its generators $\alpha, \beta, \gamma, \delta$ and the following commutation relations (the initial six of those are well known and the rest was obtained in [16]):

$$\begin{aligned} \begin{cases} \alpha\beta &= q\beta\alpha \\ \gamma\delta &= q\delta\gamma \end{cases} & \quad \begin{cases} \alpha\gamma &= q\gamma\alpha \\ \beta\delta &= q\delta\beta \end{cases} & \quad \begin{cases} \beta\gamma &= \gamma\beta \\ \alpha\delta &= \delta\alpha + (q - q^{-1})\beta\gamma \end{cases} \\ \begin{cases} \delta^*\alpha &= \alpha\delta^* \\ \delta^*\beta &= q\beta\delta^* \\ \delta^*\gamma &= q\gamma\delta^* \\ \delta^*\delta &= q^2\delta\delta^* + 1 - q^2 \end{cases} & \quad \begin{cases} \gamma^*\alpha &= q\alpha\gamma^* - (q^{-1} - q)\beta\delta^* \\ \gamma^*\beta &= \beta\gamma^* \\ \gamma^*\gamma &= q^2\gamma\gamma^* - (1 - q^2)\delta\delta^* + 1 - q^2 \end{cases} \end{aligned}$$

$$\begin{cases} \beta^* \alpha &= q\alpha\beta^* - (q^{-1} - q)\gamma\delta^* \\ \beta^* \beta &= q^2\beta\beta^* - (1 - q^2)\delta\delta^* + 1 - q^2 \end{cases}$$

$$\alpha^* \alpha = q^2\alpha\alpha^* - (1 - q^2)(\beta\beta^* + \gamma\gamma^*) + (q^{-1} - q)^2\delta\delta^* + 1 - q^2.$$

The $*$ -algebra $\text{Pol}(\text{Pl}_{2,4})_{q,x}$ is given by the generators $\alpha, \beta, \gamma, \delta, t, t^{-1}$, the commutation relations as in the above definition of $\text{Pol}(\text{Mat}_2)_q$, and the additional relations $tt^{-1} = t^{-1}t = 1$, $tt^* = t^*t$, $zt = qtz$, $zt^* = qt^*z$, with $z \in \{\alpha, \beta, \gamma, \delta\}^1$.

An application of a q -analogue for the above geometric interpretation of the Plücker manifold allows one to prove

Proposition 2.1 *i) There exists a unique structure of $U_q\mathfrak{su}_{2,2}$ -module algebra in $\text{Pol}(\text{Mat}_2)_q$ such that*

$$\begin{aligned} \begin{pmatrix} E_1\alpha & E_1\beta \\ E_1\gamma & E_1\delta \end{pmatrix} &= q^{-1/2} \begin{pmatrix} 0 & \alpha \\ 0 & \gamma \end{pmatrix}, & \begin{pmatrix} E_3\alpha & E_3\beta \\ E_3\gamma & E_3\delta \end{pmatrix} &= q^{-1/2} \begin{pmatrix} 0 & 0 \\ \alpha & \beta \end{pmatrix} \\ \begin{pmatrix} F_1\alpha & F_1\beta \\ F_1\gamma & F_1\delta \end{pmatrix} &= q^{1/2} \begin{pmatrix} \beta & 0 \\ \delta & 0 \end{pmatrix}, & \begin{pmatrix} F_3\alpha & F_3\beta \\ F_3\gamma & F_3\delta \end{pmatrix} &= q^{1/2} \begin{pmatrix} \gamma & \delta \\ 0 & 0 \end{pmatrix} \\ \begin{pmatrix} K_1\alpha & K_1\beta \\ K_1\gamma & K_1\delta \end{pmatrix} &= \begin{pmatrix} q\alpha & q^{-1}\beta \\ q\gamma & q^{-1}\delta \end{pmatrix}, & \begin{pmatrix} K_3\alpha & K_3\beta \\ K_3\gamma & K_3\delta \end{pmatrix} &= \begin{pmatrix} q\alpha & q\beta \\ q^{-1}\gamma & q^{-1}\delta \end{pmatrix} \\ \begin{pmatrix} E_2\alpha & E_2\beta \\ E_2\gamma & E_2\delta \end{pmatrix} &= -q^{1/2} \begin{pmatrix} q^{-1}\beta\gamma & \delta\beta \\ \delta\gamma & \delta^2 \end{pmatrix}, & \begin{pmatrix} F_2\alpha & F_2\beta \\ F_2\gamma & F_2\delta \end{pmatrix} &= q^{1/2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \\ \begin{pmatrix} K_2\alpha & K_2\beta \\ K_2\gamma & K_2\delta \end{pmatrix} &= \begin{pmatrix} \alpha & q\beta \\ q\gamma & q^2\delta \end{pmatrix}. \end{aligned}$$

ii) There exists a unique structure of $U_q\mathfrak{su}_{2,2}$ -module algebra in $\text{Pol}(\text{Pl}_{2,4})_{q,x}$ such that the action of $E_j, F_j, K_j^{\pm 1}$ on $\alpha, \beta, \gamma, \delta$ is given by the above equations and

$$\begin{cases} E_j t = 0 \\ F_j t = 0 \\ K_j t = t \end{cases}, \quad j = 1, 3; \quad \begin{cases} E_2 t = q^{-1/2} t \delta \\ F_2 t = 0 \\ K_2 t = q^{-1} t \end{cases}.$$

Note that a much more general result is obtained in [16].

To produce the series of quantum $(\mathfrak{g}, \mathfrak{k})$ -modules considered in the sequel we use essentially the specific dependencies of the elements $E_2 t^\lambda, F_2 t^\lambda, K_2^{\pm 1} t^\lambda, E_2((\alpha\delta - q\beta\gamma)^\lambda), F_2((\alpha\delta - q\beta\gamma)^\lambda), K_2^{\pm 1}((\alpha\delta - q\beta\gamma)^\lambda)$ on q^λ . These are easily deducible from the definitions that for all $\lambda \in \mathbb{Z}_+$

$$E_2 t^\lambda = q^{-3/2} \frac{q^{-2\lambda} - 1}{q^{-2} - 1} \delta t^\lambda, \quad F_2 t^\lambda = 0, \quad K_2^{\pm 1} t^\lambda = q^{\mp \lambda} t^\lambda,$$

$$E_2((\alpha\delta - q\beta\gamma)^\lambda) = -q^{1/2} \frac{1 - q^{2\lambda}}{1 - q^2} \delta(\alpha\delta - q\beta\gamma)^\lambda,$$

$$F_2((\alpha\delta - q\beta\gamma)^\lambda) = q^{1/2} \frac{q^{-2\lambda} - 1}{q^{-2} - 1} \alpha(\alpha\delta - q\beta\gamma)^{\lambda-1}, \quad \lambda \neq 0,$$

$$K_2^{\pm 1}((\alpha\delta - q\beta\gamma)^\lambda) = q^{\pm 2\lambda} (\alpha\delta - q\beta\gamma)^\lambda.$$

¹The notation $x = tt^*$ and $\text{Pol}(\text{Pl}_{2,4})_{q,x}$ are justified by the fact that the algebra $\text{Pol}(\text{Pl}_{2,4})_{q,x}$ in question can be derived as a localization of another useful algebra $\text{Pol}(\text{Pl}_{2,4})_q$ with respect to the multiplicative system $x^{\mathbb{N}}$.

For instance, the first relation is obvious for $\lambda = 0$, and the general case is accessible via an induction argument:

$$\begin{aligned} E_2(t^{\lambda+1}) &= (E_2 t) t^\lambda + (K_2 t)(E_2 t^\lambda) = q^{-1/2} t \delta t^\lambda + q^{-1} t q^{-3/2} \frac{q^{-2\lambda} - 1}{q^{-2} - 1} \delta t^\lambda = \\ &= \left(q^{-1/2} + q^{-5/2} \frac{q^{-2\lambda} - 1}{q^{-2} - 1} \right) q^{-1} \delta t^{\lambda+1} = q^{-3/2} \frac{q^{-2(\lambda+1)} - 1}{q^{-2} - 1} \delta t^{\lambda+1}. \end{aligned}$$

3 The analytic continuation of the holomorphic discrete series: step one

Consider the subalgebra $\mathbb{C}[\mathrm{Pl}_{2,4}]_{q,t} \subset \mathrm{Pol}(\mathrm{Pl}_{2,4})_{q,x}$ generated by $\alpha, \beta, \gamma, \delta, t, t^{-1}$. Equip it with a $\mathbb{Z}$ -grading: $\deg \alpha = \deg \beta = \deg \gamma = \deg \delta = 0$, $\deg(t^{\pm 1}) = \pm 1$. The homogeneous components of this algebra are quantum $(\mathfrak{g}, \mathfrak{k})$ -modules².

Consider the subalgebra $\mathbb{C}[\mathrm{Mat}_2]_q \subset \mathrm{Pol}(\mathrm{Mat}_2)_q$ generated by $\alpha, \beta, \gamma, \delta$. This algebra constitutes a famous subject of a research in the quantum group theory. Associate to each $\lambda \in \mathbb{Z}$ a linear operator $i_\lambda : \mathbb{C}[\mathrm{Mat}_2]_q \rightarrow \mathbb{C}[\mathrm{Pl}_{2,4}]_{q,t}$, $i_\lambda : f \mapsto f t^{-\lambda}$. This isomorphism between the vector space $\mathbb{C}[\mathrm{Mat}_2]_q$ and a homogeneous component of $\mathbb{C}[\mathrm{Pl}_{2,4}]_{q,t}$ allows one to transfer the structure of $U_q \mathfrak{sl}_4$ -module from $\mathbb{C}[\mathrm{Pl}_{2,4}]_{q,t}$ to $\mathbb{C}[\mathrm{Mat}_2]_q$. Thus we obtain a representation of $U_q \mathfrak{sl}_4$ in $\mathbb{C}[\mathrm{Mat}_2]_q$, to be denoted by π_{q^λ} . For all $\xi \in U_q \mathfrak{sl}_4$, $f \in \mathbb{C}[\mathrm{Mat}_2]_q$, the vector valued function $\pi_{q^\lambda}(\xi)f$ appears to be a Laurent polynomial of an indeterminate $\zeta = q^\lambda$. This leads to the canonical analytic continuation of the operator valued function π_{q^λ} . The term 'analytic continuation of the holomorphic discrete series' stands for the above family π_{q^λ} of representations of $U_q \mathfrak{sl}_4$.

The results of the work by H. P. Jakobsen [5] imply that the quantum $(\mathfrak{g}, \mathfrak{k})$ -modules π_{q^λ} are unitarizable for all $\lambda > 1$. We follow [17] in finding an explicit form for the related scalar product.

Consider the $\mathrm{Pol}(\mathrm{Mat}_2)_q$ -module given by a single generator v and the relations $\alpha^* v = \beta^* v = \gamma^* v = \delta^* v = 0$. The associated representation T of $\mathrm{Pol}(\mathrm{Mat}_2)_q$ in the vector space $H = \mathbb{C}[\mathrm{Mat}_2]_q v$ is faithful; it is called the vacuum representation.

Let $\check{\rho}$ be the linear operator in H that realizes the action of the 'half-sum of positive coroots':

$$\check{\rho}(\alpha^a \beta^b \gamma^c \delta^d v) = (3a + 2b + 2c + d) \alpha^a \beta^b \gamma^c \delta^d v,$$

with $a, b, c, d \in \mathbb{Z}_+$. We need also the element

$$y = 1 - (\alpha \alpha^* + \beta \beta^* + \gamma \gamma^* + \delta \delta^*) + (\alpha \delta - q \beta \gamma)(\alpha \delta - q \beta \gamma)^*,$$

which is a q -analogue of the determinant $\det(1 - \mathbf{z} \mathbf{z}^*)$, with $\mathbf{z} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$.

As a consequence of the results of [17] we have

Proposition 3.1 *i) For $\lambda > 3$ the linear functional*

$$\int_{\mathbb{U}_q} f d\nu_\lambda \stackrel{\text{def}}{=} \frac{\mathrm{tr}(T(f y^\lambda) q^{-2\check{\rho}})}{\mathrm{tr}(T(y^\lambda) q^{-2\check{\rho}})}$$

²The notation $\mathbb{C}[\mathrm{Pl}_{2,4}]_{q,t}$ can be justified in the same way as the notation $\mathrm{Pol}(\mathrm{Pl}_{2,4})_{q,x}$ introduced in the previous section.

is well defined and positive on $\text{Pol}(\text{Mat}_2)_q$.

ii) For $\lambda > 3$ the scalar product $(f_1, f_2)_{q^{2\lambda}} = \int_{\mathbb{U}_q} f_2^* f_1 d\nu_\lambda$ in $\mathbb{C}[\text{Mat}_2]_q$ is well defined, positive, and

$$(\pi_{q^\lambda}(\xi)f_1, f_2)_{q^{2\lambda}} = (f_1, \pi_{q^\lambda}(\xi^*)f_2)_{q^{2\lambda}}, \quad \xi \in U_q\mathfrak{g}, f_1, f_2 \in \mathbb{C}[\text{Mat}_2]_q.$$

The representations π_{q^λ} , $\lambda = 3, 4, 5, \dots$, are q -analogues of the holomorphic discrete series representations, and the completions of $\mathbb{C}[\text{Mat}_2]_q$ with respect to the norms $\|f\|_{q^{2\lambda}} = (f, f)_{q^{2\lambda}}^{1/2}$ are q -analogues of the weighted Bergman spaces. Our intention in what follows is to present explicit formulae for the analytic continuation of the scalar product $(f_1, f_2)_{q^{2\lambda}}$ with respect to the parameter $q^{2\lambda}$, and to prove the positivity of this scalar product for $\lambda > 1$.

To conclude, consider the $U_q\mathfrak{k}$ -invariants

$$y_1 = \alpha\alpha^* + \beta\beta^* + \gamma\gamma^* + \delta\delta^*, \quad y_2 = (\alpha\delta - q\beta\gamma)(\alpha\delta - q\beta\gamma)^*.$$

Prove that $T(y_1)T(y_2) = T(y_2)T(y_1)$, or equivalently, $y_1y_2 = y_2y_1$. In fact, observe that H admits a structure of $U_q\mathfrak{k}$ -module ($\xi(fv) = (\xi f)v$, $\xi \in U_q\mathfrak{k}$, $f \in \mathbb{C}[\text{Mat}_2]_q$) and splits into a sum of pairwise non-isomorphic simple $U_q\mathfrak{k}$ -modules $H = \bigoplus_{k_1 \geq k_2 \geq 0} H^{(k_1, k_2)}$, $H^{(k_1, k_2)} =$

$U_q\mathfrak{k}\delta^{k_1-k_2}(\alpha\delta - q\beta\gamma)^{k_2}v$. What remains is to take into account that the restrictions of $T(y_1)$, $T(y_2)$ onto $H^{(k_1, k_2)}$ are scalar operators by the 'Schur lemma'. Those scalars are easily deducible:

$$\begin{aligned} T(y_1)|_{H^{(k_1, k_2)}} &= 1 - q^{2k_1} + q^{-2}(1 - q^{2k_2}), \\ T(y_2)|_{H^{(k_1, k_2)}} &= q^{-2}(1 - q^{2k_2})(1 - q^{2(k_1+1)}). \end{aligned}$$

Just as one could expect, the joint spectrum of the operators $T(y_1)$, $T(y_2)$ tends to

$$\{(\text{tr}(zz^*), \det(zz^*)) \mid z \in \mathbb{U}\} = \{(y_1, y_2) \mid 0 \leq y_1 \leq 2 \quad \& \quad 0 \leq y_2 \leq y_1^2/4\}$$

as q goes to 1.

4 An invariant integral on the Shilov boundary

Let $c = \alpha\delta - q\beta\gamma$ and $\mathbb{C}[GL_2]_q$ be the localization of $\mathbb{C}[\text{Mat}_2]_q$ with respect to the multiplicative system $c^{\mathbb{N}}$. It is easy to prove the existence and uniqueness of an extension of the $U_q\mathfrak{g}$ -module structure from $\mathbb{C}[\text{Mat}_2]_q$ onto $\mathbb{C}[GL_2]_q$. Equip the $U_q\mathfrak{g}$ -module algebra $\mathbb{C}[GL_2]_q$ with an involution:

$$\begin{aligned} \alpha^* &= q^{-2}(\alpha\delta - q\beta\gamma)^{-1}\delta, & \beta^* &= -q^{-1}(\alpha\delta - q\beta\gamma)^{-1}\gamma, \\ \gamma^* &= -q^{-1}(\alpha\delta - q\beta\gamma)^{-1}\beta, & \delta^* &= (\alpha\delta - q\beta\gamma)^{-1}\alpha. \end{aligned}$$

and introduce the notation $\text{Pol}(S(\mathbb{U}))_q = (\mathbb{C}[GL_2]_q, *)$.

The following propositions justifies our choice of the involution.

Proposition 4.1 *For all $f \in \text{Pol}(S(\mathbb{U}))_q$, $\xi \in U_q\mathfrak{g}$ one has*

$$(\xi f)^* = (S(\xi))^* f^*.$$

Proposition 4.2 *There exists a unique homomorphism of $U_q\mathfrak{g}$ -module $*$ -algebras $j : \text{Pol}(\text{Mat}_2)_q \rightarrow \text{Pol}(S(\mathbb{U}))_q$ such that $j(\alpha) = \alpha$, $j(\beta) = \beta$, $j(\gamma) = \gamma$, $j(\delta) = \delta$.*

These statements are proved in an essentially more general form in [18]. It also follows from the results of that work that the $U_q\mathfrak{g}$ -module $*$ -algebra $\text{Pol}(S(\mathbb{U}))_q$ is a q -analogue of the polynomial algebra on the Shilov boundary $S(\mathbb{U})$ of the unit ball $\mathbb{U}$ in the space Mat_2 of complex 2×2 matrices.

The $U_q\mathfrak{k}$ -module $\text{Pol}(S(\mathbb{U}))_q$ splits into a sum of pairwise non-isomorphic simple finite dimensional submodules. In particular, the trivial $U_q\mathfrak{k}$ -module appears in $\text{Pol}(S(\mathbb{U}))_q$ with multiplicity 1 and there exists a unique $U_q\mathfrak{k}$ -invariant integral $\mu : \text{Pol}(S(\mathbb{U}))_q \rightarrow \mathbb{C}$, $f \mapsto \int_{S(\mathbb{U})_q} f d\mu$, with $\int_{S(\mathbb{U})_q} 1 d\mu = 1$.

Proposition 4.3 *The above $U_q\mathfrak{k}$ -invariant integral is positive definite.*

Proof. Consider the $*$ -algebra $\text{Pol}(U_2)_q$ of regular functions on the quantum U_2 [9], together with the $*$ -homomorphism of algebras $i : \text{Pol}(S(\mathbb{U}))_q \rightarrow \text{Pol}(U_2)_q$ given by

$$\begin{aligned} i(\alpha) &= q^{-1}\alpha, & i(\beta) &= q^{-1}\beta, \\ i(\gamma) &= \gamma, & i(\delta) &= \delta. \end{aligned}$$

The positivity of an invariant integral on the quantum group U_2 constitutes a well known fact. So, what remains is to prove the invariance of the integral

$$\text{Pol}(U_2)_q \rightarrow \mathbb{C}, \quad f \mapsto \int_{S(\mathbb{U})_q} i^{-1}(f) d\mu$$

with respect to the action of $U_q\mathfrak{u}_2$ by 'right translations' on the quantum U_2 . This is a consequence of the invariance of μ with respect to the action of the subalgebra in $U_q\mathfrak{k}$ generated by E_1 , F_1 , $K_1^{\pm 1}$, $(K_1 K_2^2 K_3)^{\pm 1}$. $\square$

Now introduce an auxiliary $U_q\mathfrak{g}$ -module $*$ -algebra $\text{Pol}(\widehat{S}(\mathbb{U}))_q$, to be used in a construction of the principal degenerate series of quantum $(\mathfrak{g}, \mathfrak{k})$ -modules.

The $*$ -algebra $\text{Pol}(\widehat{S}(\mathbb{U}))_q$ is defined by adding t , t^{-1} to the list α , β , γ , δ , c^{-1} of generators of $\text{Pol}(S(\mathbb{U}))_q$ and

$$\begin{aligned} tt^{-1} &= t^{-1}t = 1, & tt^* &= t^*t, \\ zt &= qtz, & zt^* &= qt^*z, & \text{with } z &\in \{\alpha, \beta, \gamma, \delta\} \end{aligned}$$

to the list of relations.

The next two statements follow from the results of [18].

Proposition 4.4 *i) There exists a unique extension of the structure of $U_q\mathfrak{g}$ -module $*$ -algebra from $\text{Pol}(S(\mathbb{U}))_q$ onto $\text{Pol}(\widehat{S}(\mathbb{U}))_q$ such that*

$$\begin{cases} E_j t = 0 \\ F_j t = 0 \\ K_j t = 0 \end{cases}, \quad j = 1, 3, \quad \begin{cases} E_2 t = q^{-1/2} t \delta \\ F_2 t = 0 \\ K_2 t = q^{-1} t \end{cases}.$$

ii) There exists a unique homomorphism $\widehat{j} : \text{Pol}(\text{Pl}_{2,4})_{q,x} \rightarrow \text{Pol}(\widehat{S}(\mathbb{U}))_q$ of $U_q\mathfrak{g}$ -module $$ -algebras such that*

$$\widehat{j}(\alpha) = \alpha, \quad \widehat{j}(\beta) = \beta, \quad \widehat{j}(\gamma) = \gamma, \quad \widehat{j}(\delta) = \delta, \quad \widehat{j}(t^{\pm 1}) = t^{\pm 1}.$$

Proposition 4.5 *The subspace $t^{*-2}\text{Pol}(S(\mathbb{U}))_q t^{-2}$ is a submodule of the $U_q\mathfrak{g}$ -module $\text{Pol}(\widehat{S}(\mathbb{U}))_q$, and the linear functional*

$$t^{*-2}\text{Pol}(S(\mathbb{U}))_q t^{-2} \rightarrow \mathbb{C}, \quad t^{*-2}ft^{-2} \mapsto \int_{S(\mathbb{U})_q} f d\mu$$

is an invariant integral (i.e. a morphism of $U_q\mathfrak{g}$ -modules).

5 An analytic continuation of the holomorphic discrete series: step two

Just as in the classical case $q = 1$, one has

$$\mathbb{C}[\text{Mat}_2]_q = \bigoplus_{k_1 \geq k_2 \geq 0} \mathbb{C}[\text{Mat}_2]_q^{(k_1, k_2)} = U_q\mathfrak{k}\delta^{k_1-k_2}(\alpha\delta - q\beta\gamma)^{k_2},$$

with $\mathbb{C}[\text{Mat}_2]_q^{(k_1, k_2)}$ being simple pairwise non-isomorphic $U_q\mathfrak{k}$ -submodules of the $U_q\mathfrak{k}$ -module $\mathbb{C}[\text{Mat}_2]_q$. Introduce the notation $f^{(k_1, k_2)}$ for a projection of f onto the $U_q\mathfrak{k}$ -isotypic component $\mathbb{C}[\text{Mat}_2]_q^{(k_1, k_2)}$ parallel to the sum of all other $U_q\mathfrak{k}$ -isotypic components.

By the 'Schur lemma', every $U_q\mathfrak{k}$ -invariant Hermitian form (f_1, f_2) on $\mathbb{C}[\text{Mat}_2]_q$ is given by

$$(f_1, f_2) = \sum_{k_1 \geq k_2 \geq 0} c(k_1, k_2) \int_{S(\mathbb{U})_q} (f_2^{(k_1, k_2)})^* f_1^{(k_1, k_2)} d\mu.$$

We are going to obtain this decomposition for $(f_1, f_2)_{q^{2\lambda}}$, $\lambda > 3$. Recall the notation $(a; q^2)_m = \prod_{j=0}^{m-1} (1 - aq^{2j})$.

Proposition 5.1 *For all $\lambda > 3$, $f_1, f_2 \in \mathbb{C}[\text{Mat}_2]_q$,*

$$\int_{\mathbb{U}_q} f_2^* f_1 d\nu_\lambda = \sum_{k_1 \geq k_2 \geq 0} c(k_1, k_2, q^{2\lambda}) \int_{S(\mathbb{U})_q} (f_2^{(k_1, k_2)})^* f_1^{(k_1, k_2)} d\mu,$$

with

$$c(k_1, k_2, q^{2\lambda}) = \frac{(q^4; q^2)_{k_1} (q^2; q^2)_{k_2}}{(q^{2\lambda}; q^2)_{k_1} (q^{2(\lambda-1)}; q^2)_{k_2}}. \quad (5.1)$$

Proof. In the case $q = 1$ a similar result was obtained by Faraut and Koranyi [3] in a very big generality. Our proof here imitates that of [3].

First introduce the subalgebra $\mathbb{C}[\overline{\text{Mat}}_2]_q \subset \text{Pol}(\text{Mat}_2)_q$ generated by $\alpha^*, \beta^*, \gamma^*, \delta^*$, and the algebra $\mathbb{C}[\overline{\text{Mat}}_2]_q^{\text{op}}$ which differs from $\mathbb{C}[\overline{\text{Mat}}_2]_q$ by replacement of the multiplication law with an opposite one. We use the algebra $\mathbb{C}[\text{Mat}_2 \times \overline{\text{Mat}}_2]_q \stackrel{\text{def}}{=} \mathbb{C}[\text{Mat}_2]_q \otimes \mathbb{C}[\overline{\text{Mat}}_2]_q^{\text{op}}$ as a q -analogue for the algebra of (degenerate) kernels of integral operators.

Equip $\mathbb{C}[\text{Mat}_2 \times \overline{\text{Mat}}_2]_q$ with a bigrading

$$\deg(\alpha \otimes 1) = \deg(\beta \otimes 1) = \deg(\gamma \otimes 1) = \deg(\delta \otimes 1) = (1, 0),$$

$$\deg(1 \otimes \alpha^*) = \deg(1 \otimes \beta^*) = \deg(1 \otimes \gamma^*) = \deg(1 \otimes \delta^*) = (0, 1)$$

and the associated topology. The completed algebra $\mathbb{C}[[\text{Mat}_2 \times \overline{\text{Mat}_2}]]_q$ will work as the algebra of generalized kernels of integral operators [17].

Just as in the case $q = 1$ one can deduce proposition 5.1 from the following three lemmas.

Lemma 5.2 *Given $k_1, k_2 \in \mathbb{Z}$, $k_1 \geq k_2 \geq 0$, denote by P_{k_1, k_2} the projection in $\mathbb{C}[\text{Mat}_2]_q$ onto the component $\mathbb{C}[\text{Mat}_2]_q^{(k_1, k_2)}$ parallel to the sum of all other $U_q \mathfrak{k}$ -isotypic components. There exists a unique element $p_{k_1, k_2} \in \mathbb{C}[\text{Mat}_2 \times \overline{\text{Mat}_2}]_q$ such that*

$$P_{k_1, k_2} f(z) = \int_{S(\mathbb{U})_q} p_{k_1, k_2}(z, \zeta) f(\zeta) d\mu(\zeta)$$

for all $f \in \mathbb{C}[\text{Mat}_2]_q$.

Introduce the notation $L^2(d\nu_\lambda)_q$, $L_a^2(d\nu_\lambda)_q$ for completions of vector spaces $\text{Pol}(\text{Mat}_2)_q$, $\mathbb{C}[\text{Mat}_2]_q$ respectively, with respect to the norm $\|f\|_{q^{2\lambda}} = \left(\int_{\mathbb{U}_q} f^* f d\nu_\lambda \right)^{1/2}$. These are well defined for $\lambda > 3$, and certainly $L_a^2(d\nu_\lambda)_q \subset L^2(d\nu_\lambda)_q$.

Lemma 5.3 *Given $\lambda > 3$, denote by P_λ the orthogonal projection in $L^2(d\nu_\lambda)_q$ onto $L_a^2(d\nu_\lambda)_q$. There exists a unique $K_\lambda \in \mathbb{C}[[\text{Mat}_2 \times \overline{\text{Mat}_2}]]_q$ such that*

$$P_\lambda f(z) = \int_{\mathbb{U}_q} K_\lambda(z, \zeta) f(\zeta) d\nu_\lambda(\zeta).$$

for all $f \in \text{Pol}(\text{Mat}_2)_q$.

Lemma 5.4 *In $\mathbb{C}[[\text{Mat}_2 \times \overline{\text{Mat}_2}]]_q$ one has*

$$K_\lambda = \sum_{k_1 \geq k_2 \geq 0} \frac{1}{c(k_1, k_2, \lambda)} p_{k_1, k_2},$$

with $c(k_1, k_2, \lambda)$ being given by (5.1).

Lemmas 5.2, 5.3 can be proved in the same way as in the case $q = 1$. Turn to the proof of lemma 5.4.

We are going to use the Schur polynomials

$$s_{k_1 k_2}(x_1, x_2) = (x_1 x_2)^{k_2} \cdot \frac{x_1^{k_1 - k_2 + 1} - x_2^{k_1 - k_2 + 1}}{x_1 - x_2}.$$

These are expressible in terms of elementary symmetric polynomials:

$$s_{k_1 k_2}(x_1, x_2) = u_{k_1 k_2}(x_1 + x_2, x_1 x_2).$$

(The polynomials $u_{k_1 k_2}$ are closely related to the well known Chebyshev polynomials of second kind $U_{k_1 - k_2}(x)$).

Recall the notation $[j]_q = \frac{q^j - q^{-j}}{q - q^{-1}}$, $(a; q^2)_\infty = \prod_{j=0}^{\infty} (1 - a q^{2j})$ and consider the kernels $\chi_1 = \alpha \otimes \alpha^* + \beta \otimes \beta^* + \gamma \otimes \gamma^* + \delta \otimes \delta^*$, $\chi_2 = c \otimes c^*$ with $c = (\alpha\delta - q\beta\gamma) \in \text{Pol}(\text{Mat}_2)_q$.

Lemma 5.5 *i)* $p_{k_1, k_2} = q^{k_1 + k_2} \cdot [k_1 - k_2 + 1]_q \cdot u_{k_1 k_2}(\chi_1, \chi_2),$
ii) $K_\lambda = \prod_{j=0}^{\infty} (1 - q^{2(\lambda+j)} \chi_1 + q^{4(\lambda+j)} \chi_2) \left(\prod_{j=0}^{\infty} (1 - q^{2j} \chi_1 + q^{4j} \chi_2) \right)^{-1}.$

The first statement of lemma 5.5 are easily deducible from the orthogonality relations for matrix elements of representations of the quantum group U_2 . The second statement follows from the results of [17].

Lemma 5.4 is a consequence of lemma 5.5 and the following well known relation in the theory of Schur polynomials [12]:

$$\begin{aligned} \frac{(q^{2\lambda} x_1; q^2)_\infty}{(x_1; q^2)_\infty} \cdot \frac{(q^{2\lambda} x_2; q^2)_\infty}{(x_2; q^2)_\infty} &= \\ &= \sum_{k_1 \geq k_2 \geq 0} \frac{(q^{2\lambda}; q^2)_{k_1} (q^{2(\lambda-1)}; q^2)_{k_2}}{(q^4; q^2)_{k_1} (q^2; q^2)_{k_2}} [k_1 - k_2 + 1]_q \cdot q^{(k_1 + k_2)} s_{k_1 k_2}(x_1, x_2). \end{aligned}$$

The above proof of proposition 5.1 is transferable quite literally onto the case of quantum $SU_{n,n}$ and a q-analogue of the unit ball in the space of $n \times n$ matrices.

6 Analytic continuation of the holomorphic discrete series: ladder representation of the quantum group $SU_{2,2}$

It is explained in [3] that the results like our proposition 5.1 allow one to solve the problems of irreducibility, unitarizability, and composition series of the representations π_{q^λ} . We restrict ourselves to some simplest corollaries from proposition 5.1.

Proposition 6.1 *Suppose that either $\lambda > 1$ or $\text{Im } \lambda \in \frac{\pi}{\lg q} \mathbb{Z}$. Then the sesquilinear form $(f_1, f_2)_{q^{2\lambda}}$ is positive definite, and for all $f_1, f_2 \in \mathbb{C}[\text{Mat}_2]_q$, $\xi \in U_q \mathfrak{g}$ one has*

$$(\pi_{q^\lambda}(\xi) f_1, f_2)_{q^{2\lambda}} = (f_1, \pi_{q^\lambda}(\xi^*) f_2)_{q^{2\lambda}}. \quad (6.1)$$

Proof. The positivity follows from proposition 5.1. Let $\zeta = q^\lambda$. If $\text{Im } \zeta = 0$, both sides of (6.1) are rational functions of ζ . So, what remains is to use the fact that this equality is true for $0 < \zeta < q^3$. $\square$

Turn to the case $\lambda = 1$. It follows from proposition 6.1 that the kernel of the sesquilinear form $\langle f_1, f_2 \rangle = \lim_{\lambda \rightarrow 1+0} (1 - q^{2\lambda-2})(f_1, f_2)_{q^{2\lambda}}$ is a common invariant subspace for all the operators $\pi_1(\xi)$, $\xi \in U_q \mathfrak{g}$. Explicitly, this kernel is

$$L = \bigoplus_{k=0}^{\infty} \mathbb{C}[\text{Mat}_2]_q^{(k,0)}.$$

On L one has a well defined Hermitian form $(f_1, f_2) = \lim_{\lambda \rightarrow 1+0} (f_1, f_2)_{q^{2\lambda}}$, and hence the quantum $(\mathfrak{g}, \mathfrak{k})$ -module associated to the restriction $\pi_q|_L$ is unitarizable. The representation $\pi_q|_L$ is a q-analogue of the well known *ladder representation*.

In the case $q = 1$ the subspace $\bigoplus_{k=0}^{\infty} \mathbb{C}[\text{Mat}_2]^{(k,0)}$ coincides with the kernel of the covariant differential operator $\square = \frac{\partial}{\partial \alpha} \frac{\partial}{\partial \delta} - \frac{\partial}{\partial \beta} \frac{\partial}{\partial \gamma}$. Our intention is to obtain a q-analogue of this result³.

³A similar result was obtained by V. Dobrev [2] and H. P. Jakobsen [6] in a different context.

We use a notion of the first order differential calculus over an algebra A and a covariant first order differential calculus as in [10].

Among well known $U_q\mathfrak{k}$ -invariant first order differential calculi over $\mathbb{C}[\mathrm{Mat}_2]_q$ one has to distinguish a unique $U_q\mathfrak{g}$ -invariant calculus. A general method of producing such differential calculi (with hidden symmetry) for q -Cartan domains is described in [19].

The first order differential calculus we need is determined by the following 'commutation relations between coordinates and differentials' (these are written in [16] in R-matrix form):

$$\begin{aligned}
 d\alpha \cdot \alpha &= q^2 \alpha d\alpha; & d\alpha \cdot \beta &= q\beta d\alpha - (1 - q^2)\alpha d\beta; \\
 d\alpha \cdot \gamma &= q\gamma d\alpha - (1 - q^2)\alpha d\gamma; & d\alpha \cdot \delta &= \delta d\alpha - (q^{-1} - q)(\gamma d\beta + \beta d\gamma) + (q^{-1} - q)^2 \alpha d\delta; \\
 d\beta \cdot \alpha &= q\alpha \cdot d\beta; & d\beta \cdot \beta &= q^2 \beta d\beta; \\
 d\beta \cdot \gamma &= \gamma d\beta - (q^{-1} - q)\alpha d\delta; & d\beta \cdot \delta &= q\delta d\beta - (1 - q^2)\beta d\delta; \\
 d\gamma \cdot \alpha &= q\alpha d\gamma; & d\gamma \cdot \gamma &= q^2 \gamma d\gamma; \\
 d\gamma \cdot \beta &= \beta d\gamma - (q^{-1} - q)\alpha d\delta; & d\gamma \cdot \delta &= q\delta d\gamma - (1 - q^2)\gamma d\delta; \\
 d\delta \cdot \alpha &= \alpha d\delta; & d\delta \cdot \gamma &= q\gamma d\delta; \\
 d\delta \cdot \beta &= q\beta d\delta; & d\delta \cdot \delta &= q^2 \delta d\delta.
 \end{aligned}$$

It is worthwhile to note that it admits an extension up to a $U_q\mathfrak{g}$ -module first order differential calculus over $U_q\mathfrak{g}$ -module algebra $\mathbb{C}[\mathrm{Pl}_{2,4}]_{q,t}$: $dt \cdot t = q^{-2}t dt$,

$$dz \cdot t = q^{-1}t dz, \quad dt \cdot z = q^{-1}z dt + (q^{-2} - 1)t dz \quad \text{for all } z \in \{\alpha, \beta, \gamma, \delta\}.$$

Turn back to $\mathbb{C}[\mathrm{Mat}_2]_q$. The operator d is given on the generators of this algebra in an obvious way and is extended onto the entire algebra via the Leibnitz rule. The operators $\frac{\partial}{\partial \alpha}, \frac{\partial}{\partial \beta}, \frac{\partial}{\partial \gamma}, \frac{\partial}{\partial \delta}$ in $\mathbb{C}[\mathrm{Mat}_2]_q$ are imposed in a standard way:

$$df = \frac{\partial f}{\partial \alpha} d\alpha + \frac{\partial f}{\partial \beta} d\beta + \frac{\partial f}{\partial \gamma} d\gamma + \frac{\partial f}{\partial \delta} d\delta.$$

As an easy consequence of the definitions one has

Proposition 6.2 *Let $\square_q = \frac{\partial}{\partial \alpha} \frac{\partial}{\partial \delta} - q \frac{\partial}{\partial \beta} \frac{\partial}{\partial \gamma}$.*

i) $\square_q$ intertwines π_q and π_{q^3} :

$$\pi_{q^3}(\xi) \square_q = \square_q \pi_q(\xi), \quad \xi \in U_q\mathfrak{g}.$$

ii) $L = \mathrm{Ker} \square_q$.

$$iii) (\alpha\delta - q\beta\gamma) \square_q|_{\mathbb{C}[\mathrm{Mat}_2]_q^{(k_1, k_2)}} = q^{-2} \cdot \frac{1 - q^{2k_2}}{1 - q^2} \cdot \frac{1 - q^{2(k_1+1)}}{1 - q^2}.$$

Corollary 6.3 *For all $s \in \mathbb{N}$*

$$\square_q(\alpha\delta - q\beta\gamma)^s = b_q(s)(\alpha\delta - q\beta\gamma)^{s-1},$$

$$b_q(s) = q^{-2} \cdot \frac{1 - q^{2s}}{1 - q^2} \cdot \frac{1 - q^{2(s+1)}}{1 - q^2}.$$

$b_q(s)$ is a q -analogue of the Sato-Bernstein polynomial $b(s) = s(s+1)$ for the prehomogeneous vector space Mat_2 . In a recent preprint [7] and the works cited therein, another approach to q -analogues for algebras of differential operators was used to produce q -analogues of the Bernstein polynomials.

Consider the vector space $\mathbb{C}^4$ (with its standard coordinate system t_1, t_2, t_3, t_4), together with the associated projective space $\mathbb{CP}^3$. Let $\mathbb{L} \subset \mathbb{CP}^3$ be a projectivization of the plane $t_3 = t_4 = 0$. It is well known that in the case $q = 1$ the ladder representation is isomorphic to the natural representation of $U\mathfrak{g}$ in the cohomologies $H^1(\mathbb{CP}^3 \setminus \mathbb{L}, \mathcal{O}(-2))$. A computation of these cohomologies by the Čech method leads to the Laurent polynomials:

$$H^1(\mathbb{CP}^3 \setminus \mathbb{L}, \mathcal{O}(-2)) = \left\{ \sum_{(j_1, j_2, j_3, j_4) \in J} c_{j_1, j_2, j_3, j_4} t_1^{j_1} t_2^{j_2} t_3^{j_3} t_4^{j_4} \right\},$$

with $J = \{(j_1, j_2, j_3, j_4) \in \mathbb{Z}^4 \mid j_1 \geq 0, j_2 \geq 0, j_3 < 0, j_4 < 0, j_1 + j_2 + j_3 + j_4 = -2\}$.

So, one has two geometric realizations of the ladder representation of $SU_{2,2}$ (those in $H^1(\mathbb{CP}^3 \setminus \mathbb{L}, \mathcal{O}(-2))$ and in $\text{Ker } \square$).

The lowest weight subspace in $H^1(\mathbb{CP}^3 \setminus \mathbb{L}, \mathcal{O}(-2))$ is generated by the Laurent polynomial $t_3^{-1}t_4^{-1}$, and in the kernel of $\square = \frac{\partial}{\partial \alpha} \frac{\partial}{\partial \delta} - \frac{\partial}{\partial \beta} \frac{\partial}{\partial \gamma}$ by the constant function 1. There exists a unique isomorphism between the two realizations of the ladder representation which takes $t_3^{-1}t_4^{-1}$ to 1. This operator is very essential in the mathematical physics and is called the Penrose transform [1]. A replacement of the commutation relation $t_i t_j = t_j t_i$ by $t_i t_j = q t_j t_i$, $i < j$, allows one to transfer easily the above observations onto the case $0 < q < 1$ (more precisely, everything but the notion of cohomologies for quasi-coherent sheaves). It is just the way of on which another realization of the ladder representation and the quantum Penrose transform appear.

7 The principal degenerate series of quantum Harish-Chandra modules

In the classical theory the principal series of Harish-Chandra modules are associated to parabolic subgroups P . Our purpose is to produce a q -analogue of the principal series of Harish-Chandra modules associated to a stability group P for a point of the Shilov boundary $p \in S(\mathbb{U})$.

We call a $U_q \mathfrak{g}$ -module V $\mathbb{Z}^3$ -weight module if $V = \bigoplus_{\mu} V_{\mu}$ with $\mu = (\mu_1, \mu_2, \mu_3) \in \mathbb{Z}^3$, $V_{\mu} = \{v \in V \mid K_j^{\pm 1} v = q^{\pm \mu_j} v, j = 1, 2, 3\}$.

A quantum Harish-Chandra module is a finitely generated $\mathbb{Z}^3$ -weight $U_q \mathfrak{g}$ -module V such that

- i) V is a sum of finite dimensional simple $U_q \mathfrak{k}$ -modules,
- ii) each simple finite dimensional $U_q \mathfrak{k}$ -module W occurs in V with finite multiplicity ($\dim \text{Hom}_{U_q \mathfrak{k}}(W, V) < \infty$).

Quantum Harish-Chandra modules are quantum $(\mathfrak{g}, \mathfrak{k})$ -modules, and the notion of unitarizability is applicable here. The rest of this section is devoted to producing the principal degenerate series of the unitarizable quantum Harish-Chandra modules. Note that producing and classification of simple unitarizable quantum Harish-Chandra modules still constitute an open problem even in our special case of quantum $SU_{2,2}$.

In the case $\lambda \in -2\mathbb{Z}_+$ one has a well defined linear operator $\text{Pol}(S(\mathbb{U}))_q \rightarrow \text{Pol}(\widehat{S}(\mathbb{U}))_q$, $f \mapsto f \cdot (\alpha\delta - q\beta\gamma)^{-\lambda/2} t^{-\lambda}$. The same argument as that applied in section 3 to produce π_{q^λ} , yields

Proposition 7.1 *There exists a unique one-parameter family τ_{q^λ} of representations of $U_q\mathfrak{g}$ in the space $\text{Pol}(S(\mathbb{U}))_q$ of polynomials on the Shilov boundary of the quantum matrix ball such that*

i) for all $\lambda \in -2\mathbb{Z}_+$, $\xi \in U_q\mathfrak{g}$, $f \in \text{Pol}(S(\mathbb{U}))_q$ one has

$$(\tau_{q^\lambda}(\xi)f)(\alpha\delta - q\beta\gamma)^{-\lambda/2}t^{-\lambda} = \xi(f(\alpha\delta - q\beta\gamma)^{-\lambda/2}t^{-\lambda});$$

ii) for all $\xi \in U_q\mathfrak{g}$, $f \in \text{Pol}(S(\mathbb{U}))_q$, the vector function $\tau_{q^\lambda}(\xi)f$ is a Laurent polynomial of the indeterminate $\zeta = q^\lambda$.

Note that the multiple $(\alpha\delta - q\beta\gamma)^{-\lambda/2}$ provides the integral nature for weight of τ_{q^λ} . We are to produce a q -analogue of the principal degenerate series of Harish-Chandra modules associated to the Shilov boundary $S(\mathbb{U})$.

REMARK. For $q = 1$ the construction of degenerate discrete series involves a finite dimensional irreducible representation τ of a reductive subgroup M . (This subgroup is determined in a standard way:

$$P = MAN, \quad S(\mathbb{U}) \approx P \setminus SU_{2,2}.$$

We have produced q -analogues of those representations of degenerate discrete series where τ is trivial, i.e. a q -analogue of the spherical principal degenerate series:

$$\tau_{q^\lambda}(\xi)1 = \varepsilon(\xi)1, \quad \xi \in U_q\mathfrak{k}.$$

Turn to a construction of the corresponding principal unitary series.

Proposition 7.2 *In the case $\text{Re } \lambda = 2$ the quantum Harish-Chandra module associated to τ_{q^λ} is unitarizable:*

$$\int_{S(\mathbb{U})_q} f_2^*(\tau_{q^\lambda}(\xi)f_1)d\mu = \int_{S(\mathbb{U})_q} (\tau_{q^\lambda}(\xi^*)f_2)^*f_1d\mu \quad (7.1)$$

for all $f_1, f_2 \in \text{Pol}(S(\mathbb{U}))_q$, $\xi \in U_q\mathfrak{g}$.

Proof. The representation τ_{q^λ} can be defined in a different way, as one can extend the $U_q\mathfrak{g}$ -module algebra $\text{Pol}(\widehat{S}(\mathbb{U}))_q$ via adding to the list of generators the powers t^λ , $(t^*)^\lambda$, $(\alpha\delta - q\beta\gamma)^\lambda$ for any $\lambda \in \mathbb{C}$. The relations between the generators of the extended algebra as well as the action of E_j , F_j , $K_j^{\pm 1}$, $j = 1, 2, 3$, on them are derived from the corresponding formulae for integral powers of t , t^* , and $\alpha\delta - q\beta\gamma$ via the analytic continuation which uses Laurent polynomials of the indeterminate $\zeta = q^\lambda$. Moreover, this new algebra may be endowed with an involution as follows

$$(t^\lambda)^* = (t^*)^{\bar{\lambda}}, \quad ((\alpha\delta - q\beta\gamma)^\lambda)^* = q^{-2\bar{\lambda}} \cdot (\alpha\delta - q\beta\gamma)^{-\bar{\lambda}}$$

(where bar denotes the complex conjugation), and thus it is made a $U_q\mathfrak{g}$ -module $*$ -algebra.

Now the relation (7.1) follows from

Lemma 7.3 *Let $\operatorname{Re} \lambda = 2$. The linear subspace*

$$((\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda})^* \cdot \operatorname{Pol}(S(\mathbb{U}))_q \cdot (\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda}$$

is a $U_q\mathfrak{g}$ -module, and the linear functional

$$((\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda})^* \cdot f \cdot (\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda} \mapsto \int_{S(\mathbb{U})_q} f d\mu$$

is a $U_q\mathfrak{g}$ -invariant integral.

Proof of lemma 7.3. Suppose that $\lambda = 2 + i\rho$ with $\rho \in \mathbb{R}$. Then, by definitions,

$$\begin{aligned} & ((\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda})^* \cdot f \cdot (\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda} = \\ & (t^*)^{-\bar{\lambda}} \cdot ((\alpha\delta - q\beta\gamma)^*)^{-\bar{\lambda}/2} \cdot f \cdot (\alpha\delta - q\beta\gamma)^{-\lambda/2} \cdot t^{-\lambda} = \\ & = \operatorname{const}(\rho) \cdot (t^*)^{-2} \cdot (t^*)^{i\rho} \cdot (\alpha\delta - q\beta\gamma)^{-i\rho} \cdot f \cdot t^{-i\rho} \cdot t^{-2}. \end{aligned}$$

Now it suffices to apply proposition 4.5, the equality

$$\int_{S(\mathbb{U})_q} f d\mu = \int_{S(\mathbb{U})_q} f^{(0,0)} d\mu,$$

and the observation that the element $t^*t^{-1}(\alpha\delta - q\beta\gamma)^{-1} \in \operatorname{Pol}(\widehat{S}(\mathbb{U}))_q$ commutes with any element of the subalgebra $\operatorname{Pol}(S(\mathbb{U}))_q$ and is a $U_q\mathfrak{g}$ -invariant.

A construction of the second part τ'_{q^λ} of the principal degenerate series of quantum Harish-Chandra modules we are interested in is described in the following proposition. Its proof is just the same as that of proposition 7.1.

Proposition 7.4 *There exists a unique one-parameter family τ'_{q^λ} of representations of $U_q\mathfrak{g}$ in the space $\operatorname{Pol}(S(\mathbb{U}))_q$ such that*

i) for all $\lambda \in -2\mathbb{Z}_+$, $\xi \in U_q\mathfrak{g}$, $f \in \operatorname{Pol}(S(\mathbb{U}))_q$, one has

$$(\tau'_{q^\lambda}(\xi)f)(\alpha\delta - q\beta\gamma)^{-\lambda/2}t^{-\lambda-1} = \xi(f(\alpha\delta - q\beta\gamma)^{-\lambda/2}t^{-\lambda-1})$$

ii) for all $\xi \in U_q\mathfrak{g}$, $f \in \operatorname{Pol}(S(\mathbb{U}))_q$, the vector function $\tau'_{q^\lambda}(\xi)f$ is a Laurent polynomial of the indeterminate $\zeta = q^\lambda$.

REMARK. Both parts τ_{q^λ} , τ'_{q^λ} of the series of quantum Harish-Chandra modules in question could be also derived via embeddings of vector spaces $\operatorname{Pol}(S(\mathbb{U}))_q \rightarrow \operatorname{Pol}(\widehat{S}(\mathbb{U}))_q$, $f \mapsto ft^{l_1}t^{*l_2}$. For that, with $l_1 - l_2 \in \mathbb{Z}$ being fixed, one should arrange 'an analytic continuation in $\zeta = q^{l_1+l_2}$ '. An equivalence of the two above approaches to producing the principal degenerate series follows from properties of the element $t^*t^{-1}(\alpha\delta - q\beta\gamma)^{-1}$ (see proof of lemma 7.3).

8 The principal non-degenerate series of quantum Harish-Chandra modules

The finite dimensional simple weight $U_q\mathfrak{g}$ -modules allow a plausible description in terms of generators and relations when the highest weight vectors are chosen as generators. In the infinite dimensional case the capability of this approach is much lower. The well known method of inducing from a parabolic subgroup in our case is also inapplicable due to the absence of a valuable q -analogue of the Iwasawa decomposition.

Fortunately, there exists one more approach to a description of Harish-Chandra modules, that of Beilinson and Bernstein [14]. Within the framework of this approach simple Harish-Chandra modules are produced in cohomologies with supports on K -orbits in the space of full flags $X = G/B$ (in our case $G = SL_4$, $K = S(GL_2 \times GL_2)$, and B a standard Borel subgroup). The principal non-degenerate series is related to an open orbit, which is an affine algebraic variety. This fact sharply simplifies the problem of producing the principal non-degenerate series, and makes it possible to solve the problem for the quantum case.

An application of the results of Kostant [8] allows one to obtain an analogue of proposition 4.5 for full flags and to prove that, together with every quantum Harish-Chandra module, the principal non-degenerate series contains its dual.

Appendix 1. A complete list of irreducible $*$ -representations of $\text{Pol}(\text{Mat}_2)_q$

This appendix presents an outline of the results of L. Turowska [20] on classification of irreducible $*$ -representations of $\text{Pol}(\text{Mat}_2)_q$.

To forestall the exposition, note that every irreducible representation from the list of L. Turowska possesses a distinguished vector v (determined up to a scalar multiple) and is a completion of the $\text{Pol}(\text{Mat}_2)_q$ -module $V = \text{Pol}(\text{Mat}_2)_q v$ with respect to a suitable topology. Our intention is to produce the list of relations which determine the above $\text{Pol}(\text{Mat}_2)_q$ -modules. As one can observe from the results of L. Turowska, the non-negative linear functionals

$$l_q : \text{Pol}(\text{Mat}_2)_q \rightarrow \mathbb{C}, \quad l_q : f \mapsto (fv, v)$$

lead in the classical limit $q \rightarrow 1$ to non-negative linear functionals on the polynomial algebra $\text{Pol}(\text{Mat}_2)$. The limit functionals are just the delta-functions in some points of the closure of the unit ball $\mathbb{U}$.

We list below those points, together with the lists of determining relations for the associated $\text{Pol}(\text{Mat}_2)_q$ -modules⁴.

0-dimensional leaves

$$\begin{pmatrix} e^{i\varphi_1} & 0 \\ 0 & e^{i\varphi_2} \end{pmatrix} \quad \begin{aligned} \alpha v &= e^{i\varphi_1} v, & \beta v &= 0, & \gamma v &= 0, & \delta v &= e^{i\varphi_2} v, \\ \alpha^* v &= e^{-i\varphi_1} v, & \beta^* v &= 0, & \gamma^* v &= 0, & \delta^* v &= e^{-i\varphi_2} v, \end{aligned} \quad \varphi_1, \varphi_2 \in \mathbb{R}/2\pi\mathbb{Z}.$$

2-dimensional leaves

$$\begin{pmatrix} 0 & 0 \\ 0 & e^{i\varphi} \end{pmatrix} \quad \begin{aligned} \beta v &= 0, & \gamma v &= 0, & \delta v &= e^{i\varphi} v, \\ \alpha^* v &= 0, & \beta^* v &= 0, & \gamma^* v &= 0, & \delta^* v &= e^{-i\varphi} v, \end{aligned} \quad \varphi \in \mathbb{R}/2\pi\mathbb{Z}.$$

⁴Consider the Poisson bracket $\{f_1, f_2\} = \lim_{h \rightarrow 0} \frac{f_1 f_2 - f_2 f_1}{ih}$, $h = 2\log(q^{-1})$, and associate to each of those points a bounded symplectic leaf containing this point. An important invariant of the irreducible $*$ -representation is the dimension of the associated symplectic leaf.

$$\begin{pmatrix} 0 & e^{i\varphi_1} \\ e^{i\varphi_2} & 0 \end{pmatrix} \quad \begin{aligned} \alpha v &= 0, & \beta v &= e^{i\varphi_1} v, & \gamma v &= e^{i\varphi_2} v, \\ \alpha^* v &= -q^{-1} e^{-i(\varphi_1 + \varphi_2)} \delta v, & \beta^* v &= e^{-i\varphi_1} v, & \gamma^* v &= e^{-i\varphi_2} v, & \delta^* v &= 0, \\ \varphi_1, \varphi_2 &\in \mathbb{R}/2\pi\mathbb{Z}. \end{aligned}$$

4-dimensional leaves

$$\begin{pmatrix} 0 & 0 \\ e^{i\varphi} & 0 \end{pmatrix} \quad \begin{aligned} \alpha v &= 0, & \gamma v &= e^{i\varphi} v, \\ \alpha^* v &= 0, & \beta^* v &= 0, & \gamma^* v &= e^{-i\varphi} v, & \delta^* v &= 0, & \varphi &\in \mathbb{R}/2\pi\mathbb{Z}. \end{aligned}$$

$$\begin{pmatrix} 0 & e^{i\varphi_1} \\ 0 & 0 \end{pmatrix} \quad \begin{aligned} \alpha v &= 0, & \beta v &= e^{i\varphi} v, \\ \alpha^* v &= 0, & \beta^* v &= e^{-i\varphi} v, & \gamma^* v &= 0, & \delta^* v &= 0, & \varphi &\in \mathbb{R}/2\pi\mathbb{Z}. \end{aligned}$$

6-dimensional leaves

$$\begin{pmatrix} e^{i\varphi} & 0 \\ 0 & 0 \end{pmatrix} \quad \begin{aligned} \alpha v &= e^{i\varphi} v, \\ \alpha^* v &= e^{-i\varphi} v, & \beta^* v &= \gamma^* v = \delta^* v = 0, & \varphi &\in \mathbb{R}/2\pi\mathbb{Z}. \end{aligned}$$

8-dimensional leaf

$$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \quad \alpha^* v = \beta^* v = \gamma^* v = \delta^* v = 0.$$

It follows from the results of L. Turowska that every of the above $\text{Pol}(\text{Mat}_2)_q$ -modules can be equipped with a structure of pre-Hilbert space in such a way that the $\text{Pol}(\text{Mat}_2)_q$ -action is extendable onto the associated Hilbert space, and this procedure provides a complete list of irreducible $*$ -representations of $\text{Pol}(\text{Mat}_2)_q$ ⁵.

Note that the $*$ -representation associated to the 8-dimensional symplectic leaf is faithful; it is unique (up to a unitary equivalence) faithful irreducible $*$ -representation. The uniqueness is easily deducible from the commutation relations between $\alpha, \beta, \gamma, \delta, \alpha^*, \beta^*, \gamma^*, \delta^*, y$ (the later element is defined in section 3).

Another two series of $*$ -representations are related to the leaves that contain unitary matrices

$$\begin{pmatrix} e^{i\varphi_1} & 0 \\ 0 & e^{i\varphi_2} \end{pmatrix}, \quad \begin{pmatrix} 0 & e^{i\varphi_1} \\ e^{i\varphi_2} & 0 \end{pmatrix}, \quad \varphi_1, \varphi_2 \in \mathbb{R}/2\pi\mathbb{Z}.$$

These two series are due to the $*$ -homomorphism $\text{Pol}(\text{Mat}_2)_q \rightarrow \mathbb{C}[U_2]_q$ described in the main sections of this work. They could be obtained within the theory of $*$ -representations of the algebra $\mathbb{C}[U_2]_q$ of regular functions on the quantum U_2 .

References

- [1] R. Baston, M. Eastwood. The Penrose Transform. Its Interaction with Representation Theory. Clarendon Press, Oxford, 1989.

⁵More precisely, the work by L. Turowska [20] presents explicit formulae that describe the action of the operators $\alpha, \beta, \gamma, \delta$ in the Hilbert space $l^2(\mathbb{Z}_+)^{\otimes d/2}$, with d being the dimension of the corresponding symplectic leaf.

- [2] V. Dobrev, *Representations of quantum groups and (conditionally) invariant q -difference equations*, Banach Center Publications, Vol. 40, Institute of Mathematics, Warszawa 1997, 203 – 222.
- [3] J. Faraut, A. Koranyi, *Function spaces and reproducing kernels on bounded symmetric domains*, J. Funct. Anal., **88** (1990), 64 – 89.
- [4] J. C. Jantzen. *Lectures on Quantum Groups*, Amer. Math. Soc., Providence RI, 1996.
- [5] H. P. Jakobsen, *Unitarity of highest-weight modules for quantum groups*, Letters in Math. Phys., **41** (1997), 119 – 133.
- [6] H. P. Jakobsen, *q -Differential Operators*, math.QA/9907009.
- [7] A. Kamita, *Quantum b -functions of prehomogeneous vector spaces of commutative parabolic type*, math.QA/0006209.
- [8] B. Kostant, *On the existence and irreducibility of certain series of representations*, Lie groups and Their Representations, I. M. Gelfand (ed), Halsted, New York, 1975, 231 – 329.
- [9] H. T. Koelink, *On $*$ -representations of the Hopf $*$ -algebra associated with the quantum group $U_q(n)$* , Compositio Math., **77** (1991), 199 – 231.
- [10] A. Klimyk, K. Schmüdgen. *Quantum Groups and their Representations*, Springer, Berlin-Heidelberg-New York, 1997.
- [11] G. Letzter, *Subalgebras which appear in quantum Iwasawa decomposition*, Canad. J. Math., **49** (1997), No 6, 1206 – 1223.
- [12] S. C. Milne, *A triple product identity for Schur functions*, J. Math. Anal. Appl., **160** (1991), No 2, 446 – 458.
- [13] M. Rosso, *Représentations des groupes quantiques*, Séminaire Bourbaki, exp. no 744, Astérisque, **201-203**, Société Mathématique de France, Paris, 1992, 443 – 483.
- [14] W. Schmid, *Construction and classification of irreducible Harish-Chandra modules*, in: Harmonic Analysis on Reductive Groups, W. Barker and P. Sally (eds), Birkhäuser, Boston MA, 1991, 235 – 275.
- [15] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *On function theory in quantum disc: integral representations*, math.QA/9808015.
- [16] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035.
- [17] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum matrix ball: The Bergman kernel*, math.QA/9909036.
- [18] L. Vaksman, *Quantum matrix ball: The Cauchy-Szegö kernel and the Shilov boundary*, Mathematical Physics, Analysis, and Geometry **8** (2001), No 4, 366 – 384; math.QA/0101179.
- [19] S. Sinel'shchikov and L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, q-alg/9703005; Math. Phys., Anal., and Geom., **1** (1998), 75 – 100.
- [20] L. Turowska, *On Representations of a q -Analogue of the $*$ -Algebra $Pol(Mat_2)$* , J. Phys. A **34** (2001), 2063 – 2070, math.OA/0005243.

Notes of the Editor

The results of L. Turowska mentioned in this work were applied recently in [1].

A quantum analogue of the Shilov boundary in a more general context was produced in [2].

The correspondence of the notion of Shilov boundary used in this work and its well known counterpart by W. Arveson is discussed in [3].

References

- [1] D. Proskurin, L. Turowska, *On the $*$ -algebra associated with $\text{Pol}(\text{Mat}_{2,2})_q$* , Methods of Functional Analysis and Topology, **7** (2001), No 1, 88 – 92.
- [2] L. Vaksman, *Quantum matrix ball: the Cauchy-Szegö kernel and the Shilov boundary*, Matematicheskaya Fizika. Analiz. Geometriya, **8** (2001), No 4, 366 – 384; math.QA/0101179.
- [3] L. Vaksman, *Maximum principle for holomorphic functions in quantum ball*, submitted to Matematicheskaya Fizika. Analiz. Geometriya (in Russian).

PART III
QUANTUM HARISH-CHANDRA MODULES
ASSOCIATED TO q -CARTAN DOMAINS

NON-COMPACT QUANTUM GROUPS AND HARISH-CHANDRA MODULES

D.Shklyarov[†]

S.Sinel'shchikov[†]

A.Stolin[‡]

L.Vaksman[†]

[†]Institute for Low temperature Physics & Engineering
47 Lenin Ave., 61103 Kharkov, Ukraine

[‡]Chalmers Tekniska Högskola, Matematik
412 96, Göteborg, Sweden

Abstract. An important problem of the quantum group theory is to construct and classify the Harish-Chandra modules; it is discussed in this work. The way of producing the principal non-degenerate series representations of the quantum $SU_{n,n}$ is sketched. A q-analogue for the Penrose transform is described.

A general theory of non-compact quantum groups which could include, for instance, quantum $SU_{2,2}$, does not exist. However, during the recent years, a number of problems on non-commutative geometry and harmonic analysis on homogeneous spaces of such 'groups' was solved. In these researches, the absent notion of non-compact quantum group was replaced by Harish-Chandra modules over quantum universal enveloping algebra $U_q\mathfrak{g}$. This work approaches an important and still open problem in the theory of quantum groups, the problem of constructing and classifying quantum Harish-Chandra modules. A construction of the principal non-degenerate series of quantum Harish-Chandra modules is described in the special case of the quantum $SU_{2,2}$. A q-analogue of the Penrose transform is investigated.

The last named author is grateful to V. Akulov for numerous discussions of geometric aspects of the quantum group theory.

Everywhere in the sequel $\mathfrak{g}$ stands for a simple complex Lie algebra and $\{\alpha_1, \alpha_2, \dots, \alpha_l\}$ for its system of simple roots with the standard ordering. The field $\mathbb{C}(q)$ of rational functions of the deformation parameter q normally works as a ground field (when solving the problems of harmonic analysis, it is more convenient to assume $q \in (0, 1)$ and to set $\mathbb{C}$ as a ground field).

A background in quantum universal enveloping algebras was made up by V. Drinfeld and M. Jimbo in mid-80-ies. The principal results of this theory at its early years are expounded in the review [9] and in the lectures [5]. We inherit the notation of these texts; in particular, we use the standard generators $\{E_j, F_j, K_j^{\pm 1}\}_{j=1,2,\dots,l}$ of the Hopf algebra $U_q\mathfrak{g}$ and relatively prime integers d_j , $j = 1, 2, \dots, l$, which symmetrize the Cartan matrix of $\mathfrak{g}$ (note that $d_j = 1$, $j = 1, 2, \dots, l$, in the case $\mathfrak{g} = \mathfrak{sl}_{l+1}$). We restrict ourselves to

This research was supported in part by Award No UM1-2091 of the US Civilian Research & Development Foundation

This lecture has been delivered at the Conference 'Supersymmetry and Quantum Field Theory'; published in Nucl. Physics B (Proc. Suppl.) 102& 103 (2001), 334 – 337.

considering $\mathbb{Z}^l$ -weight $U_q\mathfrak{g}$ -modules V i.e. those admitting a decomposition into a sum of weight subspaces

$$V = \bigoplus_{\mu} V_{\mu}, \quad \mu = (\mu_1, \mu_2, \dots, \mu_l) \in \mathbb{Z}^l, \quad V_{\mu} = \{v \in V \mid K_j^{\pm 1} v = q^{\pm d_j \mu_j} v, j = 1, 2, \dots, l\}.$$

Recall that some of the simple roots $\alpha_1, \alpha_2, \dots, \alpha_l$ determine Hermitian symmetric spaces of non-compact type [4]. (The coefficients of such simple roots in an expansion of the highest root of $\mathfrak{g}$ is 1.) For example, in the case $\mathfrak{g} = \mathfrak{sl}_{l+1}$ all simple roots possess this property. Choose one such root α_{j_0} and introduce the notation $U_q\mathfrak{k}$ for the Hopf subalgebra generated by $K_{j_0}^{\pm 1}, E_j, F_j, K_j^{\pm 1}, j \neq j_0$. Of course, every $U_q\mathfrak{g}$ -module V is also a $U_q\mathfrak{k}$ -module.

A finitely generated $\mathbb{Z}^l$ -weight $U_q\mathfrak{g}$ -module V is called quantum Harish-Chandra module if

1. $U_q\mathfrak{k}$ -module V is a sum of finite dimensional simple $U_q\mathfrak{k}$ -modules,
2. every finite dimensional simple $U_q\mathfrak{k}$ -module W occurs with finite multiplicity ($\dim \operatorname{Hom}_{U_q\mathfrak{k}}(W, V) < \infty$).

In the classical theory there are several methods of construction and classification of Harish-Chandra modules [10]. A similar problem for quantum Harish-Chandra modules is still open. In our opinion, it is among the most important of the quantum group theory.

To describe obstacles that appear in solving these problems, consider the Hermitian symmetric space $SU_{2,2}/S(U_2 \times U_2)$. It is determined by the Lie algebra $\mathfrak{g} = \mathfrak{sl}_4$ and the specified simple root $\alpha_{j_0} = \alpha_2$ of this Lie algebra.

Our primary desire is to construct a q-analogue for the principal non-degenerate series of Harish-Chandra modules. This interest is partially inspired by Casselman's theorem [1] which claims that every classical simple Harish-Chandra module admits an embedding into a principal non-degenerate series module. A well known method of producing this series is just the induction from a parabolic subalgebra. Regretfully, this subalgebra has no q-analogue (this obstacle does not appear if one substitutes the subalgebra $U\mathfrak{k}$, thus substituting a subject of research, cf. [7]). Fortunately, a quantization is available for another less known method of producing the principal non-degenerate series. We describe this method in a simple special case of quantum $SU_{2,2}$.

Let $G = SL_4(\mathbb{C})$, $K = S(GL_2 \times GL_2)$, $B \subset G$ be the standard Borel subgroup of upper triangular matrices, and $X = G/B$ the variety of complete flags. It is known that there exists an open K -orbit in X . It is well known that this orbit is an affine algebraic variety [10]. The regular functions on this orbit constitute a Harish-Chandra module of the principal non-degenerate series. The regular differential forms of the highest degree form another module of this series. The general case is *essentially* approached by considering a generic homogeneous algebraic line bundle on X and subsequent restricting it to the open K -orbit. The above construction procedure can be transferred to the quantum case and leads to the principal non-degenerate series of quantum Harish-Chandra modules.

In the classical representation theory, the above interplay between K -orbits on the variety and the theory of Harish-Chandra modules constitutes a generic phenomenon. In the theory of Beilinson-Bernstein, simple Harish-Chandra modules are derived from the so called standard Harish-Chandra modules. Furthermore, every series of standard Harish-Chandra modules is associated to a K -orbit in $X = G/B$. However, given an orbit Q of a codimension $s > 0$, one should consider the local cohomology $H_Q^s(X, \mathcal{F})$ instead of functions on Q , with $\mathcal{F}$ being a sheaf of sections of a homogeneous algebraic

line bundle (here our description of a standard module is somewhat naive but hopefully more plausible; the precise construction is expounded in [10]).

Our conjecture is that the standard quantum Harish-Chandra modules can be produced via some q-analogue of local cohomology $H_Q^s(X, \mathcal{F})$. An immediate obstacle that appears this way is in a lack of critical background in non-commutative algebraic geometry.

Probably the case of a closed K -orbit $Q \subset X$ is the simplest and most important one. Note that closed K -orbits are related to discrete series of Harish-Chandra modules, which are of an essential independent interest [12].

To conclude, consider a very simple example of a closed K -orbit in the space of *incomplete* flags which leads to the well known 'ladder representation' of the quantum $SU_{2,2}$ and a q-analogue of the Penrose transform [2].

We start with a purely algebraic description of the corresponding simple quantum Harish-Chandra module, to be succeeded with its two geometric realizations. Quantum Penrose transform intertwines these geometric realizations and is given by an explicit integral formula.

Consider the generalized Verma module M over $U_q \mathfrak{sl}_4$ given by its single generator v and the relations

$$\begin{aligned} F_j v &= E_j v = (K_j^{\pm 1} - 1)v = 0, & j &= 1, 3, \\ F_2 v &= 0, & K_2^{\pm 1} v &= q^{\pm 1} v. \end{aligned}$$

Consider its largest $U_q \mathfrak{sl}_4$ -submodule $J \subsetneq M$ and the associated quotient module $L = M/J$. Thus we get a simple quantum Harish-Chandra module which is a q-analogue of the $U_q \mathfrak{sl}_4$ -module related to the ladder representation of $SU_{2,2}$.

The first geometric realization is related to the K -action in the variety $\mathbb{CP}^3$ of lines in $\mathbb{C}^4$. We use the coordinate system (u_1, u_2, u_3, u_4) in $\mathbb{C}^4$. Let $Q \subset \mathbb{CP}^3$ be the subvariety of lines which are inside the plane $u_3 = u_4 = 0$. Then, as it was demonstrated in [2], the local cohomology $H_Q^2(\mathbb{CP}^3, \mathcal{O}(-2))$ can be described in terms of Laurent polynomials with complex coefficients

$$\begin{aligned} H_{[\mathbb{L}]}^2(\mathbb{CP}^3, \mathcal{O}(-2)) &\simeq \left\{ \sum_{j_1 j_2 j_3 j_4} c_{j_1 j_2 j_3 j_4} u_1^{j_1} u_2^{j_2} u_3^{j_3} u_4^{j_4} \right\} \\ &\quad \left. j_1 \geq 0 \quad \& \quad j_2 \geq 0 \quad \& \quad j_3 \leq -1 \quad \& \quad j_4 \leq -1 \quad \& \quad \sum_{j_i} = -2 \right\}. \end{aligned}$$

This space of Laurent polynomials is a $U \mathfrak{sl}_4$ -module since it is a quotient of the $U \mathfrak{sl}_4$ -module

$$\left\{ \sum_{j_1 j_2 j_3 j_4} c_{j_1 j_2 j_3 j_4} u_1^{j_1} u_2^{j_2} u_3^{j_3} u_4^{j_4} \mid j_1 \geq 0, \quad j_2 \geq 0 \right\}.$$

A similar geometric realization of the quantum Harish-Chandra module L is accessible via replacing the classical vector space $\mathbb{C}^4$ with the quantum one, which means just replacement of the commutation relations $u_j u_i = u_i u_j$ with $u_i u_j = q u_j u_i$, $1 \leq i < j \leq 4$.

The second geometric realization is in considering the vector space of polynomial solutions of the 'wave equation' in the space $\text{Mat}_{2,2}$ of 2×2 matrices:

$$\left\{ \psi \left(\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \right) \in \mathbb{C}[\text{Mat}_{2,2}] \mid \frac{\partial^2 \psi}{\partial \alpha \partial \delta} - \frac{\partial^2 \psi}{\partial \beta \partial \gamma} = 0 \right\}.$$

The structure of a $U\mathfrak{sl}_4$ -module in the space $\mathbb{C}[\text{Mat}_{2,2}]$ of polynomials on the space of matrices is introduced via embedding into the space of rational functions on the Grassmann variety of two-dimensional subspaces in $\mathbb{C}^4$:

$$\psi \left(\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \right) \mapsto \psi \left(\begin{pmatrix} t_{\{3,4\}}^{-1} t_{\{1,3\}} & t_{\{3,4\}}^{-1} t_{\{2,3\}} \\ t_{\{3,4\}}^{-1} t_{\{1,4\}} & t_{\{3,4\}}^{-1} t_{\{2,4\}} \end{pmatrix} \right) t_{\{3,4\}}^{-1}. \quad (0.1)$$

Here $t_{\{i,j\}} = t_{1i}t_{2j} - t_{1j}t_{2i}$, $i < j$, and t_{ij} are the generators of the algebra $\mathbb{C}[\text{Mat}_{2,4}]$ of polynomials on 2×4 matrices. This realization can be transferred onto the quantum case, with the ordinary wave equation being replaced by its q -analogue

$$\frac{\partial^2 \psi}{\partial \alpha \partial \delta} - q \frac{\partial^2 \psi}{\partial \beta \partial \gamma} = 0,$$

and ordinary matrices by quantum matrices (cf. [3]). It is worthwhile to note that we use $U_q\mathfrak{sl}_4$ -invariant differential calculus on the quantum space of 2×2 matrices.

The isomorphism of the two geometric realizations of the ladder representation is unique up to a constant multiple. In the case $q = 1$ it is given by the Penrose transform. It can be defined explicitly by the following integral formula:

$$f(u_1, u_2, u_3, u_4) \mapsto \int f \left(\sum_{i=1}^2 \zeta_i t_{i1}, \sum_{i=1}^2 \zeta_i t_{i2}, \sum_{i=1}^2 \zeta_i t_{i3}, \sum_{i=1}^2 \zeta_i t_{i4} \right) d\nu(\zeta),$$

where it is implicit that

$$\begin{aligned} (\zeta_1 t_{13} + \zeta_2 t_{23})^{-1} &= \frac{1}{\zeta_2 t_{23}} \cdot \sum_{k=0}^{\infty} (-1)^k \left(\frac{\zeta_1 t_{13}}{\zeta_2 t_{23}} \right)^k, \\ (\zeta_1 t_{14} + \zeta_2 t_{24})^{-1} &= \sum_{k=0}^{\infty} (-1)^k \left(\frac{\zeta_2 t_{24}}{\zeta_1 t_{14}} \right)^k \cdot \frac{1}{\zeta_1 t_{14}}, \end{aligned}$$

and the $U\mathfrak{sl}_2$ -invariant integral is given by

$$\int \left(\sum c_{i_1 i_2} \zeta_1^{i_1} \zeta_2^{i_2} \right) d\nu \stackrel{\text{def}}{=} c_{-1, -1}$$

(a passage from the Plücker coordinates $t_{\{i,j\}}$ to polynomials on $\text{Mat}_{2,2}$ is described by (0.1)). For example, in the case of a lowest weight vector $\frac{1}{u_3 u_4}$ we have

$$\frac{1}{u_3 u_4} \mapsto \frac{1}{t_{13} t_{24} - t_{14} t_{23}} \mapsto 1.$$

To pass from the classical case to the quantum one it suffices to replace the ordinary space $\mathbb{C}^2$ with its quantum analogue: $\zeta_1 \zeta_2 = q \zeta_2 \zeta_1$, the ordinary product in $\sum_i \zeta_i t_{ij}$ with the tensor product and to order multiples in the above formulae in a proper way.

We thus get the quantum Penrose transform, which is an isomorphism of the two geometric realizations of the quantum Harish-Chandra module L .

It is well known [6] that the quantum Harish-Chandra module L is unitarizable. The second geometric realization of this module allows one to find the corresponding scalar

product as in [8] (using an analytic continuation of the scalar product involved into the definition of the holomorphic discrete series [11]).

The precise formulations and complete proofs of the results announced in this work will be placed to the Eprint Archives (<http://www.arXiv.org/find/math>).

References

- [1] A. Beilinson, J. Bernstein, *A generalization of Casselman's submodule theorem*, in: Representation Theory of Reductive Groups, P. Trombi (ed), Birkhäuser, Boston MA, 1983, 35 – 52.
- [2] R. J. Baston, M. G. Eastwood. *The Penrose Transform. Its Interaction with Representation Theory*, Clarendon Press, Oxford, 1989.
- [3] V. Dobrev, *Representations of quantum groups and (conditionally) invariant q -difference equations*, Banach Center Publications, Vol. 40, Institute of Mathematics, Warszawa 1997, 203 – 222.
- [4] S. Helgason. *Differential Geometry and Symmetric Spaces*, Academic Press, New York–London, 1962.
- [5] J. C. Jantzen. *Lectures on Quantum Groups*, Amer. Math. Soc., Providence RI, 1996.
- [6] H. P. Jakobsen, *Unitarity of highest-weight modules for quantum groups*, Letters in Math. Phys., **41** (1997), 119 – 133.
- [7] G. Letzter, *Symmetric pairs for quantized enveloping algebras*, Journal of Algebra, **220** (1999), 729 – 767.
- [8] B. Ørsted, *Composition series for analytic continuation of holomorphic discrete series representations of $SU(n, n)$* , Trans. Amer. Math. Soc., **260** (1980), 563 – 573.
- [9] M. Rosso, *Représentations des groupes quantiques*, Séminaire Bourbaki, exp. no 744, Astérisque, **201-203**, Société Mathématique de France, Paris, 1992, 443 – 483.
- [10] W. Schmid, *Construction and classification of irreducible Harish-Chandra modules*, in: Harmonic Analysis on Reductive Groups, W. Baker and P. Sally (eds), Birkhäuser, Boston MA, 1991, 235 – 275.
- [11] D. Shklyarov, S. Sinel'shchikov, L. Vaksman, *Quantum matrix ball: the Bergman kernel*, math.QA/9909036, 1999, 28 p.p.
- [12] G. J. Zuckerman. *Geometric methods in representation theory*, in: Representation theory of reductive groups., P. Trombi (ed), Birkhäuser, Boston MA, 1983, 283 – 290.

Notes of the Editor

An appreciable break-through in the theory of non-compact quantum groups was advanced in the recent work [1].

References

- [1] E. Koelink, J. Kustermans, *A locally compact quantum group analogue of the normalizer of $SU_{1,1}$ in $SL(2, \mathbb{C})$* , math.QA/0105117.

ON A q -ANALOGUE OF THE PENROSE TRANSFORM

D.Shklyarov[†]

S.Sinel'shchikov[†]

A.Stolin[‡]

L.Vaksman[†]

[†]Institute for Low temperature Physics & Engineering
47 Lenin Ave., 61103 Kharkiv, Ukraine

[‡]Chalmers Tekniska Högskola, Matematik
412 96, Göteborg, Sweden

e-mail: sinelshchikov@ilt.kharkov.ua, vaksman@ilt.kharkov.ua, astolin@math.chalmers.se

1 Introduction

In the framework of the theory of quantum groups and their homogeneous spaces we consider two geometric realizations for the quantum ladder representation, together with an intertwining linear transformation — the quantum Penrose transform.

In section 2 we supply a preliminary material on the classical Penrose transform and prove (2.1). The q -analogue of (2.1) is to be used in section 3 to produce a quantum Penrose transform.

Our results hint that a great deal of constructions specific for the theory of quasi-coherent sheaves admit non-commutative analogues. This research is motivated by a possibility to use the results of non-commutative algebraic geometry for producing and studying Harish-Chandra modules over quantum universal enveloping algebras.

There is a plenty of literature on the Penrose transform, quantum groups, and non-commutative algebraic geometry. We restrict ourselves to mentioning the monographs [2, 6, 3], papers [1, 10], and the preprint [9].

Note that a noncommutative analogues for the Penrose transform and covariant differential operators are also considered in the preprints [8, 12] and in the papers [5, 4, 7] respectively in a completely different context.

2 The classical case

To recall the definition of the Penrose transform, we restrict ourselves to a simplest substantial example. In this special case, the Penrose transform intertwines the cohomology of the sheaf $\mathcal{O}(-2)$ on

$$U' = \{(u_1 : u_2 : u_3 : u_4) \in \mathbb{CP}^3 \mid u_3 \neq 0 \text{ or } u_4 \neq 0\}$$

and sections of the sheaf $\mathcal{O}(-1)$ on some open affine submanifold of the Grassmann manifold $\text{Gr}_2(\mathbb{C}^4) \hookrightarrow \mathbb{CP}^5$. Instead of the Grassmann manifold, we prefer to consider

Ukrainian Journal of Physics, **47** (2002), No 3, 288 – 292

This research was partially supported by Award No UM1-2091 of the Civilian Research & Development Foundation

the Stifel manifold of ordered linear independent pairs of vectors in $\mathbb{C}^4$. In this context, the GL_2 -covariant sections on the Stifel manifold work as sections of the sheaf $\mathcal{O}(-1)$ on $\text{Gr}_2(\mathbb{C}^4)$.

Associate to each matrix $\mathbf{t} = (t_{ij})_{i=1,2;j=1,2,3,4} \in \text{Mat}_{2,4}$ the pairs of vectors in $\mathbb{C}^4$:

$$(t_{11}, t_{12}, t_{13}, t_{14}), \quad (t_{21}, t_{22}, t_{23}, t_{24}).$$

Consider $U'' = \{\mathbf{t} \in \text{Mat}_{2,4} \mid t_{13}t_{24} - t_{14}t_{23} \neq 0\}$. Every point $\mathbf{u} = (u_1 : u_2 : u_3 : u_4) \in U'$ determines a one-dimensional subspace $L_{\mathbf{u}} \subset \mathbb{C}^4$, and every point $\mathbf{t} \in U''$ determines a two-dimensional subspace $L_{\mathbf{t}}$ generated by the vectors of the corresponding pair. Let $U = \{(\mathbf{u}, \mathbf{t}) \in U' \times U'' \mid L_{\mathbf{u}} \subset L_{\mathbf{t}}\}$. We thus get a 'double fibration' $U' \xleftarrow{\eta} U \xrightarrow{\tau} U''$, which leads to the Penrose transform. It should be noted that every line $L_{\mathbf{t}}$ is of the form

$$L = \mathbb{C}(\zeta_1, \zeta_2) \begin{pmatrix} t_{11} & t_{12} & t_{13} & t_{14} \\ t_{21} & t_{22} & t_{23} & t_{24} \end{pmatrix}, \quad (\zeta_1, \zeta_2) \in \mathbb{C}^2.$$

Hence the above double fibration is isomorphic to the double fibration

$$U' \xleftarrow{\pi} \mathbb{CP}^1 \times U'' \xrightarrow{\text{pr}_2} U'',$$

with

$$\pi : \left((\zeta_1 : \zeta_2), \begin{pmatrix} t_{11} & t_{12} & t_{13} & t_{14} \\ t_{21} & t_{22} & t_{23} & t_{24} \end{pmatrix} \right) \mapsto ((\zeta_1 t_{11} + \zeta_2 t_{21}) : (\zeta_1 t_{12} + \zeta_2 t_{22}) : (\zeta_1 t_{13} + \zeta_2 t_{23}) : (\zeta_1 t_{14} + \zeta_2 t_{24})).$$

We thus get a coordinate description for the double fibration in question; this coordinate description is going to be implicit in all subsequent computations. Let us look at the cohomologies.

Consider an open affine cover $U' = U_1 \cup U_2$,

$$U_1 = \{(u_1 : u_2 : u_3 : u_4) \in U' \mid u_3 \neq 0\}, \\ U_2 = \{(u_1 : u_2 : u_3 : u_4) \in U' \mid u_4 \neq 0\},$$

and compute the Čech cohomology $\check{H}^1(U', \mathcal{O}(-2))$. Let $\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4^{\pm 1}]$ be the Laurent polynomials in indeterminates u_3, u_4 , with coefficients from $\mathbb{C}[u_1, u_2]$. Introduce in a similar way $\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4]$, $\mathbb{C}[u_1, u_2, u_3, u_4^{\pm 1}]$; of course, these appear to be $U\mathfrak{sl}_4$ -modules.

It follows from the definition of the Čech complex that there exists a natural isomorphism of $U\mathfrak{sl}_4$ -modules:

$$\check{H}^1(U', \mathcal{O}(-2)) = \left\{ f \in \mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4^{\pm 1}] / (\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4] + \mathbb{C}[u_1, u_2, u_3, u_4^{\pm 1}]) \mid \deg f = -2 \right\}.$$

Hence the Laurent polynomials

$$\frac{u_1^{j_1} u_2^{j_2}}{u_3^{j_3} u_4^{j_4}}, \quad j_3 \geq 1 \quad \& \quad j_4 \geq 1 \quad \& \quad j_3 + j_4 = j_1 + j_2 + 2,$$

form a basis of the vector space $H^1(U', \mathcal{O}(-2))$.

Consider the trivial bundle over U'' with fiber $H^1(\mathbb{CP}^1, \mathcal{O}(-2))$. It is known that $H^1(\mathbb{CP}^1, \mathcal{O}(-2)) \simeq \mathbb{C}$, and the isomorphism is available via choosing an open affine cover $\mathbb{CP}^1 = \{(\zeta_1 : \zeta_2) \mid \zeta_1 \neq 0\} \cup \{(\zeta_1 : \zeta_2) \mid \zeta_2 \neq 0\}$. Specifically, $\sum_{j+k=-2} c_{jk} \zeta_1^j \zeta_2^k \mapsto c_{-1,-1}$.

In a different notation $f \mapsto \text{CT}(\zeta_1 \zeta_2 f)$, with $\text{CT} : \sum_{j,k} c_{jk} \zeta_1^j \zeta_2^k \mapsto c_{00}$ (the constant term of a series). Now $\mathcal{P}f$ is defined as a higher direct image of the cohomology class $\eta^* f$: $\mathcal{P}f = \tau_*^1 \eta^* f$. The linear map τ_*^1 is called the integration along the fibers of τ . We restrict ourselves to computing this 'integral' inside the infinitesimal neighborhood of $\mathbf{t}_0 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$ by using formal series in $t_{11}, t_{12}, t_{13}, t_{14}^{-1}, t_{21}, t_{23}^{-1}, t_{24}$ with coefficients from $\mathbb{C}[\zeta_1^{\pm 1}, \zeta_2^{\pm 1}]$. Of course, $\eta^* : f(\mathbf{u}) \mapsto f(\boldsymbol{\zeta} \mathbf{t})$. So, in the coordinate description

$$\mathcal{P} : f(\mathbf{u}) \mapsto \text{CT}_{\boldsymbol{\zeta}}(\zeta_1 \zeta_2 f(\boldsymbol{\zeta} \mathbf{t})), \quad f \in u_3^{-1} u_4^{-1} \mathbb{C}[u_1, u_2, u_3^{-1}, u_4^{-1}], \quad (2.1)$$

with $\text{CT}_{\boldsymbol{\zeta}}$ being the constant term in the indeterminate $\boldsymbol{\zeta}$.

EXAMPLE. Compute $\mathcal{P}(1/(u_3 u_4))$. One has:

$$\begin{aligned} \frac{1}{\zeta_1 t_{13} + \zeta_2 t_{23}} &= \frac{1}{\zeta_2 t_{23}} \sum_{i=0}^{\infty} (-1)^i \left(\frac{\zeta_1 t_{13}}{\zeta_2 t_{23}} \right)^i, \\ \frac{1}{\zeta_1 t_{14} + \zeta_2 t_{24}} &= \frac{1}{\zeta_1 t_{14}} \sum_{j=0}^{\infty} (-1)^j \left(\frac{\zeta_2 t_{24}}{\zeta_1 t_{14}} \right)^j. \end{aligned}$$

Hence,

$$\begin{aligned} \mathcal{P} \left(\frac{1}{u_3 u_4} \right) &= \text{CT}_{\boldsymbol{\zeta}} \left(\frac{1}{t_{23} t_{14}} \sum_{i,j=0}^{\infty} (-1)^{i+j} \left(\frac{\zeta_1 t_{13}}{\zeta_2 t_{23}} \right)^i \left(\frac{\zeta_2 t_{24}}{\zeta_1 t_{14}} \right)^j \right) = \\ &= \frac{1}{t_{23} t_{14}} \sum_{k=0}^{\infty} \left(\frac{t_{13} t_{24}}{t_{23} t_{14}} \right)^k = \frac{1}{t_{23} t_{14}} \cdot \frac{1}{1 - \frac{t_{13} t_{24}}{t_{23} t_{14}}} = -\frac{1}{t_{13} t_{24} - t_{14} t_{23}}. \end{aligned}$$

REMARK. It is known that the Penrose transform is an isomorphism between the two realizations for the 'ladder' representation of $\mathfrak{sl}_4$: the representation in $H^1(U', \mathcal{O}(-2))$ and the representation in

$$\{\psi(z_1^1, z_2^1, z_1^2, z_2^2)(t_{13} t_{24} - t_{14} t_{23})^{-1} \in H^0(U'', \mathcal{O}(-1)) \mid \square \psi = 0\},$$

where $\square \psi \stackrel{\text{def}}{=} \frac{\partial^2 \psi}{\partial z_1^1 \partial z_2^2} - \frac{\partial^2 \psi}{\partial z_2^1 \partial z_1^2}$, and

$$\begin{aligned} z_1^1 &= \frac{t_{11} t_{23} - t_{13} t_{21}}{t_{13} t_{24} - t_{14} t_{23}}, & z_2^1 &= \frac{t_{12} t_{23} - t_{13} t_{22}}{t_{13} t_{24} - t_{14} t_{23}} \\ z_1^2 &= \frac{t_{11} t_{24} - t_{14} t_{21}}{t_{13} t_{24} - t_{14} t_{23}}, & z_2^2 &= \frac{t_{12} t_{24} - t_{14} t_{22}}{t_{13} t_{24} - t_{14} t_{23}}. \end{aligned}$$

The vectors $1/(u_3 u_4)$ and $1/(t_{13} t_{24} - t_{14} t_{23})$ are lowest weight vectors for the above representations of $\mathfrak{sl}_4$. Of course, $z_1^1, z_2^1, z_1^2, z_2^2$ can be considered as the standard coordinates on the big cell $t_{13} t_{24} - t_{14} t_{23} \neq 0$ of the Grassmanian $\text{Gr}_2(\mathbb{C}^4)$.

3 The quantum case

In the previous section we produced the formula (2.1) which can be treated as a definition of the Penrose transform in the classical case. Now our intention is to produce a q -analogue of (2.1). The principal difference from the constructions of section 2 is in replacement of the functors η^* , τ_*^1 of the sheaf theory with the corresponding morphisms of $U_q\mathfrak{sl}_4$ -modules. (Here $U_q\mathfrak{sl}_4$ is a quantum universal enveloping algebra. It is a Hopf algebra over the ground field $\mathbb{C}(q)$ and is determined by the generators $\{E_i, F_i, K_i^{\pm 1}\}_{i=1,2,3}$ and the well known Drinfeld-Jimbo relations [6].)

The quantum projective space $\mathbb{CP}_{\text{quant}}^3$ is defined in terms of a $\mathbb{Z}_+$ -graded algebra $\mathbb{C}[u_1, u_2, u_3, u_4]_q$ whose generators u_1, u_2, u_3, u_4 are subject to the commutation relations

$$u_i u_j = q u_j u_i, \quad i < j.$$

Just as in the classical case, $\deg u_k = 1$, $k = 1, 2, 3, 4$. The localization $\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4^{\pm 1}]_q$ of $\mathbb{C}[u_1, u_2, u_3, u_4]_q$ with respect to the multiplicative system $(u_3 u_4)^{\mathbb{N}}$ is equipped in a standard way with a structure of $U_q\mathfrak{sl}_4$ -module algebra. The subalgebras $\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4]_q$, $\mathbb{C}[u_1, u_2, u_3, u_4^{\pm 1}]_q$ constitute $U_q\mathfrak{sl}_4$ -submodules of the $U_q\mathfrak{sl}_4$ -module $\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4^{\pm 1}]_q$. Thus we come to

$$V' = \mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4^{\pm 1}]_q / (\mathbb{C}[u_1, u_2, u_3^{\pm 1}, u_4]_q + \mathbb{C}[u_1, u_2, u_3, u_4^{\pm 1}]_q) \quad (3.1)$$

as a q -analogue of the $U\mathfrak{sl}_4$ -module $H^1(U', \mathcal{O}(-2))$.

We have produced a q -analogue for the first geometric realization of the 'ladder representation'. Turn to a construction of its second geometric realization.

The algebra $\mathbb{C}[\text{Mat}_{2,4}]_q$ of polynomials on the quantum matrix space is determined by its generators $\{t_{ij}\}_{i=1,2;j=1,2,3,4}$ and the well known commutation relations

$$\begin{aligned} t_{ik} t_{jk} &= q t_{jk} t_{ik}, & t_{ki} t_{kj} &= q t_{kj} t_{ki}, & i < j, \\ t_{ij} t_{kl} &= t_{kl} t_{ij}, & & & i < k \quad \& \quad j > l, \\ t_{ij} t_{kl} - t_{kl} t_{ij} &= (q - q^{-1}) t_{ik} t_{jl}, & & & i < k \quad \& \quad j < l. \end{aligned}$$

The element $t = t_{13} t_{24} - q t_{14} t_{23}$ quasi-commutes with all the generators t_{ij} , $i = 1, 2$, $j = 1, 2, 3, 4$. Let $\mathbb{C}[\text{Mat}_{2,4}]_{q,t}$ be a localization of $\mathbb{C}[\text{Mat}_{2,4}]_q$ with respect to the multiplicative system $t^{\mathbb{N}}$ and $U_q\mathfrak{sl}_2$ the quantum universal enveloping algebra (determined by the generators $E, F, K^{\pm 1}$ and the Drinfeld-Jimbo relations).

$\mathbb{C}[\text{Mat}_{2,4}]_{q,t}$ is equipped in a standard way with a structure of $U_q\mathfrak{sl}_2 \otimes U_q\mathfrak{sl}_4$ -module algebra. In particular, $\mathbb{C}[\text{Mat}_{2,4}]_{q,t}$ is a $U_q\mathfrak{sl}_4$ -module algebra.

Introduce the notation:

$$\begin{aligned} z_1^1 &= t^{-1}(t_{11} t_{23} - q t_{13} t_{21}), & z_2^1 &= t^{-1}(t_{12} t_{23} - q t_{13} t_{22}), \\ z_1^2 &= t^{-1}(t_{11} t_{24} - q t_{14} t_{21}), & z_2^2 &= t^{-1}(t_{12} t_{24} - q t_{14} t_{22}). \end{aligned}$$

It is well known and easily deducible that

$$\begin{aligned} z_k^i z_k^j &= q z_k^j z_k^i, & z_i^k z_j^k &= q z_j^k z_i^k, & i < j, \\ z_j^i z_l^k &= z_l^k z_j^i, & & & i < k \quad \& \quad j > l, \\ z_j^i z_l^k - z_l^k z_j^i &= (q - q^{-1}) z_k^i z_l^j, & & & i < k \quad \& \quad j < l. \end{aligned}$$

It follows that the subalgebra generated by $z_1^1, z_2^1, z_1^2, z_2^2$ is 'canonically' isomorphic to the algebra $\mathbb{C}[\text{Mat}_{2,2}]_q$ of 'polynomials on the quantum matrix space'. It is easy to demonstrate that $\mathbb{C}[\text{Mat}_{2,2}]_q t^{-1}$ is a $U_q \mathfrak{sl}_4$ -submodule of the $U_q \mathfrak{sl}_4$ -module $\mathbb{C}[\text{Mat}_{2,4}]_{q,t}$.

The simple submodule of the $U_q \mathfrak{sl}_4$ -module $\mathbb{C}[\text{Mat}_{2,2}]_q t^{-1}$ we are interested in is distinguished via a q -analogue $\square_q$ of the wave operator $\square$:

$$\square_q = \frac{\partial}{\partial z_1^1} \frac{\partial}{\partial z_2^2} - q \frac{\partial}{\partial z_2^1} \frac{\partial}{\partial z_1^2}.$$

Specifically, $V'' = \{\psi t^{-1} \mid \square_q \psi = 0, \psi \in \mathbb{C}[\text{Mat}_{2,2}]_q\}$. It is worthwhile to note that the operators $\frac{\partial}{\partial z_j^i}$ are defined in terms of a $U_q \mathfrak{sl}_4$ -invariant first order differential calculus in $\mathbb{C}[\text{Mat}_{2,2}]_q$:

$$df = \sum_{i,j} \frac{\partial f}{\partial z_j^i} dz_j^i.$$

In turn, this first order differential calculus is defined by the following well known 'commutation' relations:

$$\begin{cases} z_1^1 dz_1^1 = q^{-2} dz_1^1 \cdot z_1^1 \\ z_1^1 dz_2^1 = q^{-1} dz_2^1 \cdot z_1^1 \\ z_1^1 dz_1^2 = q^{-1} dz_1^2 \cdot z_1^1 \\ z_1^1 dz_2^2 = dz_2^2 \cdot z_1^1 \\ z_2^1 dz_1^1 = q^{-1} dz_1^1 \cdot z_2^1 + (q^{-2} - 1) dz_2^1 \cdot z_1^1 \\ z_2^1 dz_2^1 = q^{-2} dz_2^1 \cdot z_2^1 \\ z_2^1 dz_1^2 = dz_1^2 \cdot z_2^1 + (q^{-1} - q) dz_2^2 \cdot z_1^1 \\ z_2^1 dz_2^2 = q^{-1} dz_2^2 \cdot z_2^1 \\ z_1^2 dz_1^1 = q^{-1} dz_1^1 \cdot z_1^2 + (q^{-2} - 1) dz_1^2 \cdot z_1^1 \\ z_1^2 dz_2^1 = dz_2^1 \cdot z_1^2 + (q^{-1} - q) dz_2^2 \cdot z_1^1 \\ z_1^2 dz_1^2 = q^{-2} dz_1^2 \cdot z_1^2 \\ z_1^2 dz_2^2 = q^{-1} dz_2^2 \cdot z_1^2 \\ z_2^2 dz_1^1 = dz_1^1 \cdot z_2^2 + (q^{-1} - q) dz_2^1 \cdot z_1^2 + (q^{-1} - q) dz_1^2 \cdot z_2^1 + (q^{-1} - q)^2 dz_2^2 \cdot z_1^1 \\ z_2^2 dz_2^1 = q^{-1} dz_2^1 \cdot z_2^2 + (q^{-2} - 1) dz_2^2 \cdot z_2^1 \\ z_2^2 dz_1^2 = q^{-1} dz_1^2 \cdot z_2^2 + (q^{-2} - 1) dz_2^2 \cdot z_1^2 \\ z_2^2 dz_2^2 = q^{-2} dz_2^2 \cdot z_2^2 \end{cases}$$

We thus get the two $U_q \mathfrak{sl}_4$ -modules V', V'' ; our intention is to find an explicit form of the linear map which provides an isomorphism $\mathcal{P} : V' \rightarrow V''$.

We follow the ideas of classical constructions described in section 2 in considering the quantum projective space $\mathbb{CP}_{\text{quant}}^1$. More precisely, let us consider a $\mathbb{Z}_+$ -graded algebra $\mathbb{C}[\zeta_1, \zeta_2]_q$:

$$\zeta_1 \zeta_2 = q \zeta_2 \zeta_1, \quad \deg(\zeta_1) = \deg(\zeta_2) = 1,$$

together with its localization $\mathbb{C}[\zeta_1^{\pm 1}, \zeta_2^{\pm 1}]_q$ with respect to the multiplicative system $(\zeta_1 \zeta_2)^{\mathbb{N}}$. The algebra $\mathbb{C}[\zeta_1^{\pm 1}, \zeta_2^{\pm 1}]_q$ is equipped in a standard way with a structure of $U_q \mathfrak{sl}_2$ -module

algebra. The following homomorphism of algebras will work as the operator $f(\mathbf{u}) \mapsto f(\boldsymbol{\zeta} \mathbf{t})$:

$$\begin{aligned} \eta^* : \mathbb{C}[u_1, u_2, u_3, u_4]_q &\rightarrow \mathbb{C}[\zeta_1, \zeta_2]_q \otimes \mathbb{C}[\text{Mat}_{2,4}]_q, \\ \eta^* : u_j &\mapsto \zeta_1 \otimes t_{1j} + \zeta_2 \otimes t_{2j}, \quad j = 1, 2, 3, 4. \end{aligned}$$

To follow the constructions of section 2, we have to invert the elements $\zeta_1 \otimes t_{13} + \zeta_2 \otimes t_{23}$, $\zeta_1 \otimes t_{14} + \zeta_2 \otimes t_{24}$ in a suitable localization of $\mathbb{C}[\zeta_1^{\pm 1}, \zeta_2^{\pm 1}]_q \otimes \mathbb{C}[\text{Mat}_{2,4}]_{q,t}$. It is easy to verify that

$$\begin{aligned} (t_{14}t_{23})^2 \cdot \mathbb{C}[\text{Mat}_{2,4}]_{q,t} &\subset \mathbb{C}[\text{Mat}_{2,4}]_{q,t} \cdot (t_{14}t_{23}), \\ \mathbb{C}[\text{Mat}_{2,4}]_{q,t} \cdot (t_{14}t_{23})^2 &\subset (t_{14}t_{23}) \cdot \mathbb{C}[\text{Mat}_{2,4}]_{q,t}. \end{aligned}$$

Thus we have a well defined localization of $\mathbb{C}[\text{Mat}_{2,4}]_{q,t}$ with respect to the multiplicative system $(t_{14}t_{23})^{\mathbb{N}}$. In an appropriate completion of this algebra one has the following relations:

$$\begin{aligned} (\zeta_1 \otimes t_{13} + \zeta_2 \otimes t_{23})^{-1} &= (\zeta_2 \otimes t_{23})^{-1} \sum_{i=0}^{\infty} (-1)^i (\zeta_1 \zeta_2^{-1})^i \otimes (t_{13}t_{23}^{-1})^i, \\ (\zeta_1 \otimes t_{14} + \zeta_2 \otimes t_{24})^{-1} &= \left(\sum_{j=0}^{\infty} (-1)^j (\zeta_1^{-1} \zeta_2)^j \otimes (t_{14}^{-1}t_{24})^j \right) (\zeta_1 \otimes t_{14})^{-1}. \end{aligned}$$

We define the quantum Penrose transform by

$$\mathcal{P}_q f = (\text{CT} \otimes \text{id})(\zeta_1 \zeta_2 \otimes 1)(\eta^* f),$$

where, just as above, $\text{CT} : \sum_{ij} c_{ij} \zeta_1^i \zeta_2^j \mapsto c_{-1,-1}$, and f belongs to the linear span of the elements

$$u_1^{j_1} u_2^{j_2} u_3^{-j_3} u_4^{-j_4}, \quad j_3 \geq 1 \quad \& \quad j_4 \geq 1 \quad \& \quad j_1 + j_2 - j_3 - j_4 = -2. \quad (3.2)$$

Now (3.1) determines a $U_q \mathfrak{sl}_4$ -module structure in this linear span since the monomials (3.2) form a basis in the vector space V' .

4 Appendix

We sketch here the proof of the fact that $\mathcal{P}_q$ is an isomorphism of $U_q \mathfrak{sl}_4$ -modules $V' \xrightarrow{\sim} V''$.

It follows from the definition that $\mathcal{P}_q$ is a morphism of $U_q \mathfrak{sl}_4$ -modules. In view of the simplicity of V' and V'' , it suffices to prove that $\mathcal{P}_q$ takes the (lowest weight) vector $u_3^{-1} u_4^{-1} \in V'$ to the (lowest weight) vector $-(t_{13}t_{24} - qt_{14}t_{23})^{-1}$. We start with an auxiliary statement:

$$(1 - (t_{23}^{-1}t_{13})(t_{14}^{-1}t_{24}))^{-1} = \sum_{k=0}^{\infty} q^{-2k} (t_{23}^{-1}t_{13})^k (t_{14}^{-1}t_{24})^k. \quad (4.1)$$

It follows from the commutation relation

$$(t_{14}^{-1}t_{24})(t_{23}^{-1}t_{13}) = q^{-2}(t_{23}^{-1}t_{13})(t_{14}^{-1}t_{24}) + 1 - q^{-2}$$

and the relation (6.5) of [11].

An application of (4.1) allows one to prove that

$$\mathcal{P}_q(u_3^{-1}u_4^{-1}) = -(t_{13}t_{24} - qt_{14}t_{23})^{-1}.$$

In fact,

$$\begin{aligned} \mathcal{P}_q(u_3^{-1}u_4^{-1}) &= \text{CT} \otimes \text{id} \left(\zeta_2^{-1} \otimes t_{23}^{-1} \cdot \right. \\ &\quad \left. \left(\sum_{i=0}^{\infty} (-1)^i (\zeta_1 \zeta_2^{-1})^i \otimes q^{-i} (t_{23}^{-1} t_{13})^i \right) \left(\sum_{j=0}^{\infty} (-1)^j (\zeta_1 \zeta_2^{-1})^j \otimes (t_{14}^{-1} t_{24})^j \right) \zeta_1^{-1} \otimes t_{14}^{-1} \right). \end{aligned}$$

On the other hand,

$$\zeta_2^{-1} (\zeta_1 \zeta_2^{-1})^k (\zeta_1^{-1} \zeta_2)^k \zeta_1^{-1} = q^{-k} \zeta_2^{-1} \zeta_1^{-1} = q^{-k-1} \zeta_1^{-1} \zeta_2^{-1}.$$

Hence,

$$\begin{aligned} \mathcal{P}_q(u_3^{-1}u_4^{-1}) &= q^{-1} t_{23}^{-1} \left(\sum_{k=0}^{\infty} q^{-2k} (t_{14}^{-1} t_{24})^k (t_{23}^{-1} t_{13})^k \right) t_{14}^{-1} = \\ &= q^{-1} (t_{14} (1 - t_{23}^{-1} t_{13} t_{14}^{-1} t_{24}) t_{23})^{-1} = -(t_{13}t_{24} - qt_{14}t_{23})^{-1}, \end{aligned}$$

which completes the proof.

References

- [1] Artin M., Zhang J. J., *Noncommutative projective schemes*, Adv. in Math., **109** (1994), 228 – 287.
- [2] Baston R., Eastwood M. *The Penrose Transform. Its Interaction with Representation Theory*, Clarendon Press, Oxford, 1989.
- [3] Chari V., Pressley A. *A Guide to Quantum Groups*. Cambridge Univ. Press, 1995.
- [4] Dobrev V. K., *Representations of quantum groups and (conditional) invariant q -difference equations*, in: Quantum groups and Quantum Spaces, Banach Center Publ., vol 40, Inst. of Math., Polish Academy of Sciences, Warszawa, 1997, 203 – 222.
- [5] Jakobsen H. P., *Quantized Dirac operators*, math.QA/0010012.
- [6] Jantzen J. C. *Lectures on Quantum Groups*. – Amer. Math. Soc., Providence RI, 1996.
- [7] Kamita A., *Quantum b -functions of prehomogeneous vector spaces of commutative parabolic type*, math.QA/0006209.
- [8] Kapustin A., Kuznetsov A., Orlov D., *Noncommutative instantons and twistor transform*, hep-th/0002193.
- [9] Marastoni C., Tanisaki T., *Radon transforms for quasi-equivariant $\mathcal{D}$ -modules on generalized flag manifolds*, math.RT/9911095.
- [10] Rosenberg A. L., *Noncommutative schemes*, Compositio Mathematica, **112** (1998), 93 – 125.

- [11] Shklyarov D., Sinel'shchikov S., Vaksman L., *On function theory in quantum disc: a q -analogue of the Berezin transform*, math.QA/9809018.
- [12] Zupnik B. M., *Quantum deformations of the self-duality equation and conformal twistors*, q-alg/9509014.

SPHERICAL PRINCIPAL NON-DEGENERATE SERIES OF REPRESENTATIONS FOR THE QUANTUM GROUP $SU_{2,2}$

S. Sinel'shchikov[†] A. Stolin[‡] L. Vaksman[†]

[†]Institute for Low temperature Physics & Engineering
47 Lenin Ave., 61103 Kharkov, Ukraine

[‡]Chalmers Tekniska Högskola, Matematik
412 96, Göteborg, Sweden

1 Introduction

The first step in studying quantum bounded symmetric domains was done in [11]. Later on, an explicit form of the Plancherel measure was found in [12] for the simplest among the above domains, the quantum disc. It is still open to extend this result onto an arbitrary quantum bounded symmetric domain. Presumably, the initial step in this direction should be in producing and studying q -analogues of the principal non-degenerate series representations for automorphism groups of bounded symmetric domains. (It is certainly implicit in this setting that the quantum universal enveloping Drinfeld-Jimbo algebras work as quantum groups and the Harish-Chandra modules over those algebras work as representations of quantum groups [14].)

The present work deals with the simplest bounded symmetric domain of rank 2

$$\mathcal{D} = \{\mathbf{z} \in \text{Mat}_2(\mathbb{C}) \mid \mathbf{z}\mathbf{z}^* < I\},$$

its q -analogue (the quantum matrix ball), together with the associated Harish-Chandra modules. In section 2 we present a construction of the principal non-degenerate series of such modules. In section 3 we prove that, together with every Harish-Chandra module this series contains the dual Harish-Chandra module.

The third named author is grateful to A. Rosenberg and Ya. Soibelman for a discussion of the results expounded in section 2 of this paper.

2 A construction of the spherical principal non-degenerate series

From now on we assume q to be transcendental and $\mathbb{C}$ to be the ground field.

Consider the quantum universal enveloping algebra $U_q\mathfrak{sl}_4$. It is determined by its generators $\{E_j, F_j, K_j^{\pm 1}\}_{j=1,2,3}$ and the well known Drinfeld-Jimbo relations (see [5]). Recall that $U_q\mathfrak{sl}_4$ is a Hopf algebra. (The comultiplication $\Delta : U_q\mathfrak{sl}_4 \rightarrow U_q\mathfrak{sl}_4 \otimes U_q\mathfrak{sl}_4$ is defined

This research was supported in part by of the US Civilian Research & Development Foundation (Award No UM1-2091) and by Swedish Academy of Sciences (project No 11293562).

This lecture has been delivered at the 10th International Colloquium 'Quantum Groups and Integrable Systems' in Prague, June 2001. The text is published in Czechoslovak Journal of Physics **51** (2001), No 12, 1431 – 1440.

by $\Delta(K_j^{\pm 1}) = K_j^{\pm 1} \otimes K_j^{\pm 1}$, $\Delta(E_j) = E_j \otimes 1 + K_j \otimes E_j$, $\Delta(F_j) = F_j \otimes K_j^{-1} + 1 \otimes F_j$, $j = 1, 2, 3$, the counit $\varepsilon : U_q \mathfrak{sl}_4 \rightarrow \mathbb{C}$ by $\varepsilon(E_j) = \varepsilon(F_j) = \varepsilon(K_j^{\pm 1} - 1) = 0$, $j = 1, 2, 3$, and the antipode $S : U_q \mathfrak{sl}_4 \rightarrow U_q \mathfrak{sl}_4$ by $S(K_j^{\pm 1}) = K_j^{\mp 1}$, $S(E_j) = -K_j^{-1} E_j$, $S(F_j) = -F_j K_j$, $j = 1, 2, 3$. An ordinary universal enveloping algebra $U \mathfrak{sl}_4$ is derivable from $U_q \mathfrak{sl}_4$ via substituting $q = e^{-h/2}$, $K_j^{\pm 1} = e^{\mp h H_j/2}$ and a formal passage to the limit $h \rightarrow 0$.

Introduce the notation $U_q \mathfrak{g}$ for the Hopf algebra $U_q \mathfrak{sl}_4$ and $U_q \mathfrak{k}$ for its Hopf subalgebra generated by $K_2^{\pm 1}$, E_j , F_j , $K_j^{\pm 1}$, $j = 1, 3$. A $U_q \mathfrak{g}$ -module V is said to be $\mathbb{Z}^3$ -weight module if $V = \bigoplus_{\mu} V_{\mu}$, $\mu = (\mu_1, \mu_2, \mu_3) \in \mathbb{Z}^3$, $V_{\mu} = \{v \in V \mid K_j^{\pm 1} v = q^{\pm \mu_j} v, j = 1, 2, 3\}$.

In what follows we are going to consider only $U_q \mathfrak{g}$ -modules V of the above form, which allows one to introduce the linear operators H_1, H_2, H_3 , in V with $K_i^{\pm 1} = q^{\pm H_i}$, $i = 1, 2, 3$. Specifically, $H_i|_{V_{\mu}} = \mu_i I$, $i = 1, 2, 3$.

REMARK. For every simple Lie algebra and every lattice L which is between the lattices of roots and weights, a class of L -weight $U_q \mathfrak{g}$ -modules can be introduced in the same way.

A $\mathbb{Z}^3$ -weight $U_q \mathfrak{g}$ -module V is called a Harish-Chandra module if

1. $U_q \mathfrak{k}$ -module V is a sum of finite dimensional simple $U_q \mathfrak{k}$ -modules,
2. every finite dimensional simple $U_q \mathfrak{k}$ -module W occurs with finite multiplicity $(\dim \operatorname{Hom}_{U_q \mathfrak{k}}(W, V) < \infty)^1$.

The initial step in studying such $U_q \mathfrak{g}$ -modules was done by the authors in [14]. To produce the principal non-degenerate series of Harish-Chandra modules, it was suggested to use the approach of Beilinson-Bernstein [1]. Within this approach, in the classical case ($q = 1$) the principal non-degenerate series admits a geometric realization on the open $S(GL_2 \times GL_2)$ -orbit U in the projective variety X of flags in $\mathbb{C}^4$.

A passage to the quantum case should be started with producing a q -analogue for the open orbit U . We need a well known q -analogue of the the affine algebraic variety $\widehat{X}$ associated to X .

We are about to introduce a q -analogue $\mathbb{C}[\widehat{X}]_q$ of the algebra $\mathbb{C}[\widehat{X}]$ of regular functions on $\widehat{X}$. Start with the algebra $\mathbb{C}[SL_4]_q$ of 'regular functions' on the quantum group SL_4 . Its description in terms of generators t_{ij} , $i, j = 1, 2, 3$, and relations is well known [3]. It admits a canonical embedding $\mathbb{C}[SL_4]_q \hookrightarrow (U_q \mathfrak{g})^*$ which sends t_{ij} into matrix elements of the vector representation of $U_q \mathfrak{g}$ in the standard basis [2]. Thus $\mathbb{C}[SL_4]_q$ is equipped with a structure of $(U_q \mathfrak{g})^{\text{op}} \otimes U_q \mathfrak{g}$ -module algebra².

Assume $J \subset \{1, 2, 3, 4\}$ and $\operatorname{card}(J) \leq 3$. Introduce the notation t_J for the quantum minor of the matrix (t_{ij}) formed by the initial $\operatorname{card}(J)$ lines and the columns with indices from J . For example, $t_{\{1,2\}} = t_{11}t_{22} - qt_{12}t_{21}$, $t_{\{3,4\}} = t_{13}t_{24} - qt_{14}t_{23}$.

A complete list of relations between t_J , $\operatorname{card}(J) \leq 3$, includes commutation relations and the generalized Plücker relations [15, 10]. $\mathbb{C}[\widehat{X}]_q$ is defined as a unital subalgebra of $\mathbb{C}[SL_4]_q$ generated by t_J , $\operatorname{card}(J) \leq 3$. As a subalgebra of $\mathbb{C}[SL_4]_q$, $\mathbb{C}[\widehat{X}]_q$ has no divisors of zero. It inherits a structure of $U_q \mathfrak{g}$ -module algebra from $\mathbb{C}[SL_4]_q$. Furthermore, it

¹The class of $U_q \mathfrak{g}$ -modules we are interested in differs essentially from that considered by G. Letzter [8]. Both classes are accessible via the general approach of [7, Definition 2.2].

²For this and some other notions widespread in the quantum group theory the reader is referred to [3]. The Hopf algebra $(U_q \mathfrak{g})^{\text{op}}$ is derivable from $(U_q \mathfrak{g})$ via a replacement of its multiplication law with the opposite one.

admits a structure of $\mathbb{Z}^3$ -graded algebra :

$$\deg t_J = \begin{cases} (1, 0, 0), & \text{card}(J) = 1 \\ (0, 1, 0), & \text{card}(J) = 2 \\ (0, 0, 1), & \text{card}(J) = 3. \end{cases}$$

We use the $(U_q \mathfrak{g})^{\text{op}}$ -module structure in $\mathbb{C}[SL_4]_q^{\text{op}}$ in the above definition.

Turn to the classical case $q = 1$. Let B, N stand for the standard Borel and the unipotent subgroups of SL_4 , respectively. One has $X \simeq B \backslash SL_4$, $\widehat{X} \setminus \{0\} \simeq N \backslash SL_4$, which leads to a canonical morphism $\pi : \widehat{X} \setminus \{0\} \rightarrow X$. An inverse image of the open orbit $U \subset X$ with respect to π is the open set

$$t_{\{1,2\}}t_{\{3,4\}} (t_{\{1\}}t_{\{2,3,4\}} - t_{\{2\}}t_{\{1,3,4\}}) \neq 0.$$

Thus we obtain an embedding of the algebra of regular functions $\mathbb{C}[U]$ into the localization of $\mathbb{C}[\widehat{X}]$ with respect to the multiplicative subset $(t_{\{1,2\}}t_{\{3,4\}} (t_{\{1\}}t_{\{2,3,4\}} - t_{\{2\}}t_{\{1,3,4\}}))^{\mathbb{N}}$. Of course, the above $\mathbb{Z}^3$ -grading admits a unique extension onto this localization, and the image $\mathbb{C}[U]$ coincides with the subalgebra of all elements of degree zero. This description of $\mathbb{C}[U]$ is to be used to produce its q -analogue. The $U_q \mathfrak{sl}_2 \otimes U_q \mathfrak{sl}_2$ -invariant element

$$\eta = t_{\{1\}}t_{\{2,3,4\}} - qt_{\{2\}}t_{\{1,3,4\}}$$

of $\mathbb{C}[\widehat{X}]_q$ will work as the $SL_2 \times SL_2$ -invariant element $t_{\{1\}}t_{\{2,3,4\}} - t_{\{2\}}t_{\{1,3,4\}} \in \mathbb{C}[\widehat{X}]$.

Proposition 2.1 *The multiplicative subset $(t_{\{1,2\}}t_{\{3,4\}}\eta)^{\mathbb{N}} \subset \mathbb{C}[\widehat{X}]_q$ satisfies the Ore condition.*

Proof. It is easy to deduce commutation relations between $t_{\{1,2\}}$, $t_{\{3,4\}}$, η , and the generators t_J of $\mathbb{C}[\widehat{X}]_q$.

Describe some of those. The general properties of the universal R -matrix and the fact that all t_J are matrix elements of simple $U_q \mathfrak{g}$ -modules, imply the following

Lemma 2.2 *The elements $t_{\{1,2\}}$, $t_{\{3,4\}}$ quasi-commute with all the generators t_J of $\mathbb{C}[\widehat{X}]_q$.*

All the generators t_J are derivable from the elements

$$t_{\{2\}}, \quad t_{\{4\}}, \quad t_{\{1,2\}}, \quad t_{\{3,4\}}, \quad t_{\{2,4\}}, \quad t_{\{1,2,4\}}, \quad t_{\{2,3,4\}},$$

via an application of $E_1, E_3, E_1 E_3$. Hence, in view of the relations

$$E_j \eta = 0, \quad K_j^{\pm 1} \eta = \eta, \quad j \neq 2,$$

it suffices to obtain commutation relations between η and the generators listed above. An application of the well known properties of the quantum determinants allows one to deduce the following relations:

Lemma 2.3

$$\begin{aligned} \eta t_{\{2\}} &= q^{-1} t_{\{2\}} \eta, & \eta t_{\{4\}} &= q t_{\{4\}} \eta, \\ \eta t_{\{1,2\}} &= q^{-1} t_{\{1,2\}} \eta, & \eta t_{\{3,4\}} &= q t_{\{3,4\}} \eta, \\ \eta t_{\{2,4\}} &= q t_{\{2,4\}} \eta - (1 - q^2) t_{\{2\}} t_{\{1,2,4\}} t_{\{3,4\}}, \\ \eta t_{\{1,2,4\}} &= q^{-1} t_{\{1,2,4\}} \eta, & \eta t_{\{2,3,4\}} &= q t_{\{2,3,4\}} \eta. \end{aligned}$$

It follows from the above commutation relations that for all the generators t_J of $\mathbb{C}[\widehat{X}]_q$ one has

$$\begin{aligned} (t_{\{1,2\}}t_{\{3,4\}}\eta)^2 t_J &\in \mathbb{C}[\widehat{X}]_q t_{\{1,2\}}t_{\{3,4\}}\eta \\ t_J (t_{\{1,2\}}t_{\{3,4\}}\eta)^2 &\in t_{\{1,2\}}t_{\{3,4\}}\eta \mathbb{C}[\widehat{X}]_q, \end{aligned}$$

that is, the Ore condition is satisfied. $\square$

Corollary 2.4 *For all $l \in \mathbb{N}$*

$$\eta^l t_{\{2,4\}} = (qt_{\{2,4\}}\eta - q^2(q^{-l} - q^l)t_{\{2\}}t_{\{1,2,4\}}t_{\{3,4\}})\eta^{l-1}.$$

Let $\widehat{U} = \left\{ x \in \widehat{X} \mid t_{\{1,2\}}t_{\{3,4\}}(t_{\{1\}}t_{\{2,3,4\}} - t_{\{2\}}t_{\{1,3,4\}}) \neq 0 \right\}$, and denote by $\mathbb{C}[\widehat{U}]_q$ the localization of $\mathbb{C}[\widehat{X}]_q$ with respect to the multiplicative subset $(t_{\{1,2\}}t_{\{3,4\}}\eta)^\mathbb{N}$. Obviously, $\mathbb{C}[\widehat{X}]_q \hookrightarrow \mathbb{C}[\widehat{U}]_q$, and $\mathbb{Z}^3$ -grading is uniquely extendable from $\mathbb{C}[\widehat{X}]_q$ onto $\mathbb{C}[\widehat{U}]_q$.

Proposition 2.5 *There exists a unique extension of the structure of $U_q\mathfrak{g}$ -module algebra from $\mathbb{C}[\widehat{X}]_q$ onto $\mathbb{C}[\widehat{U}]_q$.*

Proof. The uniqueness of the extension is obvious. We are going to construct such extension by applying the following statement.

Lemma 2.6 *For every $\xi \in U_q\mathfrak{g}$, $f \in \mathbb{C}[\widehat{X}]_q$, there exists a unique Laurent polynomial $p_{\xi,f}(\lambda)$ with coefficients from $\mathbb{C}[\widehat{U}]_q$ such that*

$$p_{\xi,f}(q^l) = \xi \left(f \cdot (t_{\{1,2\}}t_{\{3,4\}}\eta)^l \right) (t_{\{1,2\}}t_{\{3,4\}}\eta)^{-l}, \quad l \in \mathbb{Z}_+.$$

We can use the same Laurent polynomials for defining $\xi \left(f \cdot (t_{\{1,2\}}t_{\{3,4\}}\eta)^l \right)$ for all integers l . Of course, we need to verify that the action of $\xi \in U_q\mathfrak{g}$ in $\mathbb{C}[\widehat{U}]_q$ is well defined, and that we obtain this way a structure of $U_q\mathfrak{g}$ -module algebra.

For that, we have to prove some identities for $\xi \in U_q\mathfrak{g}$, $\left(f_1 \cdot (t_{\{1,2\}}t_{\{3,4\}}\eta)^l \right)$, $\left(f_2 \cdot (t_{\{1,2\}}t_{\{3,4\}}\eta)^l \right)$, $f_1, f_2 \in \mathbb{C}[\widehat{U}]_q$, $l \in \mathbb{Z}$. Observe that the left and right hand sides of those identities (up to multiplying by the same powers of $t_{\{1,2\}}t_{\{3,4\}}\eta$) are just Laurent polynomials of the indeterminate $\lambda = q^l$. So, it suffices to prove them for $l \in \mathbb{Z}_+$, due to the well known uniqueness theorem for Laurent polynomials. What remains is to use the fact that $\mathbb{C}[\widehat{X}]_q$ is a $U_q\mathfrak{g}$ -module algebra. $\square$

A more general but less elementary approach to proving statements like proposition 2.5 have been obtained in a recent work by Lunts and Rosenberg [9].

Consider the subalgebra $\mathbb{C}[U]_q = \left\{ f \in \mathbb{C}[\widehat{U}]_q \mid \deg f = (0, 0, 0) \right\}$. This $U_q\mathfrak{g}$ -module algebra is a q -analogue of the algebra $\mathbb{C}[U]$ of regular functions on the open $S(GL_2 \times GL_2)$ -orbit $U \subset X$.

The following quite plausible statement allows one to embed the above $U_q\mathfrak{g}$ -module into a two-parameter family, the principal non-degenerate series.

Proposition 2.7 *There exists a unique two-parameter family $\pi_{\lambda', \lambda''}$ of representations of $U_q \mathfrak{g}$ in the vector space $\mathbb{C}[U]_q$ such that*

i) for all $l', l'' \in \mathbb{Z}$, $\xi \in U_q \mathfrak{g}$, $f \in \mathbb{C}[U]_q$,

$$(\pi_{q', q''}(\xi)f) (t_{\{1,2\}} t_{\{3,4\}})^{l'} \eta^{l''} = \xi \left(f \cdot (t_{\{1,2\}} \cdot t_{\{3,4\}})^{l'} \cdot \eta^{l''} \right)$$

ii) for all $\xi \in U_q \mathfrak{g}$, $f \in \mathbb{C}[U]_q$, the vector function $\pi_{\lambda', \lambda''}(\xi)f$ is a Laurent polynomial with coefficients from $\mathbb{C}[U]_q$.

Note that the method of 'analytic continuation' we use was suggested in [16].

Proposition 2.8 *The $U_q \mathfrak{g}$ -modules associated to $\pi_{q', q''}$, are Harish-Chandra modules.*

Proof. It suffices to consider the $U_q \mathfrak{g}$ -module $\mathbb{C}[U]_q$, i.e. the case $l' = l'' = 0$. Obviously, this $U_q \mathfrak{g}$ -module is a $\mathbb{Z}^3$ -weight module. Prove that the multiplicity of occurrence of simple $U_q \mathfrak{k}$ -modules in $\mathbb{C}[U]_q$ is the same as in the classical case $q = 1$. Equip the vector space $\mathbb{C}[U]_q$ with a filtration: $\mathbb{C}[U]_q = \bigcup_j \mathbb{C}[U]_q^{(j)}$, where

$$\mathbb{C}[U]_q^{(j)} = \left\{ f \cdot (t_{\{1,2\}} t_{\{3,4\}} \eta)^{-j} \mid f \in \mathbb{C}[\widehat{X}]_q, \deg f = (j, 2j, j) \right\}.$$

The $U_q \mathfrak{g}$ -module

$$\mathbb{C}[U]_q^{(j)} \cdot (t_{\{1,2\}} t_{\{3,4\}} \eta)^j = \left\{ f \in \mathbb{C}[\widehat{X}]_q \mid \deg f = (j, 2j, j) \right\}$$

is simple, and $t_{\{1\}}^j t_{\{1,2\}}^{2j} t_{\{1,2,3\}}^j$ is its highest weight vector. A restriction of this representation onto $U_q \mathfrak{k}$ splits into a direct sum of irreducible representations of this subalgebra with the same multiplicities as in the classical case $q = 1$. Hence the multiplicities in $\mathbb{C}[U]_q^{(j)} / \mathbb{C}[U]_q^{(j-1)}$ are also classical. What remains is to observe that in the classical case $\mathbb{C}[U]$ is a Harish-Chandra module. $\square$

We thus obtain the spherical principal non-degenerate series:

$$\pi_{q', q''}(\xi)1 = \varepsilon(\xi)1, \quad \xi \in U_q \mathfrak{k}.$$

3 Duality

Introduce the notation $V_{q', q''}$ for the Harish-Chandra modules corresponding to the representations $\pi_{q', q''}$.

The outcome of the previous section was in producing the $U_q \mathfrak{g}$ -modules $V_{q', q''}$ via demonstrating their geometric realization. This geometric realization is to be used in the present section to produce a $U_q \mathfrak{g}$ -invariant pairing $V_{q', q''} \times V_{q^{-1-l'}, q^{-2-l''}} \rightarrow \mathbb{C}$ (the associated linear map $V_{q', q''} \otimes V_{q^{-1-l'}, q^{-2-l''}} \rightarrow \mathbb{C}$ is going to be a morphism of $U_q \mathfrak{g}$ -modules). We restrict ourselves to the special case $l', l'' \in \mathbb{Z}$. A passage to the general case can be performed using the analytic continuation in the parameters $u' = q^{l'}$, $u'' = q^{l''}$ as in the proof of proposition 2.5.

Recall that $\mathbb{C}[\widehat{U}]_q = \bigoplus_{j', j'', j'''} \mathbb{C}[\widehat{U}]_q^{(j', j'', j''')}$,

$$\mathbb{C}[\widehat{U}]_q^{(j', j'', j''')} = \left\{ f \in \mathbb{C}[\widehat{U}]_q \mid \deg f = (j', j'', j''') \right\},$$

and one has well defined bilinear maps

$$\mathbb{C}[\widehat{U}]_q^{(i', i'', i''')} \times \mathbb{C}[\widehat{U}]_q^{(j', j'', j''')} \rightarrow \mathbb{C}[\widehat{U}]_q^{(i'+j', i''+j'', i''' + j''')}, \quad f_1 \times f_2 \mapsto f_1 f_2.$$

Use the canonical isomorphisms

$$V_{q^{l'}, q^{l''}} \simeq \mathbb{C}[\widehat{U}]_q^{(l'', 2l', l'')}, \quad V_{q^{-1-l'}, q^{-2-l''}} \simeq \mathbb{C}[\widehat{U}]_q^{(-2-l'', -2-2l', -2-l'')}$$

to obtain a bilinear map

$$V_{q^{l'}, q^{l''}} \times V_{q^{-1-l'}, q^{-2-l''}} \rightarrow \mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)}.$$

The non-degenerate $U_q \mathfrak{g}$ -invariant pairing

$$V_{q^{l'}, q^{l''}} \times V_{q^{-1-l'}, q^{-2-l''}} \rightarrow \mathbb{C}$$

we are interested in can be obtained as a composition of the above bilinear map and a $U_q \mathfrak{g}$ -invariant integral

$$\mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)} \rightarrow \mathbb{C}, \quad f \mapsto \int f d\nu, \quad (3.1)$$

normalized as follows:

$$\int (t_{\{1,2\}} t_{\{3,4\}} \eta^2)^{-1} d\nu = 1. \quad (3.2)$$

What remains is to prove the existence and uniqueness of the above $U_q \mathfrak{g}$ -invariant integral ν .

Proposition 3.1 *There exists a unique morphism of $U_q \mathfrak{g}$ -modules (3.1) which satisfies (3.2).*

Proof. We start with proving the uniqueness. Consider the classical case ($q = 1$). Obviously, the $U\mathfrak{k}$ -module $\mathbb{C}[U]$ of regular functions on the orbit U splits as a sum of finite dimensional isotypical components. Hence, the multiplicity of a simple $U\mathfrak{k}$ -module W in $\mathbb{C}[U]^{(j)} = \left\{ f \cdot (t_{\{1,2\}} t_{\{3,4\}} \eta)^{-j} \mid f \in \mathbb{C}[\widehat{X}], \deg f = (j, 2j, j) \right\}$ is constant for j large enough. It is easy to prove, using the standard methods of the quantum group theory [3], the existence of a similar decomposition and stabilization of the multiplicity for all the transcendental q . Hence, the multiplicity of simple $U_q \mathfrak{k}$ -modules in the $U_q \mathfrak{k}$ -module $\mathbb{C}[U]_q$ is finite. This multiplicity is the same as in the classical case since it is determined by the multiplicity of weights of finite dimensional $U_q \mathfrak{k}$ -modules $\mathbb{C}[U]_q^{(j)}$. In particular, the subspace of $U_q \mathfrak{k}$ -invariants in $\mathbb{C}[U]_q$ is one-dimensional, so we get the following

Lemma 3.2 *There exists a unique morphism of $U_q \mathfrak{k}$ -modules (3.1) which satisfies (3.2).*

Turn to the proof of existence of a morphism of $U_q \mathfrak{g}$ -modules (3.1) which satisfies (3.2).

Lemma 3.3

- i) *The element $\omega = (t_{\{1,2\}} t_{\{3,4\}} \eta^2)^{-1}$ generates the $U_q \mathfrak{g}$ -module $\mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)}$.*
- ii) *The following relations are valid: $\xi \omega = \varepsilon(\xi) \omega$ for $\xi \in U_q \mathfrak{k}$; $z \omega = \varepsilon(z) \omega$, for $z \in Z(U_q \mathfrak{g})$. The standard notation $Z(U_q \mathfrak{g})$ is used here to denote the center of $U_q \mathfrak{g}$.*

Proof. i) Since q is transcendental, one can easily reduce the proof of the first statement to the proof of a similar statement in the classical case $q = 1$. For this case the result is a consequence of the well known Kostant theorem about cyclic vectors for principal series [4] (observe that ω is the primitive weight vector with respect to the left action of $U_q\mathfrak{g}$ and its weight is -2ρ , with ρ being the half sum of positive roots).

ii) The relation $\xi\omega = \varepsilon(\xi)\omega$, $\xi \in U_q\mathfrak{k}$, is obvious.

Consider an element $z \in Z(U_q\mathfrak{g})$, together with its image ψ_z under the Harish-Chandra isomorphism. One has for $m \in \mathbb{Z}_+$:

$$z \left((t_{\{1,2\}} t_{\{3,4\}} \eta^2)^m \right) = \psi_z(2m\rho) (t_{\{1,2\}} t_{\{3,4\}} \eta^2)^m. \quad (3.3)$$

That is,

$$(t_{\{1,2\}} t_{\{3,4\}} \eta^2)^{-m} z \left((t_{\{1,2\}} t_{\{3,4\}} \eta^2)^m \right) = \psi_z(2m\rho). \quad (3.4)$$

Both sides of (3.4) are Laurent polynomials of q^m . Hence (3.3), (3.4) are valid for all $m \in \mathbb{Z}$. In particular, $z \left((t_{\{1,2\}} t_{\{3,4\}} \eta^2)^{-1} \right) = \psi_z(-2\rho) (t_{\{1,2\}} t_{\{3,4\}} \eta^2)^{-1}$. What remains is to prove that $\psi_z(-2\rho) = \varepsilon(z)$. It suffices to prove the equalities $\psi_z(-2\rho) = \psi_z(0)$, $\psi_z(0) = \varepsilon(z)$. The first of them is valid due to the invariance of ψ_z with respect to the standard action of the Weyl group. The second equality is obvious. $\square$

REMARK. It is convenient to pass from the Hopf algebras $U_q\mathfrak{k} \subset U_q\mathfrak{g}$ to their 'extensions' $U_q\mathfrak{k}^{\text{ext}} \subset U_q\mathfrak{g}^{\text{ext}}$ as in [3, chapter 8.5.3].³ (These are derived via adding certain products of $K_i^{\pm 1/4}$, $i = 1, 2, 3$). Obviously, $\mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)}$ is a $U_q\mathfrak{g}^{\text{ext}}$ -module, and lemma 3.3 can be proved for the $U_q\mathfrak{g}^{\text{ext}}$ -module structure (in fact in the proof of lemma 3.3 it is possible to use a quantum analogue of the Harish-Chandra isomorphism for algebras over the formal series of h).

Corollary 3.4 *Consider the $U_q\mathfrak{g}^{\text{ext}}$ -module V with a single generator v subject to the relations*

$$\xi v = \varepsilon(\xi)v, \quad \xi \in U_q\mathfrak{k}^{\text{ext}}; \quad zv = \varepsilon(z)v, \quad z \in Z(U_q\mathfrak{g}^{\text{ext}}).$$

The map $v \mapsto \omega$ is uniquely extendable up to an onto morphism of $U_q\mathfrak{g}^{\text{ext}}$ -modules $\varphi : V \rightarrow \mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)}$.

Turn back to the proof of proposition 3.1. Evidently, there exists a unique morphism of $U_q\mathfrak{g}^{\text{ext}}$ -modules $\mu : V \rightarrow \mathbb{C}$ such that $\mu(1) = 1$. So, the existence of a morphism $\nu : \mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)} \rightarrow \mathbb{C}$ follows from the isomorphism $\mathbb{C}[\widehat{U}]_q^{(-2, -2, -2)} \simeq V/\text{Ker } \varphi$ and the following

Lemma 3.5 $\text{Ker } \mu \supset \text{Ker } \varphi$.

Proof. It suffices to prove firstly, that the $U_q\mathfrak{k}^{\text{ext}}$ -module V splits into a sum of finite dimensional simple $U_q\mathfrak{k}^{\text{ext}}$ -modules and, secondly, that $\text{Ker } \varphi$ contains no non-zero $U_q\mathfrak{k}^{\text{ext}}$ -invariant vectors. Consider the Hopf subalgebras $U_q\mathfrak{p}_{\pm}^{\text{ext}}$ generated respectively by $\{E_j, F_j\}_{j \neq 2}, E_2, U_q\mathfrak{k}^{\text{ext}}; \{E_j, F_j\}_{j \neq 2}, F_2, U_q\mathfrak{k}^{\text{ext}}$. The first statement follows from the fact that the $U_q\mathfrak{k}^{\text{ext}}$ -module $U_q\mathfrak{g}^{\text{ext}}/U_q\mathfrak{k}^{\text{ext}} \simeq (U_q\mathfrak{p}_{+}^{\text{ext}}/U_q\mathfrak{k}^{\text{ext}}) \otimes (U_q\mathfrak{p}_{-}^{\text{ext}}/U_q\mathfrak{k}^{\text{ext}})$ splits as a sum of simple finite dimensional $U_q\mathfrak{k}^{\text{ext}}$ -modules (to prove this, one can use suitable bases in $U_q\mathfrak{g}^{\text{ext}}$, cf. [13]). The second statement is due to the following

³The reason is that the well known q -analogues of standard generators of $Z(U\mathfrak{g})$ belong to $U_q\mathfrak{g}^{\text{ext}}$ [6]

Lemma 3.6

$$\dim V^{U_q \mathfrak{g}^{\text{ext}}} = 1. \quad ^4$$

Proof. Consider a $U_q \mathfrak{g}^{\text{ext}}$ -module $U_q \mathfrak{g}^{\text{ext}} / U_q \mathfrak{k}^{\text{ext}}$. It suffices to prove that the canonical map

$$Z(U_q \mathfrak{g}^{\text{ext}}) \rightarrow (U_q \mathfrak{g}^{\text{ext}} / U_q \mathfrak{k}^{\text{ext}})^{U_q \mathfrak{k}^{\text{ext}}} \quad (3.5)$$

is onto.

Equip $U_q \mathfrak{g}^{\text{ext}}$ with a $U_q \mathfrak{k}^{\text{ext}}$ -invariant filtration via setting

$$\deg(E_2) = \deg(F_2) = 1; \quad \deg(\xi) = 0, \quad \xi \in U_q \mathfrak{k}^{\text{ext}}.$$

Evidently, the subspaces $\{\xi \in U_q \mathfrak{g}^{\text{ext}} / U_q \mathfrak{k}^{\text{ext}} \mid \deg \xi \leq j\}$ are finite dimensional. It suffices to prove that for all j the canonical map

$$\{\xi \in Z(U_q \mathfrak{g}^{\text{ext}}) \mid \deg \xi \leq j\} \rightarrow \{\xi \in (U_q \mathfrak{g}^{\text{ext}} / U_q \mathfrak{k}^{\text{ext}})^{U_q \mathfrak{k}^{\text{ext}}} \mid \deg \xi \leq j\}$$

is onto.

An application of the standard generators of $Z(U \mathfrak{g})$ allows one to reduce easily the quantum case (q transcendental) to the classical one ($q = 1$). In the classical case the surjectivity of the map (3.5) is a well known fact (see [4, Th. 1.3.1]). $\square$

References

- [1] A. Beilinson, J. Bernstein, *A generalization of Casselman's submodule theorem*, in: Representation Theory of Reductive Groups, P. Trombi (ed), Birkhäuser, Boston MA, 1983, 35 – 52.
- [2] V. G. Drinfeld, "Quantum groups", in *Proceedings of the International Congress of Mathematicians*, Berkeley, 1986, A. M. Gleason (ed.), 1987, American Mathematical Society, Providence, R. I., 798 – 820.
- [3] A. Klimyk, K. Schmüdgen, *Quantum Groups and Their Representations*, Springer, Berlin et al., 1997.
- [4] B. Kostant, *On the existence and irreducibility of certain series of representations*, Lie groups and Their Representations, I. M. Gelfand (ed), Halsted, New York, 1975, 231 – 329.
- [5] J. C. Jantzen. *Lectures on Quantum Groups*, Amer. Math. Soc., Providence RI, 1996.
- [6] M. Jimbo, *Quantum R-matrix related to the generalized Toda system: an algebraic approach*, Lecture Notes Phys. **246**, Springer, Berlin, 1986, pp. 335 – 361.
- [7] G. Letzter, *Coideal subalgebras and quantum symmetric pairs*, math.QA/0103228.
- [8] G. Letzter, *Harish-Chandra modules for quantum symmetric pairs*, Representation Theory, **4** (2000), 64 – 96.
- [9] V. A. Lunts, A. L. Rosenberg, *Localization for quantum groups*, Selecta Mathematica, New Series, **5** (1999), 123 – 150.
- [10] M. Noumi, H. Yamada, K. Mimachi, *Finite dimensional representations of the quantum group $GL_q(n, \mathbb{C})$ and the zonal spherical functions on $U_q(n-1) \backslash U_q(n)$* , Japanese J. Math. **19** (1993), No 1, 31 – 80.

⁴ $V^{U_q \mathfrak{g}^{\text{ext}}} = \{v' \in V \mid \xi v' = \varepsilon(\xi) v', \quad \xi \in U_q \mathfrak{g}^{\text{ext}}\}$

- [11] S. Sinel'shchikov and L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, Math. Phys., Anal., and Geom., **1** (1998), 75 – 100; q-alg/9703005.
- [12] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman. *On function theory in quantum disc: integral representations*, math.QA/9808015.
- [13] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. *Quantum matrix ball: differential and integral calculi*, math.QA/9905035.
- [14] D. Shklyarov, A. Stolin, S. Sinel'shchikov, and L. Vaksman, *Non-compact quantum groups and quantum Harish-Chandra modules*, Nuclear Physics B (Proc. Suppl.) 102&103 (2001), 334 – 337.
- [15] E. Taft, J. Towber, *Quantum deformations of flag schemes and Grassmann schemes I. A q -deformation of the shape-algebra of $GL(n)$* , J. Alg. **142** (1991), 1 – 36.
- [16] L. Vaksman, *Intertwining operators and quantum homogeneous spaces*, Matematicheskaya Fizika. Analiz. Geometriya, **1** (1994), No.3, 392 – 409; q-alg/9511007.

Notes of the Editor

The methods of this work can be used to produce the principal **unitary** non-degenerate series, that is, to prove the unitarizability of the Harish-Chandra modules $V_{q',q''}$ in the case $\text{Im } l' = -1/2$, $\text{Im } l'' = -1$.

Here is an outline of the proof. Let us consider the pseudo-Hermitian space $\mathbb{C}^4$ endowed with the metric $-|t_1|^2 - |t_2|^2 + |t_3|^2 + |t_4|^2$ and the projective algebraic variety of uncomplete 'Lagrange flags'

$$X_0 = \{0 \subsetneq L_1 \subsetneq L_2 \subset \mathbb{C}^4 \mid L_2 = L_2^\perp\} \subset X.$$

Let $\widehat{X}_0 \subset \widehat{X}$ be the **real** affine algebraic variety associated to X_0 . Introduce a $U_q \mathfrak{su}_{2,2}$ -module algebra $\text{Pol}(\widehat{X}_0)_q$, which is a q -analogue of the algebra of regular functions on $\widehat{X}_0$.

In the category of non-involutive algebras $\text{Pol}(\widehat{X}_0)_q$ can be defined as a subalgebra of $\mathbb{C}[\widehat{X}]_q$ generated by

$$\begin{aligned} t_{\{i\}} t_{\{j_1, j_2, j_3\}}, & \quad 1 \leq i \leq 4, \quad 1 \leq j_1 < j_2 < j_3 \leq 4, \\ t_{\{i_1, i_2\}} t_{\{j_1, j_2\}}, & \quad 1 \leq i_1 < i_2 \leq 4, \quad 1 \leq j_1 < j_2 \leq 4. \end{aligned}$$

It is easy to prove the existence and uniqueness of the involution $*$ which equips $\text{Pol}(\widehat{X}_0)_q$ with a structure of $U_q \mathfrak{su}_{2,2}$ -module algebra and such that the elements

$$(t_{\{1,2\}} t_{\{3,4\}})^j \eta^k, \quad j, k \in \mathbb{Z}_+$$

appear to be selfadjoint.⁵

⁵The uniqueness follows from the selfadjointness of the monomials $(t_{\{1,2\}} t_{\{3,4\}})^j \eta^k$ and from the fact that $\text{Pol}(\widehat{X}_0)_q$ is a sum of simple $U_q \mathfrak{su}_{2,2}$ -modules generated by the above monomials: $\text{Pol}(\widehat{X}_0)_q = \bigoplus_{j,k=0}^{\infty} L(k, 2j, k)$.

Sketch the proof of the existence. Equip every subspace $L(k, 2j, k)$ with an antilinear map $*$ which satisfies all the necessary requirements, possibly except

$$(f_1 f_2)^* = f_2^* f_1^*, \quad f_1, f_2 \in L(k, 2j, k).$$

What remains is to extend it onto $\text{Pol}(\widehat{X}_0)_q$ by antilinearity and to prove the latter identity using the above claim of uniqueness.

The involution in $\text{Pol}(\widehat{X_0})_q$ is canonically extendable onto the localization of this $*$ -algebra with respect to the multiplicatively closed set $(t_{\{1,2\}}t_{\{3,4\}})^j\eta^k$, $j, k \in \mathbb{Z}_+$.

Consider the $U_q\mathfrak{k}$ -module subalgebra of zero degree homogeneous elements. Its $U_q\mathfrak{k}$ -isotypic components are finite dimensional and are smooth functions of $q \in (0, 1]$ (this could be well rephrased rigorously). In particular, the dimensions of $U_q\mathfrak{k}$ -isotypic components are independent of $q \in (0, 1]$. Hence there exists a unique $U_q\mathfrak{k}$ -invariant integral μ on this $*$ -subalgebra with the property $\mu(1) = 1$. This integral is positive in the classical case $q = 1$, hence for all $0 < q < 1$ (otherwise for some $U_q\mathfrak{k}$ -isotypic component and some $0 < q < 1$ the $U_q\mathfrak{k}$ -invariant scalar product given by $(f_1, f_2) = \mu(f_2^* f_1)$ would appear to be non-negative but not strictly positive, hence degenerate). In view of proposition 3.1 the scalar product

$$\left(f_1 \cdot (t_{\{1,2\}}t_{\{3,4\}})^{l''}\eta^{l'''}, f_2 \cdot (t_{\{1,2\}}t_{\{3,4\}})^{l''}\eta^{l'''} \right) \stackrel{\text{def}}{=} \mu(f_2^* f_1)$$

is $U_q\mathfrak{su}_{2,2}$ -invariant in the case $\text{Im } l' = -1/2$, $\text{Im } l'' = -1$.

PART IV
ADDITIONAL RESULTS ON SOME QUANTUM
VECTOR SPACES

HIDDEN SYMMETRY OF THE DIFFERENTIAL CALCULUS ON THE QUANTUM MATRIX SPACE

S. Sinel'shchikov* L. Vaksman†

Mathematics Department, Institute for Low Temperature Physics and
Engineering, 47 Lenin Avenue, 61103 Kharkov, Ukraine

PAX: 02.40-k geometry

PAX: 03.65-Fd algebraic methods, differential geometry, topology

Abstract. A standard bicovariant differential calculus on the quantum matrix space $\text{Mat}(m, n)_q$ is considered. Our main result is in proving that the $U_q\mathfrak{sl}_m \times \mathfrak{sl}_n$ -module differential algebra $\Omega^*(\text{Mat}(m, n))_q$ is in fact a $U_q\mathfrak{sl}(m+n)$ -module differential algebra.

1. This work solves a problem whose simple special case occurs in a construction of a quantum unit ball of $\mathbb{C}^n$ (in the spirit of [10]). Within the framework of that theory, the action of the subgroup $SU(n, 1) \subset SL(n+1)$ by automorphisms of the unit matrix ball is essential. The problem is that the Wess-Zumino differential calculus in quantum $\mathbb{C}^n$ [11] at a first glance seems to be only $U_q\mathfrak{sl}_n$ -invariant. In that particular case the lost $U_q\mathfrak{sl}_{n+1}$ -symmetry can be easily detected. The main result of this work is in disclosing the hidden $U_q\mathfrak{sl}_n$ -symmetry for bicovariant differential calculus in the quantum matrix space $\text{Mat}(m, n)$. (Note that for $n = 1$ we have the case of a ball).

The authors are grateful to V. Akulov and G. Maltsiniotis for a helpful discussion of the results.

2. We start with recalling the definition of the Hopf algebra $U_q\mathfrak{sl}_N$, $N > 1$, over the field $\mathbb{C}(q)$ of rational functions of an indeterminate q [4, 5]. (We follow the notations of [3]).

For $i, j \in \{1, \dots, N-1\}$ let

$$a_{ij} = \begin{cases} 2, & i - j = 0 \\ -1, & |i - j| = 1 \\ 0, & |i - j| > 1. \end{cases}$$

The algebra $U_q\mathfrak{sl}_N$ is defined by the generators $\{E_i, F_i, K_i, K_i^{-1}\}$ and the relations

This research was partially supported by Award No UM1-2091 of the Civilian Research & Development Foundation

This lecture has been delivered at the P. Vogel seminar, Paris, 1996; published in J. Phys. A: Math. Gen., **30** (1997), L23 – L26

*Partially supported by ISF grant U2B200

E-mail address: sinelshchikov@ilt.kharkov.ua

†Partially supported by ISF grant U21200 and INTAS grant 4720

E-mail address: vaksman@ilt.kharkov.ua

$$\begin{aligned}
K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1 \\
K_i E_j &= q^{a_{ij}} E_j K_i, & K_i F_j &= q^{-a_{ij}} F_j K_i \\
E_i F_j - F_j E_i &= \delta_{ij} (K_i - K_i^{-1}) / (q - q^{-1}) \\
E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0, & |i - j| &= 1 \\
F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0, & |i - j| &= 1 \\
[E_i, E_j] &= [F_i, F_j] = 0, & |i - j| &\neq 1.
\end{aligned}$$

A comultiplication Δ , an antipode S and a counit ε are defined by

$$\begin{aligned}
\Delta E_i &= E_i \otimes 1 + K_i \otimes E_i, & \Delta F_i &= F_i \otimes K_i^{-1} + 1 \otimes F_i, \\
\Delta K_i &= K_i \otimes K_i, & S(E_i) &= -K_i^{-1} E_i, \\
S(F_i) &= -F_i K_i, & S(K_i) &= K_i^{-1}, \\
\varepsilon(E_i) &= \varepsilon(F_i) = 0, & \varepsilon(K_i) &= 1.
\end{aligned}$$

3. Remind a description of a differential algebra $\Omega^*(\text{Mat}(m, n))_q$ on a quantum matrix space [2, 8].

Let $i, j, i', j' \in \{1, 2, \dots, m + n\}$, and

$$\check{R}_{ij}^{i'j'} = \begin{cases} q^{-1}, & i = j = i' = j' \\ 1, & i' = j \text{ and } j' = i \text{ and } i \neq j \\ q^{-1} - q, & i = i' \text{ and } j = j' \text{ and } i < j \\ 0, & \text{otherwise} \end{cases}$$

$\Omega^*(\text{Mat}(m, n))_q$ is given by the generators $\{t_a^\alpha\}$ and relations

$$\begin{aligned}
\sum_{\gamma, \delta} \check{R}_{\gamma\delta}^{\alpha\beta} t_a^\gamma t_b^\delta &= \sum_{c, d} \check{R}_{ab}^{cd} t_d^\beta t_c^\alpha \\
\sum_{a', b', \gamma', \delta'} \check{R}_{\gamma'\delta'}^{\alpha\beta} \check{R}_{ab}^{a'b'} t_{a'}^{\gamma'} dt_{b'}^{\delta'} &= dt_a^\alpha t_b^\beta \\
\sum_{a', b', \gamma', \delta'} \check{R}_{\gamma'\delta'}^{\alpha\beta} \check{R}_{ab}^{a'b'} dt_{a'}^{\gamma'} dt_{b'}^{\delta'} &= -dt_a^\alpha dt_b^\beta
\end{aligned}$$

$(a, b, c, d, a', b' \in \{1, \dots, n\}; \alpha, \beta, \gamma, \delta, \gamma', \delta' \in \{1, \dots, m\})$.

Let us define a grading by $\deg(t_a^\alpha) = 0$, $\deg(dt_a^\alpha) = 1$. With that, $\mathbb{C}[\text{Mat}(m, n)]_q = \Omega^0(\text{Mat}(m, n))_q$ will stand for a subalgebra of zero degree elements.

4. Let A be a Hopf algebra and F an algebra with unit and an A -module the same time. F is said to be a A -module algebra [1] if the multiplication $m : F \otimes F \rightarrow F$ is a morphism of A -modules, and $1 \in F$ is an invariant (that is

$$a(f_1 f_2) = \sum_j a'_j f_1 \otimes a''_j f_2, \quad a1 = \varepsilon(a)1 \text{ for all } a \in A; f_1, f_2 \in F, \text{ with } \Delta(a) = \sum_j a'_j \otimes a''_j).$$

An important example of an A -module algebra appears if one supplies A^* with the structure of an A -module: $\langle af, b \rangle = \langle f, ba \rangle$, $a, b \in A$, $f \in A^*$.

5. Our immediate goal is to furnish $\mathbb{C}[\text{Mat}(m, n)]_q$ with a structure of a $U_q\mathfrak{sl}_{m+n}$ -module algebra via an embedding $\mathbb{C}[\text{Mat}(m, n)]_q \hookrightarrow (U_q\mathfrak{sl}_{m+n})^*$.

Let $\{e_{ij}\}$ be a standard basis in $\text{Mat}(m+n)$ and $\{f_{ij}\}$ the dual basis in $\text{Mat}(m+n)^*$. Consider a natural representation π of $U_q\mathfrak{sl}_{m+n}$:

$$\pi(E_i) = e_{i\ i+1}, \quad \pi(F_i) = e_{i+1\ i}, \quad \pi(K_i) = qe_{ii} + q^{-1}e_{i+1\ i+1} + \sum_{j \neq i, i+1} e_{jj}.$$

The matrix elements $u_{ij} = f_{ij}\pi \in (U_q\mathfrak{sl}_{m+n})^*$ of the natural representation may be treated as "coordinates" on the quantum group SL_{m+n} [4]. To construct "coordinate" functions on a big cell of the Grassmann manifold, we need the following elements of $\mathbb{C}[\text{Mat}(m, n)]_q$

$$x(j_1, j_2, \dots, j_m) = \sum_{w \in S_m} (-q)^{l(w)} u_{1j_{w(1)}} u_{2j_{w(2)}} \dots u_{mj_{w(m)}},$$

with $1 \leq j_1 < j_2 < \dots < j_m \leq m+n$, and $l(w) = \text{card}\{(a, b) \mid a < b \text{ and } w(a) > w(b)\}$ being the "length" of a permutation $w \in S_m$.

Proposition 0.7 $x(1, 2, \dots, m)$ is invertible in $(U_q\mathfrak{sl}_{m+n})^*$, and the map

$$t_a^\alpha \mapsto x(1, 2, \dots, m)^{-1} x(1, \dots, \widehat{m+1-\alpha}, \dots, m, m+a)$$

can be extended up to an embedding

$$i : \mathbb{C}[\text{Mat}(m, n)]_q \hookrightarrow (U_q\mathfrak{sl}_{m+n})^*.$$

(The sign $\widehat{}$ here indicates the item in a list that should be omitted).

Proposition 0.7 allows one to equip $\mathbb{C}[\text{Mat}(m, n)]_q$ with the structure of a $U_q\mathfrak{sl}_{m+n}$ -module algebra :

$$i\xi t_a^\alpha = \xi i t_a^\alpha, \quad \xi \in U_q\mathfrak{sl}_{m+n}, \quad a \in \{1, \dots, n\}, \quad \alpha \in \{1, \dots, m\}.$$

6. The main result of our work is the following

Theorem 0.8 $\Omega^*(\text{Mat}(m, n))_q$ admits a unique structure of a $U_q\mathfrak{sl}_{m+n}$ -module algebra such that the embedding

$$i : \mathbb{C}[\text{Mat}(m, n)]_q \hookrightarrow \Omega^*(\text{Mat}(m, n))_q$$

and the differential

$$d : \Omega^*(\text{Mat}(m, n))_q \rightarrow \Omega^*(\text{Mat}(m, n))_q$$

are the morphisms of $U_q\mathfrak{sl}_{m+n}$ -modules.

REMARK 1. The bicovariance of the differential calculus on the quantum matrix space allows one to equip the algebra $\Omega^*(\text{Mat}(m, n))_q$ with a structure of $U_q\mathfrak{sl}(\mathfrak{gl}_m \times \mathfrak{gl}_n)$ -module, which is compatible with multiplication in $\Omega^*(\text{Mat}(m, n))_q$ and differential d . Theorem 0.8 implies that $\Omega^*(\text{Mat}(m, n))_q$ possess an additional hidden symmetry since $U_q\mathfrak{sl}_{m+n} \supsetneq U_q\mathfrak{sl}(\mathfrak{gl}_m \times \mathfrak{gl}_n)$.

REMARK 2. Let $q_0 \in \mathbb{C}$ and q_0 is not a root of unity. It follows from the explicit formulae for $E_m t_a^\alpha$, $F_m t_a^\alpha$, $K_m^{\pm 1} t_a^\alpha$, $a \in \{1, \dots, n\}$, $\alpha \in \{1, \dots, m\}$, that the "specialization" $\Omega^*(\text{Mat}(m, n))_{q_0}$ is a $U_{q_0} \mathfrak{sl}_{m+n}$ -module algebra.

7. Supply the algebra $U_q \mathfrak{sl}_{m+n}$ with a grading as follows:

$$\deg(K_i) = \deg(E_i) = \deg(F_i) = 0, \quad \text{for } i \neq m,$$

$$\deg(K_m) = 0, \quad \deg(E_m) = 1, \quad \deg(F_m) = 0.$$

The proofs of Proposition 0.7 and Theorem 0.8 reduce to the construction of graded $U_q \mathfrak{sl}_{m+n}$ -modules which are dual respectively to the modules of functions $\Omega^0(\text{Mat}(m, n))_q$ and that of 1-forms $\Omega^1(\text{Mat}(m, n))_q$. The dual modules are defined by their generators and correlations. While proving the completeness of the correlation list we implement the "limit specialization" $q_0 = 1$ (see [3, p. 416]).

The passage from the order one differential calculus $\Omega^0(\text{Mat}(m, n))_q \xrightarrow{d} \Omega^1(\text{Mat}(m, n))_q$ to $\Omega^*(\text{Mat}(m, n))_q$ is done via a universal argument described in a paper by G. Maltsiniotis [9]. This argument doesn't break $U_q \mathfrak{sl}_{m+n}$ -symmetry.

8. Our approach to the construction of order one differential calculus is completely analogous to that of V. Drinfel'd [4] used initially to produce the algebra of functions on a quantum group by means of a universal enveloping algebra.

9. The space of matrices is the simplest example of an irreducible prehomogeneous vector space of parabolic type [7]. Such space can be also associated to a pair constituted by a Dynkin diagram of a simple Lie algebra $\mathcal{G}$ and a distinguished vertex of this diagram. Our method can work as an efficient tool for producing $U_q \mathcal{G}$ -invariant differential calculi on the above prehomogeneous vector spaces.

Note that $U_q \mathcal{G}$ -module algebras of polynomials on quantum prehomogeneous spaces of parabolic type were considered in a recent work of M. S. Kebe [6].

Acknowledgement

The authors would like to express their gratitude to Prof. A. Boutet de Monvel at University Paris VII for the warm hospitality during the work on this paper.

References

- [1] E. Abe. Hopf Algebras, Cambridge Tracts in Mathematics 74, Cambridge University Press, Cambridge, 1980.
- [2] V. Chari and A. Pressley. A guide to quantum groups. Cambridge Univ. Press, 1994.
- [3] C. De Concini and V. G. Kac, *Representations of quantum groups at roots of 1*, in Operator Algebras, Unitary representations, Enveloping Algebras and Invariant Theory, A. Connes, M. Duflo, A. Joseph, R. Rentschler (eds.), pp. 471- 506, Birkhäuser, Boston, 1990.
- [4] V. G. Drinfel'd , *Quantum groups*, in Proceedings of International Congress of Mathematicians, Berkeley, 1986, A. M. Gleason (ed.), pp. 798 – 820, American Mathematical Society, Providence RI, 1987.

- [5] M. Jimbo, *Quantum R-matrix related to the generalized Toda system: an algebraic approach*, in Field Theory, Quantum Gravity and Strings, H. J. de Vega & N. Sanchez (eds.), Lecture Notes in Phys., **246**, 335 – 361, Springer, Berlin, 1986.
- [6] Mâlek Stefan Kebe, *$\mathcal{O}$ -algèbres quantiques*, C. R. Acad. Sci. Paris, **322** (1996), Série 1, 1 – 4.
- [7] T. Kimura, *A classification theory of prehomogeneous vector spaces*, Advanced Studies in Pure Mathematics **14**, 1988, Representations of Lie groups, Kyoto, Hiroshima, 1986, p. 223 – 256.
- [8] G. Maltsiniotis, *Groupes quantiques et structures différentielles*, C. R. Acad. Sci., Paris, Série I, **311** (1990), 831 – 834.
- [9] G. Maltsiniotis, *Le langage des espaces et des groupes quantiques*, Comm. Math. Phys., **151** (1993), 275 – 302.
- [10] W. Rudin. Function Theory in the Unit Ball of $\mathbb{C}^n$. Springer – Verlag, N.-Y., 1980.
- [11] J. Wess and B. Zumino, *Covariant differential calculus on the quantum hyperplane*, Nucl. Phys. B, Proc. Suppl. **18 B**, 1991, 302 – 312.

Notes of the Editor

The proofs of the results announced in this work can be found in [1].

References

- [1] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035.

q-ANALOGS OF CERTAIN PREHOMOGENEOUS VECTOR SPACES: COMPARISON OF SEVERAL APPROACHES

D. Shklyarov

1 Introduction: prehomogeneous vector spaces of commutative parabolic type

Let $\mathfrak{g}$ be a complex simple Lie algebra, $\mathfrak{h}$ its Cartan subalgebra, $\{\alpha_i\}_{i=\overline{1,l}}$ the simple roots with respect to $\mathfrak{h}$. Let us associate to each $\alpha_0 \in \{\alpha_i\}_{i=\overline{1,l}}$ the element $H_0 \in \mathfrak{h}$ such that

$$\alpha_i(H_0) = \begin{cases} 2, & \alpha_i = \alpha_0, \\ 0, & \text{otherwise.} \end{cases}$$

Such H_0 can be used to equip $\mathfrak{g}$ with a $\mathbb{Z}$ -grading as follows

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_i, \quad \mathfrak{g}_i \stackrel{\text{def}}{=} \{\xi \in \mathfrak{g} \mid [H_0, \xi] = 2i\xi\}.$$

If $\mathfrak{g}_i$ is nonzero only for $i \in \{-1, 0, 1\}$ then the subspace $\mathfrak{g}_{-1}$ is said to be a prehomogeneous vector space of commutative parabolic type (see [5]).

REMARK 1. Let us explain the adjective "prehomogeneous". Let $K \subset \text{Aut } \mathfrak{g}$ be the subgroup of those automorphisms which preserve the decomposition

$$\mathfrak{g} = \mathfrak{g}_{-1} \bigoplus \mathfrak{g}_0 \bigoplus \mathfrak{g}_{+1}.$$

Then $\mathfrak{g}_0 = \text{Lie } K$. The group K acts in $\mathfrak{g}_{-1}$ in such way that there exists a Zariski open K -orbit. A pair (G, V) (G is an algebraic group acting in the vector space V) which possesses this property is called a prehomogeneous vector space.

Since K acts in $\mathfrak{g}_{-1}$ and $\mathfrak{g}_0 = \text{Lie } K$, one may consider the corresponding representation of $U\mathfrak{g}_0$ in the space $\mathbb{C}[\mathfrak{g}_{-1}]$ of holomorphic polynomials on $\mathfrak{g}_{-1}$.

REMARK 2. The Killing form of $\mathfrak{g}$ makes the vector spaces $\mathfrak{g}_{-1}$ and $\mathfrak{g}_{+1}$ dual to each other. This allows one to identify the algebras $\mathbb{C}[\mathfrak{g}_{-1}]$ and $S(\mathfrak{g}_{+1})$ (the symmetric algebra over $\mathfrak{g}_{+1}$). The latter algebra is isomorphic to $U\mathfrak{g}_{+1}$ for $\mathfrak{g}_{+1}$ is an Abelian Lie subalgebra in $\mathfrak{g}$. The action of $U\mathfrak{g}_0$ in $\mathbb{C}[\mathfrak{g}_{-1}]$ we deal with corresponds (under the isomorphism $\mathbb{C}[\mathfrak{g}_{-1}] \simeq U\mathfrak{g}_{+1}$) to the adjoint action of $U\mathfrak{g}_0$ in $U\mathfrak{g}_{+1}$.

There exist several approaches to constructing a q -analog of the algebra $\mathbb{C}[\mathfrak{g}_{-1}]$. In the present paper we concern with those developed in [2], [3], [9]. Within framework of each

This research was supported in part by Award No UM1-2091 of the US Civilian Research & Development Foundation

This lecture has been delivered at the seminar 'Quantum groups', Kharkov, April 2000; published in *Matematicheskaya Fizika. Analiz. Geometriya*, **8** (2001), No 3, 325 – 345.

approach a (noncommutative) analog of $\mathbb{C}[\mathfrak{g}_{-1}]$ is endowed with an action of the quantum universal enveloping algebra $U_q \mathfrak{g}_0$. We prove that q -analogs of $\mathbb{C}[\mathfrak{g}_{-1}]$ constructed in [2], [3], [9] are isomorphic as $U_q \mathfrak{g}_0$ -module algebras.

For the sake of simplicity we carry out this program for $\mathfrak{g} = \mathfrak{sl}_4(\mathbb{C})$, $\alpha_0 = \alpha_2$ (in this case $\mathfrak{g}_{-1}$ is the space of 2×2 complex matrices, $K = S(GL_2(\mathbb{C}) \times GL_2(\mathbb{C}))$). But we present such proofs that are transferable to the case of an arbitrary prehomogeneous vector space of commutative parabolic type.

The author thanks L. Vaksman for stating the problem and discussing the results.

2 Notation and auxiliary facts

Let $\mathfrak{g} = \mathfrak{sl}_4(\mathbb{C})$. For convenience, we identify $\mathfrak{g}$ with the Lie algebra of 4×4 complex matrices with zero trace. Let $\mathfrak{h} \subset \mathfrak{g}$ be the Cartan subalgebra of diagonal matrices. Denote by Δ and W the root system of $\mathfrak{g}$ with respect to $\mathfrak{h}$ and the Weyl group of this system, respectively. Let also $\alpha_1, \alpha_2, \alpha_3$ be the simple roots in Δ given by

$$\alpha_i(H) = a_i - a_{i+1}$$

with $H = \text{diag}(a_1, a_2, a_3, a_4) \in \mathfrak{h}$.

There exists an isomorphism of the group W onto the group S_4 such that the simple reflections $s_{\alpha_1}, s_{\alpha_2}, s_{\alpha_3}$ correspond to the transpositions $(1, 2), (2, 3), (3, 4)$. Let $\Delta_+ \subset \Delta$ be the system of positive roots:

$$\Delta_+ = \{\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3\}.$$

Denote by $(\cdot | \cdot)$ the W -invariant scalar product in $\mathfrak{h}^*$ such that $(\alpha_i | \alpha_i) = 2$.

The root α_2 plays the role of the 'distinguished' root α_0 (see Introduction). The associated element $H_0 \in \mathfrak{h}$ is given by $H_0 = \text{diag}(1, 1, -1, -1)$, or

$$H_0 = H_1 + 2H_2 + H_3 \tag{2.1}$$

with $H_1 = \text{diag}(1, -1, 0, 0)$, $H_2 = \text{diag}(0, 1, -1, 0)$, $H_3 = \text{diag}(0, 0, 1, -1)$.

Let $\Delta_c \stackrel{\text{def}}{=} \{\alpha_1, \alpha_3, -\alpha_1, -\alpha_3\} \subset \Delta$, $\Delta_n \stackrel{\text{def}}{=} \Delta \setminus \Delta_c$. Then

$$\mathfrak{g}_0 = \mathfrak{h} \oplus \left(\bigoplus_{\alpha \in \Delta_c} \mathfrak{g}_\alpha \right),$$

$$\mathfrak{g}_{+1} = \bigoplus_{\alpha \in \Delta_+ \cap \Delta_n} \mathfrak{g}_\alpha, \quad \mathfrak{g}_{-1} = \bigoplus_{-\alpha \in \Delta_+ \cap \Delta_n} \mathfrak{g}_\alpha,$$

with $\mathfrak{g}_\alpha$ being the root subspace in $\mathfrak{g}$ corresponding to $\alpha \in \Delta$.

Let W_c be the subgroup in W generated by $s_{\alpha_1}, s_{\alpha_3}$. Thus, $W_c \simeq S_2 \times S_2$ is the Weyl group of the Lie subalgebra $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2 \in \mathfrak{g}$.

In the rest of this paper the ground field will be the field of rational functions $\mathbb{C}(q^{1/4})$.

Let us recall some definitions and facts of the quantum group theory (we follow [4]).

The quantum universal enveloping algebra $U_q \mathfrak{g}$ is the algebra with the generators $\{E_i, F_i, K_i^{\pm 1}\}_{i=\overline{1,3}}$ satisfying the following relations

$$K_i K_j = K_j K_i, \quad K_i K_i^{-1} = K_i^{-1} K_i = 1,$$

$$\begin{aligned}
 K_i E_j &= q^{a_{ij}} E_j K_i, & K_i F_j &= q^{-a_{ij}} F_j K_i, \\
 E_i F_j - F_j E_i &= \delta_{ij} \frac{K_i - K_i^{-1}}{q - q^{-1}}, \\
 \sum_{s=0}^{1-a_{ij}} (-1)^s \begin{bmatrix} 1-a_{ij} \\ s \end{bmatrix}_q E_i^{1-a_{ij}-s} E_j E_i^s &= 0, & i \neq j, \\
 \sum_{s=0}^{1-a_{ij}} (-1)^s \begin{bmatrix} 1-a_{ij} \\ s \end{bmatrix}_q F_i^{1-a_{ij}-s} F_j F_i^s &= 0, & i \neq j,
 \end{aligned}$$

where (a_{ij}) is the Cartan matrix for $\mathfrak{g}$:

$$a_{ij} = \begin{cases} 2 & , \quad i - j = 0 \\ -1 & , \quad |i - j| = 1 \\ 0 & , \quad \text{otherwise} \end{cases}$$

$$\begin{bmatrix} n \\ j \end{bmatrix}_q \stackrel{\text{def}}{=} \frac{[n]_q!}{[n-j]_q! [j]_q!}, \quad [n]_q! \stackrel{\text{def}}{=} [n]_q \cdot [n-1]_q \cdot \dots \cdot [1]_q, \quad [n]_q \stackrel{\text{def}}{=} \frac{q^n - q^{-n}}{q - q^{-1}}.$$

The algebra $U_q \mathfrak{g}$ is endowed with a structure of a Hopf algebra as follows

$$\begin{aligned}
 \Delta(E_i) &= E_i \otimes 1 + K_i \otimes E_i, & \Delta(F_i) &= F_i \otimes K_i^{-1} + 1 \otimes F_i, & \Delta(K_i) &= K_i \otimes K_i, \\
 S(E_i) &= -K_i^{-1} E_i, & S(F_i) &= -F_i K_i, & S(K_i) &= K_i^{-1}, \\
 \varepsilon(E_i) &= \varepsilon(F_i) = 0, & \varepsilon(K_i) &= 1,
 \end{aligned}$$

with Δ , S , ε being the comultiplication, the antipode, and the counit, respectively.

Let us use the short notation $x_{(1)} \otimes x_{(2)}$ for the element $\Delta(x) \in U_q \mathfrak{g} \otimes U_q \mathfrak{g}$ ($x \in U_q \mathfrak{g}$). For example, coassociativity of the comultiplication $\Delta : U_q \mathfrak{g} \rightarrow U_q \mathfrak{g} \otimes U_q \mathfrak{g}$ looks in this notation as follows

$$x_{(1)} \otimes x_{(2)(1)} \otimes x_{(2)(2)} = x_{(1)(1)} \otimes x_{(1)(2)} \otimes x_{(2)}. \quad (2.2)$$

Sometimes we use the notation $x_{(1)} \otimes x_{(2)} \otimes x_{(3)}$ for the right (and left) hand side of (2.2). Then the obvious meaning has the notation $x_{(1)} \otimes x_{(2)} \otimes x_{(3)} \otimes x_{(4)}$ etc.

The adjoint representation of the algebra $U_q \mathfrak{g}$ is defined as follows

$$\text{ad}x(y) \stackrel{\text{def}}{=} x_{(1)} \cdot y \cdot S(x_{(2)})$$

with $x, y \in U_q \mathfrak{g}$. This adjoint action makes $U_q \mathfrak{g}$ a $U_q \mathfrak{g}$ -module algebra. It means that the product map $U_q \mathfrak{g} \otimes U_q \mathfrak{g} \rightarrow U_q \mathfrak{g}$ is a morphism of $U_q \mathfrak{g}$ -modules and the unit $1 \in U_q \mathfrak{g}$ is $U_q \mathfrak{g}$ -invariant.

We fix the following notation for some subalgebras in $U_q \mathfrak{g}$:

$$\begin{aligned}
 U_q^{\geq 0} &= \langle E_i, K_i^{\pm 1} \mid i = \overline{1, 3} \rangle, & U_q^{\leq 0} &= \langle F_i, K_i^{\pm 1} \mid i = \overline{1, 3} \rangle, \\
 U_q^+ &= \langle E_i \mid i = \overline{1, 3} \rangle, & U_q^- &= \langle F_i \mid i = \overline{1, 3} \rangle, \\
 U_q^0 &= \langle K_i^{\pm 1} \mid i = \overline{1, 3} \rangle, & U_q \mathfrak{g}_0 &= \langle K_i^{\pm 1}, E_j, F_j \mid i = \overline{1, 3}, \quad j \neq 2 \rangle.
 \end{aligned}$$

Let us recall a definition of the Lusztig automorphisms T_i , $i = \overline{1, 3}$, of the algebra $U_q \mathfrak{g}$. The action of T_i on the subalgebra $U_q^{\geq 0}$ is given by

$$\begin{aligned} T_i(K_j) &= K_j \cdot K_i^{-a_{ij}}, & T_i(E_i) &= -F_i K_i, \\ T_i(E_j) &= (-\text{ad } E_i)^{-a_{ij}}(E_j), & i &\neq j. \end{aligned}$$

To define T_i completely one sets

$$T_i \circ k = k \circ T_i,$$

where k is the conjugate linear antiautomorphism of the $\mathbb{C}(q^{1/4})$ -algebra $U_q \mathfrak{g}$ given by

$$k(E_i) = F_i, \quad k(F_i) = E_i, \quad k(K_i) = K_i^{-1}, \quad k(q^{1/4}) = q^{-1/4}.$$

Let $w \in W$ and $w = s_{i_1} s_{i_2} \dots s_{i_k}$ be a reduced expression (we write ' s_i ' instead of s_{α_i}). It is well known that the automorphism $T_w \stackrel{\text{def}}{=} T_{i_1} T_{i_2} \dots T_{i_k}$ does not depend on particular choice of a reduced expression of w .

All $U_q \mathfrak{g}$ -modules we consider possess the property

$$V = \bigoplus_{\mu \in \mathbb{Z}^3} V_\mu, \quad V_\mu \stackrel{\text{def}}{=} \{v \in V \mid K_i v = q^{\mu_i} v, i = \overline{1, 3}\}$$

with $\mu = (\mu_1, \mu_2, \mu_3)$. This allows one to introduce endomorphisms H_i , $i = \overline{1, 3}$, of any $U_q \mathfrak{g}$ -module V by

$$H_i v = \mu_i v \quad \Leftrightarrow \quad v \in V_\mu, \quad \mu = (\mu_1, \mu_2, \mu_3).$$

Formally this can be written as $K_i = q^{H_i}$.

Let $K_0 \stackrel{\text{def}}{=} K_1 \cdot K_2^2 \cdot K_3$ (i.e. $K_0 = q^{H_0}$ with H_0 given by (2.1)). It is an important consequence of definitions that K_0 belongs to the centre of the algebra $U_q \mathfrak{g}_0$:

$$\text{ad } K_0(\xi) = \xi, \quad \xi \in U_q \mathfrak{g}_0. \quad (2.3)$$

Let us recall one some facts concerning the universal R -matrix for $U_q \mathfrak{g}$ (in context of the present paper the universal R -matrix have to be understood just as in [4]).

R satisfies some well known identities. We don't adduce a full list of these identities but recall one those important for us:

$$\text{id} \otimes \Delta^{\text{op}}(R) = R^{12} \cdot R^{13}, \quad (2.4)$$

$$\Delta^{\text{op}}(\eta) \cdot R = R \cdot \Delta(\eta), \quad \eta \in U_q \mathfrak{g}, \quad (2.5)$$

with $\Delta^{\text{op}}(x) \stackrel{\text{def}}{=} x_{(2)} \otimes x_{(1)}$, $R^{12} \stackrel{\text{def}}{=} \sum_i a_i \otimes b_i \otimes 1$, $R^{13} \stackrel{\text{def}}{=} \sum_i a_i \otimes 1 \otimes b_i$.

Remind an explicit formula for the R -matrix (the so-called multiplicative formula). Let $w_0 \in W$ be the maximal length element. Identifying W with S_4 we get

$$w_0 = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix}.$$

The length of any reduced expression of w_0 is equal to 6. To a reduced expression $w_0 = s_{i_1} s_{i_2} \dots s_{i_6}$ one attaches the following data:

i) the total order on the set Δ_+ of positive roots:

$$\beta_1 = \alpha_{i_1}, \quad \beta_2 = s_{i_1}(\alpha_{i_2}), \quad \dots \quad \beta_6 = s_{i_1}s_{i_2}\dots s_{i_5}(\alpha_{i_6});$$

ii) the set of elements $E_{\beta_1}, E_{\beta_2}, \dots, E_{\beta_6} \in U_q^+$, $F_{\beta_1}, F_{\beta_2}, \dots, F_{\beta_6} \in U_q^-$ which are q -analogs of root vectors in $\mathfrak{g}$:

$$E_{\beta_1} = E_{i_1}, \quad E_{\beta_2} = T_{i_1}(E_{i_2}), \quad \dots \quad E_{\beta_6} = T_{i_1}T_{i_2}\dots T_{i_5}(E_{i_6}),$$

$$F_{\beta_1} = F_{i_1}, \quad F_{\beta_2} = T_{i_1}(F_{i_2}), \quad \dots \quad F_{\beta_6} = T_{i_1}T_{i_2}\dots T_{i_5}(F_{i_6});$$

iii) the multiplicative formula for the R -matrix:
 $R =$

$$\exp_{q^2}((q^{-1}-q)E_{\beta_6} \otimes F_{\beta_6}) \cdot \dots \cdot \exp_{q^2}((q^{-1}-q)E_{\beta_2} \otimes F_{\beta_2}) \cdot \exp_{q^2}((q^{-1}-q)E_{\beta_1} \otimes F_{\beta_1}) \cdot q^t \quad (2.6)$$

with $t \stackrel{\text{def}}{=} -\sum_{i,j} c_{ij} H_i \otimes H_j$, the matrix (c_{ij}) being the inverse to the Cartan matrix, and

$$\exp_{q^2}(t) \stackrel{\text{def}}{=} \sum_{k=0}^{\infty} \frac{t^k}{(k)_{q^2}!}, \quad (k)_{q^2}! \stackrel{\text{def}}{=} \prod_{j=1}^k \frac{1-q^{2j}}{1-q^2}.$$

3 Three approaches to quantization of prehomogeneous vector spaces of commutative parabolic type

3.1 First approach

In this subsection we describe an approach to constructing q -analogs of prehomogeneous vector spaces of commutative parabolic type developed in [9].

Let us consider the generalized Verma module $V(0)$ over $U_q\mathfrak{g}$ given by its generator $v(0)$ and the relations

$$E_i v(0) = 0, \quad K_i v(0) = v(0), \quad i = \overline{1, 3}, \quad (3.1)$$

$$F_i v(0) = 0, \quad i \neq 2. \quad (3.2)$$

$V(0)$ splits into direct sum of its finite dimensional subspaces $V(0)_k$, $-k \in \mathbb{Z}_+$, with

$$V(0)_k \stackrel{\text{def}}{=} \{v \in V(0) \mid H_0 v = 2kv\}.$$

Consider the *graded* dual $U_q\mathfrak{g}$ -module:

$$\mathbb{C}[\mathfrak{g}_{-1}]_q \stackrel{\text{def}}{=} \bigoplus_{-k \in \mathbb{Z}_+} (V(0)_k)^*.$$

Let us equip the tensor product $V(0) \otimes V(0)$ with a $U_q\mathfrak{g}$ -module structure as follows

$$\xi(v_1 \otimes v_2) = \xi_{(2)}(v_1) \otimes \xi_{(1)}(v_2), \quad \xi \in U_q\mathfrak{g}, \quad v_1, v_2 \in V(0). \quad (3.3)$$

Due to (3.1), (3.2) the maps

$$v(0) \mapsto v(0) \otimes v(0), \quad v(0) \mapsto 1 \quad (3.4)$$

are extendable up to morphisms of $U_q\mathfrak{g}$ -modules

$$\Delta_- : V(0) \rightarrow V(0) \bigotimes V(0), \quad \varepsilon_- : V(0) \rightarrow \mathbb{C}(q^{1/4}).$$

It can be shown that Δ_- and ε_- make $V(0)$ a coassociative coalgebra with a counit. Thus, the dual maps

$$m = (\Delta_-)^* : \mathbb{C}[\mathfrak{g}_{-1}]_q \bigotimes \mathbb{C}[\mathfrak{g}_{-1}]_q \rightarrow \mathbb{C}[\mathfrak{g}_{-1}]_q, \quad 1 = (\varepsilon_-)^* : \mathbb{C}(q^{1/4}) \rightarrow \mathbb{C}[\mathfrak{g}_{-1}]_q \quad (3.5)$$

make $\mathbb{C}[\mathfrak{g}_{-1}]_q$ an associative unital algebra. Moreover, the product map m is a morphism of $U_q\mathfrak{g}$ -modules and the unit 1 is $U_q\mathfrak{g}$ -invariant, i.e. $\mathbb{C}[\mathfrak{g}_{-1}]_q$ is a $U_q\mathfrak{g}$ -module algebra. In particular, it is a $U_q\mathfrak{g}_0$ -module algebra. This $U_q\mathfrak{g}_0$ -module structure is just the one we mentioned in the Introduction.

3.2 Second approach

Now we are going to describe briefly an approach of H. P. Jakobsen [2] to quantization of $\mathbb{C}[\mathfrak{g}_{-1}]$.

It follows from the definition of $U_q\mathfrak{g}$ that

$$\text{ad}K_2(E_2) = q^2 E_2, \quad \text{ad}K_1(E_2) = \text{ad}K_3(E_2) = q^{-1} E_2,$$

$$\text{ad}F_1(E_2) = \text{ad}F_3(E_2) = 0,$$

$$(\text{ad}E_1)^2(E_2) = (\text{ad}E_3)^2(E_2) = 0.$$

Thus $\text{ad}U_q\mathfrak{g}_0(E_2)$ is a finite dimensional $U_q\mathfrak{g}_0$ -submodule in $U_q\mathfrak{g}$. Let us denote by $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}$ the minimal subalgebra in $U_q\mathfrak{g}$ which contains the subspace $\text{ad}U_q\mathfrak{g}_0(E_2)$. Evidently, it is a $U_q\mathfrak{g}_0$ -module subalgebra in $U_q\mathfrak{g}$. The algebra $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}$ can be treated as a q -analog of $\mathbb{C}[\mathfrak{g}_{-1}]$ (see Remark 2 above).

3.3 Third approach

Let us turn to description of an approach of A. Kamita, Y. Morita, and T. Tanisaki [3]. Note that notation in [3] differs from ours.

Let $w'_0 \in W_c$ be the maximal length element. Evidently, $w'_0 = (1, 2) \cdot (3, 4)$. Consider the subspace $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$ in $U_q\mathfrak{g}$ defined by

$$\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}} = U_q^+ \cap T_{w'_0}^{-1}(U_q^+). \quad (3.6)$$

Obviously the subspace $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$ is a subalgebra in $U_q\mathfrak{g}$. It is shown in [3] that $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$ is a $U_q\mathfrak{g}_0$ -module subalgebra in $U_q\mathfrak{g}$ with respect to the adjoint action. It is one more q -analog of the algebra $\mathbb{C}[\mathfrak{g}_{-1}]$.

4 Comparison of the approaches

Note that both the algebras $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}$ and $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$ lie within the quantum universal enveloping algebra $U_q\mathfrak{g}$. Our aim is:

- i) to construct an embedding T of the algebra $\mathbb{C}[\mathfrak{g}_{-1}]_q$ into $U_q\mathfrak{g}$ which intertwines the $U_q\mathfrak{g}_0$ -action in $\mathbb{C}[\mathfrak{g}_{-1}]_q$ mentioned in subsection 3.1 and the adjoint $U_q\mathfrak{g}_0$ -action;
- ii) to show that the subalgebras $T(\mathbb{C}[\mathfrak{g}_{-1}]_q)$, $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}$, and $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$ in $U_q\mathfrak{g}$ coincide.

4.1 $\mathbb{C}[\mathfrak{g}_{-1}]_q \simeq \mathbb{C}[\mathfrak{g}_{-1}]_q^I$

In this subsection we construct an $U_q\mathfrak{g}_0$ -invariant embedding T of the algebra $\mathbb{C}[\mathfrak{g}_{-1}]_q$ into $U_q\mathfrak{g}$, and then we show that $T(\mathbb{C}[\mathfrak{g}_{-1}]_q) = \mathbb{C}[\mathfrak{g}_{-1}]_q^I$. The embedding T is constructed via a standard technique due to [1].

Let $R = \sum_i a_i \otimes b_i$ be the universal R -matrix for $U_q\mathfrak{g}$. Consider the linear map $T : \mathbb{C}[\mathfrak{g}_{-1}]_q \rightarrow U_q\mathfrak{g}$ given by

$$T(f) \stackrel{\text{def}}{=} \sum_i a_i \langle b_i v(0), f \rangle \quad (4.1)$$

with $f \in \mathbb{C}[\mathfrak{g}_{-1}]_q$, $v(0)$ being the generator of the generalized Verma module $V(0)$, $\langle \cdot, \cdot \rangle$ being the pairing $V(0) \times \mathbb{C}[\mathfrak{g}_{-1}]_q \rightarrow \mathbb{C}(q^{1/4})$ arising from the equality $\mathbb{C}[\mathfrak{g}_{-1}]_q = (V(0))^*$ (see subsection 3.1).

Let us comment the definition of T . Using the multiplicative formula (2.6) and the definition of the $U_q\mathfrak{g}$ -module $V(0)$ one shows that $\sum_i a_i \otimes b_i v(0)$ is a formal series of elements from $U_q^+ \otimes V(0)$. We will prove later (see proof of Proposition 4.5) that the right hand side of (4.1) is a finite sum for any $f \in \mathbb{C}[\mathfrak{g}_{-1}]_q$.

Proposition 4.1 *T is a homomorphism of algebras.*

Proof. Let $f, \varphi \in \mathbb{C}[\mathfrak{g}_{-1}]_q$. Then due to (3.5), (3.4), (3.3)

$$\begin{aligned} T(f \cdot \varphi) &= \sum_i a_i \langle b_i v(0), f \cdot \varphi \rangle = \sum_i a_i \langle b_i v(0), m(f \otimes \varphi) \rangle = \\ &= \sum_i a_i \langle \Delta_-(b_i v(0)), f \otimes \varphi \rangle = \sum_i a_i \langle \Delta^{\text{op}}(b_i)(v(0) \otimes v(0)), f \otimes \varphi \rangle. \end{aligned}$$

By (2.4)

$$\begin{aligned} \sum_i a_i \langle \Delta^{\text{op}}(b_i)(v(0) \otimes v(0)), f \otimes \varphi \rangle &= \sum_{i,j} a_i \cdot a_j \langle b_i \otimes b_j(v(0) \otimes v(0)), f \otimes \varphi \rangle = \\ &= \left(\sum_i a_i \langle b_i v(0), f \rangle \right) \cdot \left(\sum_j a_j \langle b_j v(0), \varphi \rangle \right) = T(f) \cdot T(\varphi). \end{aligned} \quad \square$$

Proposition 4.2 *T is a morphism of $U_q\mathfrak{g}_0$ -modules.*

Proof. Denote by $U_q\mathfrak{g}\widehat{\otimes}V(0)$ the vector space of formal series of the form

$$\sum_i \xi_i \otimes v_i, \quad \xi_i \otimes v_i \in U_q\mathfrak{g} \otimes V(0).$$

Equip this vector space with a $U_q\mathfrak{g}$ -module structure as follows

$$\xi(\eta \otimes v) \stackrel{\text{def}}{=} \xi_{(3)} \cdot \eta \cdot S^{-1}(\xi_{(1)}) \otimes \xi_{(2)}v, \quad \xi, \eta \in U_q\mathfrak{g}, \quad v \in V(0).$$

Lemma 4.3 *The linear map*

$$V(0) \rightarrow U_q\mathfrak{g}\widehat{\otimes}V(0), \quad v \mapsto \sum_i a_i \otimes b_i v,$$

is a morphism of $U_q\mathfrak{g}$ -modules (here $\sum_i a_i \otimes b_i$ is the universal R -matrix).

Proof of the lemma. The map is well defined (it can be explained using the definition (3.1), (3.2) of the $U_q\mathfrak{g}$ -module $V(0)$ and the multiplicative formula (2.6) for the universal R -matrix). Let $\xi \in U_q\mathfrak{g}$, $v \in V(0)$. Then

$$\begin{aligned} \xi \left(\sum_i a_i \otimes b_i v \right) &= \sum_i \xi_{(3)} a_i S^{-1}(\xi_{(1)}) \otimes \xi_{(2)} b_i v = \sum_i \varepsilon(\xi_{(3)(2)}) \xi_{(3)(1)} a_i S^{-1}(\xi_{(1)}) \otimes \xi_{(2)} b_i v = \\ &= \sum_i \xi_{(3)(1)} a_i S^{-1}(\xi_{(1)}) \otimes \xi_{(2)} b_i \varepsilon(\xi_{(3)(2)}) v = \sum_i \xi_{(2)(2)} a_i S^{-1}(\xi_{(1)}) \otimes \xi_{(2)(1)} b_i \varepsilon(\xi_{(3)}) v. \end{aligned}$$

Let us make use of the property (2.5). We get

$$\begin{aligned} \xi \left(\sum_i a_i \otimes b_i v \right) &= \sum_i \xi_{(2)(2)} a_i S^{-1}(\xi_{(1)}) \otimes \xi_{(2)(1)} b_i \varepsilon(\xi_{(3)}) v = \\ &= \sum_i a_i \xi_{(2)(1)} S^{-1}(\xi_{(1)}) \otimes b_i \xi_{(2)(2)} \varepsilon(\xi_{(3)}) v = \sum_i a_i \xi_{(1)(2)} S^{-1}(\xi_{(1)(1)}) \otimes b_i \xi_{(2)(1)} \varepsilon(\xi_{(2)(2)}) v = \\ &= \sum_i a_i \varepsilon(\xi_{(1)}) \otimes b_i \xi_{(2)} v = \sum_i a_i \otimes b_i \varepsilon(\xi_{(1)}) \xi_{(2)} v = \sum_i a_i \otimes b_i \xi v. \end{aligned}$$

□

Lemma 4.4 *An element $\eta \otimes v \in U_q\mathfrak{g}\widehat{\otimes}V(0)$ is $U_q\mathfrak{g}_0$ -invariant iff for any $\xi \in U_q\mathfrak{g}_0$*

$$\xi_{(1)}\eta S(\xi_{(2)}) \otimes v = \eta \otimes S(\xi)v. \quad (4.2)$$

Proof of the lemma. Let $\eta \otimes v \in U_q\mathfrak{g}\widehat{\otimes}V(0)$ satisfies (4.2) for any $\xi \in U_q\mathfrak{g}_0$. Rewrite (4.2) for $\xi := S^{-1}(\zeta)$:

$$S^{-1}(\zeta_{(2)})\eta\zeta_{(1)} \otimes v = \eta \otimes \zeta v. \quad (4.3)$$

Using (4.3) one gets

$$\begin{aligned} \xi(\eta \otimes v) &= \xi_{(3)}\eta S^{-1}(\xi_{(1)}) \otimes \xi_{(2)}v = \xi_{(3)}S^{-1}(\xi_{(2)(2)})\eta\xi_{(2)(1)}S^{-1}(\xi_{(1)}) \otimes v = \\ &= \xi_{(2)(2)}S^{-1}(\xi_{(2)(1)})\eta\xi_{(1)(2)}S^{-1}(\xi_{(1)(1)}) \otimes v = \varepsilon(\xi_{(2)})\eta\varepsilon(\xi_{(1)}) \otimes v = \varepsilon(\xi)\eta \otimes v. \end{aligned}$$

Thus $\eta \otimes v$ is $U_q \mathfrak{g}_0$ -invariant.

Conversely, suppose that $\eta \otimes v$ is $U_q \mathfrak{g}_0$ -invariant. Let us prove (4.3) (obviously, it is equivalent to (4.2)).

$$S^{-1}(\xi_{(2)})\eta\xi_{(1)} \otimes v = S^{-1}(\xi_{(2)})\eta\varepsilon(\xi_{(1)(2)})\xi_{(1)(1)} \otimes v = S^{-1}(\xi_{(3)})\eta\varepsilon(\xi_{(2)})\xi_{(1)} \otimes v.$$

$U_q \mathfrak{g}_0$ -invariance of $\eta \otimes v$ implies

$$\varepsilon(\xi_{(2)})\eta \otimes v = \xi_{(2)(3)}\eta S^{-1}(\xi_{(2)(1)}) \otimes \xi_{(2)(2)}v.$$

Thus

$$\begin{aligned} S^{-1}(\xi_{(2)})\eta\xi_{(1)} \otimes v &= S^{-1}(\xi_{(3)})\xi_{(2)(3)}\eta S^{-1}(\xi_{(2)(1)})\xi_{(1)} \otimes \xi_{(2)(2)}v = \\ &= S^{-1}(\xi_{(3)(2)})\xi_{(3)(1)}\eta S^{-1}(\xi_{(1)(2)})\xi_{(1)(1)} \otimes \xi_{(2)}v = \varepsilon(\xi_{(3)})\eta\varepsilon(\xi_{(1)}) \otimes \xi_{(2)}v = \eta \otimes \xi v. \end{aligned}$$

□

Let us complete the proof of Proposition 4.2. By (3.1), (3.2) $v(0)$ is $U_q \mathfrak{g}_0$ -invariant. Due to Lemma 4.3 the element $\sum_i a_i \otimes b_i v(0) \in U_q \mathfrak{g} \widehat{\otimes} V(0)$ is $U_q \mathfrak{g}_0$ -invariant. By (4.2) we have: for $f \in \mathbb{C}[\mathfrak{g}_{-1}]_q$, $\xi \in U_q \mathfrak{g}_0$

$$\begin{aligned} T(\xi f) &= \sum_i a_i \langle b_i v(0), \xi f \rangle = \sum_i a_i \langle S(\xi) b_i v(0), f \rangle = \\ &= \sum_i \xi_{(1)} a_i S(\xi_{(2)}) \langle b_i v(0), f \rangle = \text{ad} \xi(T(f)). \end{aligned}$$

□

We have constructed the mapping $T : \mathbb{C}[\mathfrak{g}_{-1}]_q \rightarrow U_q \mathfrak{g}$ which is a morphism of $U_q \mathfrak{g}_0$ -module algebras. It turns out to be an embedding.

Proposition 4.5 *T is injective.*

Proof. Let $w_0 \in W$ be the maximal length element. We fix a reduced expression for w_0 :

$$w_0 = (1, 2)(3, 4)(2, 3)(1, 2)(3, 4)(2, 3) = s_1 s_3 s_2 s_1 s_3 s_2. \quad (4.4)$$

Obviously $w_0 = w'_0 s_2 s_1 s_3 s_2$ with w'_0 being the maximal length element in W_c . Describe explicitly the order in Δ_+ attached to the expression (4.4) (see section 2):

$$\beta_1 = \alpha_1, \quad \beta_2 = \alpha_3, \quad \beta_3 = \alpha_1 + \alpha_2 + \alpha_3,$$

$$\beta_4 = \alpha_2 + \alpha_3, \quad \beta_5 = \alpha_1 + \alpha_2, \quad \beta_6 = \alpha_2.$$

Thus $\beta_3, \beta_4, \beta_5, \beta_6$ are 'noncompact' roots (they belong to Δ_n), and β_1, β_2 are 'compact' roots (they belong to Δ_c). This implies $E_{\beta_1}, F_{\beta_1}, E_{\beta_2}, F_{\beta_2} \in U_q \mathfrak{g}_0$. Indeed, let us show this, for example, for E_{β_2} . Consider the $\mathbb{Z}_+$ -grading in U_q^+ given by

$$\deg E_i = \begin{cases} 1 & , \quad i = 2 \\ 0 & , \quad \text{otherwise} \end{cases}. \quad (4.5)$$

This grading can be described in another way:

$$(U_q^+)_j = \{\xi \in U_q^+ \mid \text{ad } K_0(\xi) = q^{2j} \xi\}. \quad (4.6)$$

Using (4.5) one shows that

$$(U_q^+)_0 = U_q^+ \cap U_q \mathfrak{g}_0. \quad (4.7)$$

It follows from the definition of E_{β_2} and K_0 that

$$\text{ad} K_0(E_{\beta_2}) = q^{\beta_2(H_0)} E_{\beta_2} = E_{\beta_2}.$$

Thus, according to (4.6) and (4.7) $E_{\beta_2} \in U_q \mathfrak{g}_0$.

It follows from the relations (3.1) and (3.2) that

$$\sum_i a_i \otimes b_i v(0) = \sum_j A_j \otimes B_j v(0),$$

where $\sum_i a_i \otimes b_i$ is the universal R -matrix,

$$\begin{aligned} \sum_j A_j \otimes B_j &\stackrel{\text{def}}{=} \exp_{q^2}((q^{-1} - q)E_{\beta_6} \otimes F_{\beta_6}) \cdot \exp_{q^2}((q^{-1} - q)E_{\beta_5} \otimes F_{\beta_5}) \times \\ &\times \exp_{q^2}((q^{-1} - q)E_{\beta_4} \otimes F_{\beta_4}) \cdot \exp_{q^2}((q^{-1} - q)E_{\beta_3} \otimes F_{\beta_3}). \end{aligned} \quad (4.8)$$

It is clear that

$$\sum_j A_j \otimes B_j = \sum_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4} a_{k_1, \dots, k_4} E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4} \otimes F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4}, \quad (4.9)$$

where all $a_{k_1, \dots, k_4}$ are nonzero elements of $\mathbb{C}(q^{1/4})$. Thus we get the formula

$$T(f) = \sum_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4} a_{k_1, \dots, k_4} E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4} \langle F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4} v(0), f \rangle. \quad (4.10)$$

To complete the proof of Proposition 4.5 it is sufficient to prove that the vectors

$$\{E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$$

are linearly independent, and the vectors

$$\{F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4} v(0)\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$$

constitute a basis in $V(0)$ (the latter statement implies also the well definiteness of T !). For this purpose we need the following Theorem [4, page 14]:

Theorem 4.6 *i) The monomials*

$$\{F_{\beta_6}^{k_1} F_{\beta_5}^{k_2} \dots F_{\beta_1}^{k_6} K_1^{m_1} K_2^{m_2} K_3^{m_3} E_{\beta_1}^{l_1} E_{\beta_2}^{l_2} \dots E_{\beta_6}^{l_6}\}_{(k_1, \dots, k_6) \in \mathbb{Z}_+^6, (m_1, m_2, m_3) \in \mathbb{Z}^3, (l_1, \dots, l_6) \in \mathbb{Z}_+^6}$$

constitute a basis in $U_q \mathfrak{g}$;

ii) for $i < j$ one has:

$$E_{\beta_i} E_{\beta_j} - q^{(\beta_i | \beta_j)} E_{\beta_j} E_{\beta_i} = \sum_{(k_1, \dots, k_6) \in \mathbb{Z}_+^6} a_{k_1, \dots, k_6} E_{\beta_1}^{k_1} E_{\beta_2}^{k_2} \dots E_{\beta_6}^{k_6},$$

where $a_{k_1, \dots, k_6} \neq 0$ only when $k_s = 0$ for $s \leq i$ or $s \geq j$.

It is not hard to prove that this Theorem implies linear independence of the vectors $\{E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$. Let us prove that the vectors $\{F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4} v(0)\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ constitute a basis in $V(0)$.

Obviously, $V(0)$ is the linear span of $\{F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4} v(0)\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$. Indeed, this follows from statement i) of the Theorem, from the observations that the map $U_q \mathfrak{g} \rightarrow V(0)$, $\xi \mapsto \xi v(0)$ is surjective, and from the relations

$$\begin{aligned} E_{\beta_i} v(0) &= 0, \quad i = \overline{1, 6}, \\ K_i v(0) &= v(0), \quad i = \overline{1, 3}, \\ F_{\beta_1} v(0) &= F_{\beta_2} v(0) = 0. \end{aligned}$$

What remains is to prove that $\{F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4} v(0)\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ are linearly independent. We prove this statement using its correctness for $q = 1$.

Let A be the ring $\mathbb{C}[q, q^{-1}]$. Consider the A -algebra $U_{q,A}^-$ generated by $\{F_i^A\}_{i=\overline{1,3}}$ which satisfy the same relations as $\{F_i\}_{i=\overline{1,3}}$. Evidently, as $\mathbb{C}(q^{1/4})$ -algebras

$$U_q^- \simeq \mathbb{C}(q^{1/4}) \bigotimes_A U_{q,A}^-. \quad (4.11)$$

Let $\{F_{\beta_i}^A\}_{i=\overline{1,6}}$ be the elements of $U_{q,A}^-$ derived from $\{F_{\beta_i}\}_{i=\overline{1,6}}$ via substitution $F_i \leftrightarrow F_i^A$. Consider the $U_{q,A}^-$ -module $V(0)^A$ given by the generator $v(0)^A$ and the relations

$$F_i^A v(0)^A = 0, \quad i \neq 2.$$

Similarly to (4.11)

$$V(0) \simeq \mathbb{C}(q^{1/4}) \bigotimes_A V(0)_q^A \quad (4.12)$$

as $\mathbb{C}(q^{1/4})$ -modules.

There is an evident homomorphism of $\mathbb{C}$ -algebras

$$J : U_{q,A}^- \rightarrow U_{q,A}^- / (q - 1) \cdot U_{q,A}^- \simeq U^-,$$

where U^- is the subalgebra in the classical universal enveloping algebra $U \mathfrak{g}$ generated by $\{J(F_i^A)\}_{i=\overline{1,3}}$. Denote by $\widetilde{V(0)}$ a U^- -module given by the generator $\widetilde{v(0)}$ and the relations

$$J(F_i^A) \widetilde{v(0)} = 0, \quad i \neq 2.$$

It is clear that the map $v(0)^A \mapsto \widetilde{v(0)}$ can be extended up to a $\mathbb{C}$ -linear map

$$J_0 : V(0)^A \rightarrow \widetilde{V(0)}$$

such that for any $\xi \in U_{q,A}^-$

$$J(\xi) \widetilde{v(0)} = J_0(\xi v(0)^A). \quad (4.13)$$

It is well known that the vectors $\{J(F_{\beta_6}^A)^{k_1} \dots J(F_{\beta_3}^A)^{k_4} \widetilde{v(0)}\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ constitute a basis in the $\mathbb{C}$ -module $\widetilde{V(0)}$. Thus by (4.13) the vectors $\{(F_{\beta_6}^A)^{k_1} \dots (F_{\beta_3}^A)^{k_4} v(0)^A\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ are linearly independent. Due to (4.12) $\{F_{\beta_6}^{k_1} \dots F_{\beta_3}^{k_4} v(0)\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ are linearly independent.

We have completed the proof of Proposition 4.5. $\square$

The following statement is a direct consequence of Propositions 4.1, 4.5 and formula (4.10).

Corollary 4.7 *Linear span of $\{E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ is a $\text{ad } U_q \mathfrak{g}_0$ -invariant subalgebra in U_q^+ . It coincides with $T(\mathbb{C}[\mathfrak{g}_{-1}]_q)$.*

REMARK 3. The fact that linear span of $\{E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ is a subalgebra in U_q^+ easily follows from the statement ii) of the Theorem 4.6.

The main statement of this section is

Proposition 4.8

$$T(\mathbb{C}[\mathfrak{g}_{-1}]_q) = \mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}.$$

Proof. Let us start with

Lemma 4.9 *For some $c \in \mathbb{C}(q^{1/4})$*

$$E_{\beta_6} = c \cdot E_2.$$

Proof of the Lemma. The elements E_{β_6} , E_2 of the U_q^0 -module U_q^+ are weight vectors of the weight $\alpha_2 \in \mathfrak{h}^*$. But the subspace in U_q^+ of weight vectors of that weight is 1-dimensional (this follows from linear independence of the weights $\alpha_1, \alpha_2, \alpha_3 \in \mathfrak{h}^*$ of the generators E_1, E_2, E_3). $\square$

Remind that we have equipped U_q^+ with a $\mathbb{Z}_+$ -grading (see (4.6)). Obviously,

$$\deg(E_{\beta_3}) = \deg(E_{\beta_4}) = \deg(E_{\beta_5}) = \deg(E_{\beta_6}) = 1.$$

It follows from (2.3), (4.6) that endomorphisms from $\text{ad } U_q \mathfrak{g}_0$ preserve this grading: for $\xi \in U_q \mathfrak{g}_0$ and $\eta \in U_q^+$

$$\deg(\text{ad} \xi(\eta)) = \deg(\eta)$$

provided $\text{ad} \xi(\eta) \in U_q^+$. Using this observation and Corollary 4.7 we get

$$\text{ad } U_q \mathfrak{g}_0(E_{\beta_6}) \subseteq \text{linear span}\{E_{\beta_3}, E_{\beta_4}, E_{\beta_5}, E_{\beta_6}\}. \quad (4.14)$$

Actually the spaces in the both sides of (4.14) coincide: dimension of $\text{ad } U_q \mathfrak{g}_0(E_{\beta_6})$ should be equal to 4 just as in the classical case $q = 1$. Thus, by Lemma 4.9 and by the definition of the algebra $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}$

$$\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}} = \langle E_{\beta_3}, E_{\beta_4}, E_{\beta_5}, E_{\beta_6} \rangle.$$

What remains is to use Corollary 4.7. $\square$

$$\mathbf{4.2} \quad \mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}} = \mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$$

In this subsection we use notation of the previous one.

Proposition 4.10

$$\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}} \subseteq \mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}.$$

Proof. By Corollary 4.7 and Proposition 4.8 the linear span of $\{E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}\}_{(k_1, \dots, k_4) \in \mathbb{Z}_+^4}$ coincides with $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{I}}$. Thus, due to the definition (3.6) of $\mathbb{C}[\mathfrak{g}_{-1}]_q^{\text{II}}$ it is sufficient to prove that

$$T_{w'_0}^{-1}(E_{\beta_k}) \in U_q^+, \quad k = \overline{3, 6}.$$

Let us prove this, for example, for E_{β_6} . By definition

$$E_{\beta_6} = T_1 T_3 T_2 T_1 T_3(E_2) = T_{w'_0} T_2 T_1 T_3(E_2).$$

One gets

$$T_{w'_0}^{-1}(E_{\beta_6}) = T_2 T_1 T_3(E_2).$$

It remains to make use of the following well known fact.

Lemma 4.11 *If $w(\alpha_i) \in \Delta_+$ for some $i = \overline{1, 3}$ and $w \in W$ then $T_w(E_i) \in U_q^+$.*

□

Now we are ready to prove

Proposition 4.12

$$\mathbb{C}[\mathfrak{g}_{-1}]_q^I = \mathbb{C}[\mathfrak{g}_{-1}]_q^{II}.$$

Proof. Let $\sum a_{k_1, \dots, k_6} E_{\beta_6}^{k_1} \dots E_{\beta_1}^{k_6} \in \mathbb{C}[\mathfrak{g}_{-1}]_q^{II}$, i.e.

$$T_{w'_0}^{-1} \left(\sum a_{k_1, \dots, k_6} E_{\beta_6}^{k_1} \dots E_{\beta_1}^{k_6} \right) = \sum a_{k_1, \dots, k_6} T_{w'_0}^{-1} (E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}) T_{w'_0}^{-1} (E_{\beta_2}^{k_5} \cdot E_{\beta_1}^{k_6}) \in U_q^+. \quad (4.15)$$

By Proposition 4.10

$$T_{w'_0}^{-1} (E_{\beta_6}^{k_1} \dots E_{\beta_3}^{k_4}) \in U_q^+. \quad (4.16)$$

Lemma 4.13 $T_{w'_0}^{-1} (E_{\beta_2}) \in U_q^{\leq 0}$, $T_{w'_0}^{-1} (E_{\beta_1}) \in U_q^{\leq 0}$.

Proof of the Lemma. Suppose that β_k is a 'compact' root ($\beta_k = \beta_1$ or $\beta_k = \beta_2$). Let $s_{i_1} s_{i_2} \dots s_{i_M}$ be a reduced expression of w'_0 (of course, in the case we consider $M = 2$ and there are only two different reduced expression for w'_0). One has

$$\beta_k = s_{i_1} s_{i_2} \dots s_{i_{k-1}}(\alpha_k), \quad E_{\beta_k} = T_{i_1} T_{i_2} \dots T_{i_{k-1}}(E_k).$$

Since $T_{w'_0} = T_{i_1} T_{i_2} \dots T_{i_M}$ we get

$$T_{w'_0}^{-1}(E_{\beta_k}) = T_{i_M}^{-1} T_{i_{M-1}}^{-1} \dots T_{i_k}^{-1}(E_k).$$

So we have to prove that

$$T_{i_M}^{-1} T_{i_{M-1}}^{-1} \dots T_{i_k}^{-1}(E_k) \in U_q^{\leq 0}. \quad (4.17)$$

Consider the antiautomorphism τ of the algebra $U_q \mathfrak{g}$ given by

$$\tau(K_i) = K_i^{-1}, \quad \tau(E_i) = E_i, \quad \tau(F_i) = F_i.$$

It is not hard to prove that

$$\tau \circ T_i = T_i^{-1} \circ \tau.$$

Thus the inclusion

$$T_{i_M} T_{i_{M-1}} \dots T_{i_k}(E_k) \in U_q^{\leq 0}. \quad (4.18)$$

is equivalent to (4.17). One has $T_{i_k}(E_k) = -F_k K_k$. Therefore (4.18) is equivalent to

$$T_{i_M} T_{i_{M-1}} \dots T_{i_{k+1}}(F_k) \cdot T_{i_M} T_{i_{M-1}} \dots T_{i_{k+1}}(K_k) \in U_q^{\leq 0}$$

and thus to

$$T_{i_M} T_{i_{M-1}} \dots T_{i_{k+1}}(F_k) \in U_q^-. \quad (4.19)$$

Applying the antiautomorphism k (see section 2) to both hand sides of (4.19) we get the equivalent inclusion

$$T_{i_M} T_{i_{M-1}} \dots T_{i_{k+1}}(E_k) \in U_q^+. \quad (4.20)$$

But (4.20) is a direct consequence of Lemma 4.11. $\square$

The following result is well known.

Lemma 4.14 *The multiplication in $U_q \mathfrak{g}$ induces the isomorphism of vector spaces*

$$U_q^+ \bigotimes U_q^{\leq 0} \rightarrow U_q \mathfrak{g}.$$

It follows from (4.16) and Lemmas 4.13, 4.14 that (4.15) holds iff $a_{k_1, \dots, k_6} = 0$ for $k_5 \neq 0$ or $k_6 \neq 0$. We have completed the proof of Proposition 4.12. $\square$

Comparing results of the two previous sections we get:
the algebras $\mathbb{C}[\mathfrak{g}_{-1}]_q$, $\mathbb{C}[\mathfrak{g}_{-1}]_q^I$, and $\mathbb{C}[\mathfrak{g}_{-1}]_q^{II}$ are isomorphic to each other as $U_q \mathfrak{g}_0$ -module algebras.

References

- [1] V. G. Drinfeld. *Quantum groups*. In: Proceedings of the International Congress of Mathematicians, Berkeley, 1986, A. M. Gleason (ed.), 1987, American Mathematical Society, Providence RI, 798 – 820.
- [2] H. P. Jakobsen. *Quantized Hermitian Symmetric Spaces*. In: Lie theory and its applications in physics (Clausthal, 1995), 105 – 116.
- [3] A. Kamita, Y. Morita, and T. Tanisaki, *Quantum deformations of certain prehomogeneous spaces I*, Hiroshima Math. J., **28** (1998), 527 – 540.
- [4] I. Damiani, C. de Concini, *Quantum groups and Poisson groups*, In: Proceedings of the session of the European School of Group Theory, Trento, Italy, 1993, V. Baldoni (ed.) et al., 'Representations of Lie groups and quantum groups', 1994, Longman Scientific & Technical, Pitman Res. Notes Math., ser. 311, 1 – 45.
- [5] H. Rubenthaler, *Les paires duales dans les algèbres de Lie réductives*, Asterisque, **219** (1994).
- [6] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035, 1999.
- [7] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: the Bergman kernel*, math.QA/9909036, 1999.
- [8] S. Sinel'shchikov and L. Vaksman, *Hidden symmetry of the differential calculus on the quantum matrix space*, J. Phys. A. **30** (1997), 23 – 26 (see also this volume).
- [9] S. Sinel'shchikov and L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, Math. Phys., Anal., and Geom., **1** (1998), 75 – 100; q-alg/9703005, 1997.

HIDDEN SYMMETRY OF SOME ALGEBRAS OF q -DIFFERENTIAL OPERATORS

D. Shklyarov, S. Sinel'shchikov* L. Vaksman[†]

Institute for Low Temperature Physics & Engineering
47 Lenin Avenue, 61103 Kharkov, Ukraine

1 Introduction

Let us explain the meaning of the words "q-differential operators" and "hidden symmetry". Let $\mathbb{C}[z]_q$ be the algebra of polynomials in z over the field of rational functions $\mathbb{C}(q^{1/2})$ (we assume this field to be the ground field throughout the paper). We denote by $\Lambda^1(\mathbb{C})_q$ the $\mathbb{C}[z]_q$ -bimodule with the generator dz such that

$$z \cdot dz = q^{-2} dz \cdot z.$$

Let d be the linear map $\mathbb{C}[z]_q \rightarrow \Lambda^1(\mathbb{C})_q$ given by the two conditions:

$$d : z \mapsto dz,$$

$$d(f_1(z)f_2(z)) = d(f_1(z))f_2(z) + f_1(z)d(f_2(z)).$$

(The later condition is just the Leibniz rule). The bimodule $\Lambda^1(\mathbb{C})_q$ (together with the map d) is a well known first order differential calculus over the algebra $\mathbb{C}[z]_q$. The differential d allows one to introduce an operator of "partial derivative" $\frac{d}{dz}$ in $\mathbb{C}[z]_q$:

$$df = dz \cdot \frac{df}{dz}.$$

Let us introduce also the notation $\widehat{z}$ for the operator in $\mathbb{C}[z]_q$ of multiplication by z :

$$\widehat{z} : f(z) \mapsto zf(z).$$

Let $D(\mathbb{C})_q$ be the subalgebra in the algebra $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[z]_q)$ (of all endomorphisms of the linear space $\mathbb{C}[z]_q$) containing 1 and generated by $\frac{d}{dz}$, $\widehat{z}$. It is easy to check that

$$\frac{d}{dz} \cdot \widehat{z} = q^{-2} \widehat{z} \cdot \frac{d}{dz} + 1.$$

Thus the algebra $D(\mathbb{C})_q$ is an analogue of the Weyl algebra $A_1(\mathbb{C})$.

This lecture has been delivered at the NATO Advanced Research Workshop 'Non-commutative Structures in Mathematics and Physics', Kiev, September 2000; published in Noncommutative Structures in Mathematical Physics, S. Duplij and J. Wess (eds), Kluwer AP, Netherlands, 2001, 309 – 320.

*email: sinelshchikov@ilt.kharkov.ua

[†]email: vaksman@ilt.kharkov.ua

Let $\lambda \in \mathbb{C}(q^{1/2})$. One checks that the map

$$\widehat{z} \mapsto \lambda \cdot \widehat{z}, \quad \frac{d}{dz} \mapsto \lambda^{-1} \cdot \frac{d}{dz}$$

is extendable up to an automorphism of the algebra $D(\mathbb{C})_q$. Such automorphisms are "evident" symmetries of $D(\mathbb{C})_q$. It turn out that they belong to a wider set of symmetries of $D(\mathbb{C})_q$. This set does not consists of automorphisms only. Let us turn to precise formulations.

To start with, we recall the definition of the quantum universal enveloping algebra $U_q \mathfrak{sl}_2$ [1]. It is

i) the algebra given by the generators E, F, K, K^{-1} , and the relations

$$KK^{-1} = K^{-1}K = 1, \quad KE = q^2EK, \quad KF = q^{-2}FK,$$

$$EF - FE = \frac{K - K^{-1}}{q - q^{-1}};$$

ii) the Hopf algebra: the comultiplication Δ , the antipode S , and the counit ε are determined by

$$\Delta(E) = E \otimes 1 + K \otimes E, \quad \Delta(F) = F \otimes K^{-1} + 1 \otimes F, \quad \Delta(K) = K \otimes K,$$

$$S(E) = -K^{-1}E, \quad S(F) = -FK, \quad S(K) = K^{-1},$$

$$\varepsilon(E) = \varepsilon(F) = 0, \quad \varepsilon(K) = 1.$$

There is a well known structure of $U_q \mathfrak{sl}_2$ -module in the space $\mathbb{C}[z]_q$. Let us describe it explicitly:

$$E : f(z) \mapsto -q^{1/2}z^2 \frac{f(z) - f(q^2z)}{z - q^2z},$$

$$F : f(z) \mapsto q^{1/2} \frac{f(z) - f(q^{-2}z)}{z - q^{-2}z},$$

$$K^{\pm 1} : f(z) \mapsto f(q^{\pm 2}z).$$

It can be checked that $\mathbb{C}[z]_q$ is a $U_q \mathfrak{sl}_2$ -module algebra, i.e. for any $\xi \in U_q \mathfrak{sl}_2$, $f_1, f_2 \in \mathbb{C}[z]_q$

$$\xi(1) = \varepsilon(\xi) \cdot 1, \tag{1.1}$$

$$\xi(f_1 f_2) = \sum_j \xi'_j(f_1) \xi''_j(f_2), \tag{1.2}$$

with $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$.

REMARK. This observation is an analogue of the following one. The group $SL_2(\mathbb{C})$ acts on $\mathbb{CP}^1$ via the fractional-linear transformations. Thus the universal enveloping algebra $U \mathfrak{sl}_2$ acts via differential operators in the space of holomorphic functions on the open cell $\mathbb{C} \subset \mathbb{CP}^1$.

Let V be a $U_q \mathfrak{sl}_2$ -module. Then the algebra $\text{End}(V)$ admits a "canonical" structure of $U_q \mathfrak{sl}_2$ -module: for $\xi \in U_q \mathfrak{sl}_2$, $T \in \text{End}(V)$

$$\xi(T) = \sum_j \xi'_j \cdot T \cdot S(\xi''_j), \tag{1.3}$$

where $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$, S is the antipode, and the elements in the right-hand side are multiplied within the algebra $\text{End}(V)$. It is well known that this action of $U_q \mathfrak{sl}_2$ in $\text{End}(V)$ makes $\text{End}(V)$ into a $U_q \mathfrak{sl}_2$ -module algebra (i.e. for $\xi \in U_q \mathfrak{sl}_2$, $T_1, T_2 \in \text{End}(V)$ (1.1), (1.2) hold with f_1, f_2 being replaced by T_1, T_2 , respectively).

The objects considered above are the simplest among ones we deal with in the present paper. In this simplest case our main result can be formulated as follows: the algebra $D(\mathbb{C})_q$ is a $U_q \mathfrak{sl}_2$ -module subalgebra in the $U_q \mathfrak{sl}_2$ -module algebra $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[z]_q)$ (where the $U_q \mathfrak{sl}_2$ -action is given by (1.3)). This $U_q \mathfrak{sl}_2$ -module structure in the algebra $D(\mathbb{C})_q$ is what we call "hidden symmetry" of $D(\mathbb{C})_q$.

REMARK. In the setting of the previous Remark the analogous fact is evident: for $\xi \in \mathfrak{sl}_2$ the action (1.3) is just the commutator of the differential operators ξ and T in the space of holomorphic functions on $\mathbb{C}$. The commutator is again a differential operator.

We can describe the $U_q \mathfrak{sl}_2$ -action in $D(\mathbb{C})_q$ explicitly:

$$E(\widehat{z}) = -q^{1/2} \widehat{z}^2, \quad F(\widehat{z}) = q^{1/2}, \quad K^{\pm 1}(\widehat{z}) = q^{\pm 2} \widehat{z},$$

$$E\left(\frac{d}{dz}\right) = q^{-3/2}(q^{-2} + 1) \widehat{z} \frac{d}{dz}, \quad F\left(\frac{d}{dz}\right) = 0, \quad K^{\pm 1}\left(\frac{d}{dz}\right) = q^{\mp 2} \frac{d}{dz}.$$

(The action of $U_q \mathfrak{sl}_2$ on an arbitrary element of $D(\mathbb{C})_q$ can be produced via the rule (1.2).)

The paper is organized as follows.

In Section 2 we recall one definitions of the quantum universal enveloping algebra $U_q \mathfrak{sl}_N$, a $U_q \mathfrak{sl}_N$ -module algebra of holomorphic polynomials on a quantum space of $m \times n$ matrices ($N = m + n$), and a well known first order differential calculus over this algebra (in this Introduction the case $m = n = 1$ was considered). Then we introduce an algebra of q -differential operators and formulate a main theorem concerning a hidden symmetry of this algebra.

Section 3 contains a sketch of the proof of the main theorem.

In Section 4 we discuss briefly q -analogues of the notions of a holomorphic vector bundle and a differential operator in sections of holomorphic vector bundles. We formulate an analogue of our main theorem for such differential operators.

Section 'Concluding notes' discusses one of possible generalizations of our results, specifically, the case when the matrix space is replaced by an arbitrary prehomogeneous vector space of commutative parabolic type [7].

Appendix deals with q -analogues of constant coefficient differential operators in functions and in sections of holomorphic vector bundles.

We take this opportunity to thank Prof. H. P. Jakobsen and Prof. T. Tanisaki who attracted our attention to other approaches to the notion of quantum differential operators.

This research was partially supported by Award No.UM1-2091 of the U.S. Civilian Research and Development Foundation.

2 The main theorem

Throughout the paper (except for the section 'Concluding notes') we suppose that $\mathfrak{g} = \mathfrak{sl}_N$, $\mathfrak{k} = \mathfrak{sl}_n \times \mathfrak{sl}_m$, $N = n + m$, and $\mathfrak{p}^-$ is the space of complex $m \times n$ matrices.

The algebra $\mathbb{C}[\mathfrak{p}^-]_q$ is the unital algebra given by its generators z_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, and the following relations

$$z_a^\alpha z_b^\beta = \begin{cases} q z_b^\beta z_a^\alpha & , \quad a = b \ \& \ \alpha < \beta \quad \text{or} \quad a < b \ \& \ \alpha = \beta \\ z_b^\beta z_a^\alpha & , \quad a < b \ \& \ \alpha > \beta \\ z_b^\beta z_a^\alpha + (q - q^{-1}) z_a^\beta z_b^\alpha & , \quad a < b \ \& \ \alpha < \beta \end{cases}, \quad (2.1)$$

Let (a_{ij}) be the Cartan matrix for $\mathfrak{g}$. The Hopf algebra $U_q \mathfrak{g}$ is determined by the generators E_i, F_i, K_i, K_i^{-1} , $i = 1, \dots, N-1$, and the relations

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1, & K_i E_j &= q^{a_{ij}} E_j K_i, \\ K_i F_j &= q^{-a_{ij}} F_j K_i, & E_i F_j - F_j E_i &= \delta_{ij} (K_i - K_i^{-1}) / (q - q^{-1}) \end{aligned}$$

$$\begin{aligned} E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0, & |i - j| &= 1 \\ F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0, & |i - j| &= 1 \\ [E_i, E_j] &= [F_i, F_j] = 0, & |i - j| &\neq 1. \end{aligned} \quad (2.2)$$

The comultiplication Δ , the antipode S , and the counit ε are determined by

$$\Delta(E_i) = E_i \otimes 1 + K_i \otimes E_i, \quad \Delta(F_i) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \quad \Delta(K_i) = K_i \otimes K_i, \quad (2.3)$$

$$\begin{aligned} S(E_i) &= -K_i^{-1} E_i, & S(F_i) &= -F_i K_i, & S(K_i) &= K_i^{-1}, \\ \varepsilon(E_i) &= \varepsilon(F_i) = 0, & \varepsilon(K_i) &= 1. \end{aligned} \quad (2.4)$$

Denote by $U_q \mathfrak{k}$ the Hopf subalgebra in $U_q \mathfrak{g}$ generated by E_j, F_j, K_i, K_i^{-1} , $i, j = 1, \dots, N-1$, $j \neq n$.

The algebra $\mathbb{C}[\mathfrak{p}^-]_q$ possesses the well known structure of $U_q \mathfrak{k}$ -module algebra:

$$K_n z_a^\alpha = \begin{cases} q^2 z_a^\alpha & , \quad a = n \ \& \ \alpha = m \\ q z_a^\alpha & , \quad a = n \ \& \ \alpha \neq m \quad \text{or} \quad a \neq n \ \& \ \alpha = m \\ z_a^\alpha & , \quad \text{otherwise} \end{cases}, \quad (2.5)$$

and with $k \neq n$

$$K_k z_a^\alpha = \begin{cases} q z_a^\alpha & , \quad k < n \ \& \ a = k \quad \text{or} \quad k > n \ \& \ \alpha = N - k \\ q^{-1} z_a^\alpha & , \quad k < n \ \& \ a = k + 1 \quad \text{or} \quad k > n \ \& \ \alpha = N - k + 1 \\ z_a^\alpha & , \quad \text{otherwise} \end{cases}, \quad (2.6)$$

$$F_k z_a^\alpha = q^{1/2} \cdot \begin{cases} z_{a+1}^\alpha & , \quad k < n \ \& \ a = k \\ z_{a+1}^{\alpha+1} & , \quad k > n \ \& \ \alpha = N - k \\ 0 & , \quad \text{otherwise} \end{cases}, \quad (2.7)$$

$$E_k z_a^\alpha = q^{-1/2} \cdot \begin{cases} z_{a-1}^\alpha & , \quad k < n \ \& \ a = k + 1 \\ z_{a-1}^{\alpha-1} & , \quad k > n \ \& \ \alpha = N - k + 1 \\ 0 & , \quad \text{otherwise} \end{cases}. \quad (2.8)$$

This $U_q\mathfrak{k}$ -module algebra structure in $\mathbb{C}[\mathfrak{p}^-]_q$ can be extended up to a $U_q\mathfrak{g}$ -module algebra structure as follows (see [8]):

$$F_n z_a^\alpha = q^{1/2} \cdot \begin{cases} 1 & , \quad a = n \ \& \ \alpha = m \\ 0 & , \quad \text{otherwise} \end{cases} , \quad (2.9)$$

$$E_n z_a^\alpha = -q^{1/2} \cdot \begin{cases} q^{-1} z_a^m z_n^\alpha & , \quad a \neq n \ \& \ \alpha \neq m \\ (z_n^m)^2 & , \quad a = n \ \& \ \alpha = m \\ z_n^m z_a^\alpha & , \quad \text{otherwise} \end{cases} . \quad (2.10)$$

REMARKS. i) In the classical case the corresponding action of $U\mathfrak{g}$ in the space of holomorphic functions on $\mathfrak{p}^-$ can be produced via an embedding $\mathfrak{p}^-$ into the Grassmanian $Gr_{m,N}$ as an open cell (we describe a q-analogue of the embedding in [8]).

Now let us recall a definition of a well known first order differential calculus over $\mathbb{C}[\mathfrak{p}^-]_q$. Let $\Lambda^1(\mathfrak{p}^-)_q$ be the $\mathbb{C}[\mathfrak{p}^-]_q$ -bimodule given by its generators dz_a^α , $a = 1, \dots, n$, $\alpha = 1, \dots, m$, and the relations

$$z_b^\beta dz_a^\alpha = \sum_{\alpha', \beta'=1}^m \sum_{a', b'=1}^n R_{\beta\alpha}^{\beta'\alpha'} R_{ba}^{b'a'} dz_{a'}^{\alpha'} \cdot z_{b'}^{\beta'} , \quad (2.11)$$

with

$$R_{ba}^{b'a'} = \begin{cases} q^{-1} & , \quad a = b = a' = b' \\ 1 & , \quad a \neq b \ \& \ a = a' \ \& \ b = b' \\ q^{-1} - q & , \quad a < b \ \& \ a = b' \ \& \ b = a' \\ 0 & , \quad \text{otherwise} \end{cases} . \quad (2.12)$$

The map $d : z_a^\alpha \mapsto dz_a^\alpha$ can be extended up to a linear operator $d : \mathbb{C}[\mathfrak{p}^-]_q \rightarrow \Lambda^1(\mathfrak{p}^-)_q$ satisfying the Leibniz rule. It was noted for the first time in [9], that there exists a unique structure of a $U_q\mathfrak{g}$ -module $\mathbb{C}[\mathfrak{p}^-]_q$ -bimodule in $\Lambda^1(\mathfrak{p}^-)_q$ such that the map d is a morphism of $U_q\mathfrak{g}$ -modules. The pair $(\Lambda^1(\mathfrak{p}^-)_q, d)$ is the first order differential calculus over $\mathbb{C}[\mathfrak{p}^-]_q$.

Let us introduce an algebra $D(\mathfrak{p}^-)_q$ of q-differential operators on $\mathfrak{p}^-$. Define the linear operators $\frac{\partial}{\partial z_a^\alpha}$ in $\mathbb{C}[\mathfrak{p}^-]_q$ via the differential d :

$$df = \sum_{a=1}^n \sum_{\alpha=1}^m dz_a^\alpha \cdot \frac{\partial f}{\partial z_a^\alpha}, \quad f \in \mathbb{C}[\mathfrak{p}^-]_q,$$

and the operators $\widehat{z}_a^\alpha$ by

$$\widehat{z}_a^\alpha f = z_a^\alpha \cdot f, \quad f \in \mathbb{C}[\mathfrak{p}^-]_q.$$

Then $D(\mathfrak{p}^-)_q$ is the unital subalgebra in $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$ generated by the operators $\frac{\partial}{\partial z_a^\alpha}$, $\widehat{z}_a^\alpha$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$.

To start with, we describe $D(\mathfrak{p}^-)_q$ in terms of generators and relations.

Proposition 2.1 *The complete list of relations between the generators $\widehat{z}_a^\alpha$, $\frac{\partial}{\partial z_a^\alpha}$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$, of $D(\mathfrak{p}^-)_q$ is as follows*

$$\widehat{z}_a^\alpha \widehat{z}_b^\beta = \begin{cases} q \widehat{z}_b^\beta \widehat{z}_a^\alpha & , \quad a = b \ \& \ \alpha < \beta \quad \text{or} \quad a < b \ \& \ \alpha = \beta \\ \widehat{z}_b^\beta \widehat{z}_a^\alpha & , \quad a < b \ \& \ \alpha > \beta \\ \widehat{z}_b^\beta \widehat{z}_a^\alpha + (q - q^{-1}) \widehat{z}_a^\beta \widehat{z}_b^\alpha & , \quad a < b \ \& \ \alpha < \beta \end{cases} , \quad (2.13)$$

$$\frac{\partial}{\partial z_b^\beta} \frac{\partial}{\partial z_a^\alpha} = \begin{cases} q \frac{\partial}{\partial z_a^\alpha} \frac{\partial}{\partial z_b^\beta} & , \quad a = b \text{ \& } \alpha < \beta \quad \text{or} \quad a < b \text{ \& } \alpha = \beta \\ \frac{\partial}{\partial z_a^\alpha} \frac{\partial}{\partial z_b^\beta} & , \quad a < b \text{ \& } \alpha > \beta \\ \frac{\partial}{\partial z_a^\alpha} \frac{\partial}{\partial z_b^\beta} + (q - q^{-1}) \frac{\partial}{\partial z_a^\beta} \frac{\partial}{\partial z_b^\alpha} & , \quad a < b \text{ \& } \alpha < \beta \end{cases}, \quad (2.14)$$

$$\frac{\partial}{\partial z_a^\alpha} \widehat{z}_b^\beta = \sum_{a', b'=1}^n \sum_{\alpha', \beta'=1}^m R_{ba'}^{b'a} R_{\beta\alpha'}^{\beta'\alpha} \widehat{z}_{b'}^{\beta'} \frac{\partial}{\partial z_{a'}^{\alpha'}} + \delta_{ab} \delta^{\alpha\beta}, \quad (2.15)$$

with $\delta_{ab}, \delta^{\alpha\beta}$ being the Kronecker symbols, and $R_{ba'}^{b'a}$ given by (2.12).

Using the $U_q \mathfrak{g}$ -module structure in $\mathbb{C}[\mathfrak{p}^-]_q$, we can define the structure of $U_q \mathfrak{g}$ -module algebra in $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$ via (1.3) with $\xi \in U_q \mathfrak{g}$, $T \in \text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$.

Our main result is

Theorem 2.2 *i) The algebra $D(\mathfrak{p}^-)_q$ is a $U_q \mathfrak{g}$ -module subalgebra in the $U_q \mathfrak{g}$ -module algebra $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$.*

ii) The $U_q \mathfrak{g}$ -module structure in $D(\mathfrak{p}^-)_q$ is described explicitly as follows:

$U_q \mathfrak{g}$ acts on the generators $\widehat{z}_a^\alpha$ via formulae (2.5)-(2.8) (where z_a^α should be replaced by $\widehat{z}_a^\alpha$); for the generators $\frac{\partial}{\partial z_a^\alpha}$ the formulae are

$$K_n \frac{\partial}{\partial z_a^\alpha} = \begin{cases} q^{-2} \frac{\partial}{\partial z_a^\alpha} & , \quad a = n \text{ \& } \alpha = m \\ q^{-1} \frac{\partial}{\partial z_a^\alpha} & , \quad a = n \text{ \& } \alpha \neq m \quad \text{or} \quad a \neq n \text{ \& } \alpha = m \\ \frac{\partial}{\partial z_a^\alpha} & , \quad \text{otherwise} \end{cases}, \quad (2.16)$$

$$F_n \frac{\partial}{\partial z_a^\alpha} = 0 \quad a = 1, \dots, n, \quad \alpha = 1, \dots, m, \quad (2.17)$$

$$E_n \frac{\partial}{\partial z_a^\alpha} = q^{-3/2}.$$

$$\cdot \begin{cases} \sum_{b=1}^n \widehat{z}_b^m \frac{\partial}{\partial z_b^m} + \sum_{\beta=1}^m \widehat{z}_n^\beta \frac{\partial}{\partial z_n^\beta} + (q^{-2} - 1) \sum_{b=1}^n \sum_{\beta=1}^m \widehat{z}_b^\beta \frac{\partial}{\partial z_b^\beta} & , \quad a = n \text{ \& } \alpha = m \\ \sum_{\beta=1}^m \widehat{z}_n^\beta \frac{\partial}{\partial z_a^\beta} & , \quad a \neq n \text{ \& } \alpha = m \\ \sum_{b=1}^n \widehat{z}_b^m \frac{\partial}{\partial z_b^\alpha} & , \quad a = n \text{ \& } \alpha \neq m \\ 0 & , \quad \text{otherwise} \end{cases}, \quad (2.18)$$

and with $k \neq n$

$$K_k \frac{\partial}{\partial z_a^\alpha} = \begin{cases} q^{-1} \frac{\partial}{\partial z_a^\alpha} & , \quad k < n \ \& \ a = k \quad \text{or} \quad k > n \ \& \ \alpha = N - k \\ q \frac{\partial}{\partial z_a^\alpha} & , \quad k < n \ \& \ a = k + 1 \quad \text{or} \quad k > n \ \& \ \alpha = N - k + 1 \\ \frac{\partial}{\partial z_a^\alpha} & , \quad \text{otherwise} \end{cases} \quad (2.19)$$

$$F_k \frac{\partial}{\partial z_a^\alpha} = -q^{3/2} \cdot \begin{cases} \frac{\partial}{\partial z_{a-1}^\alpha} & , \quad k < n \ \& \ a = k + 1 \\ \frac{\partial}{\partial z_a^{\alpha-1}} & , \quad k > n \ \& \ \alpha = N - k + 1 \\ 0 & , \quad \text{otherwise} \end{cases} \quad (2.20)$$

$$E_k \frac{\partial}{\partial z_a^\alpha} = -q^{-3/2} \cdot \begin{cases} \frac{\partial}{\partial z_{a+1}^\alpha} & , \quad k < n \ \& \ a = k \\ \frac{\partial}{\partial z_a^{\alpha+1}} & , \quad k > n \ \& \ \alpha = N - k \\ 0 & , \quad \text{otherwise} \end{cases} \quad (2.21)$$

3 Sketch of the proof

Let us outline an idea of the proof of the main theorem. To prove the statement i) of the theorem we have to explain why for arbitrary $\xi \in U_q \mathfrak{g}$, $T \in D(\mathfrak{p}^-)_q$

$$\xi(T) \in D(\mathfrak{p}^-)_q. \quad (3.1)$$

The map $z_a^\alpha \mapsto \widehat{z}_a^\alpha$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$, is extendable up to an embedding of algebras $\mathbb{C}[\mathfrak{p}^-]_q \hookrightarrow \text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$. This embedding intertwines the actions of $U_q \mathfrak{g}$ in $\mathbb{C}[\mathfrak{p}^-]_q$ and $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$ (this is a corollary of the fact that $\mathbb{C}[\mathfrak{p}^-]_q$ is a $U_q \mathfrak{g}$ -module algebra). This observation proves (3.1) for T of the form $\widehat{f}$, $f \in \mathbb{C}[\mathfrak{p}^-]_q$, as well as the first part of the statement ii) of the theorem. What remains is to prove (3.1) for $T = \frac{\partial}{\partial z_a^\alpha}$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$.

The space $\text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$ can be made into a left $\mathbb{C}[\mathfrak{p}^-]_q$ -module as follows:

$$z_a^\alpha(T) = \widehat{z}_a^\alpha \cdot T,$$

with $a = 1, \dots, n$, $\alpha = 1, \dots, m$, $T \in \text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$. This structure is compatible with the action of $U_q \mathfrak{g}$. Consider the $U_q \mathfrak{g}$ -module $\Lambda^1(\mathfrak{p}^-)_q \otimes_{\mathbb{C}[\mathfrak{p}^-]_q} \text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$. The differential $d : \mathbb{C}[\mathfrak{p}^-]_q \rightarrow \Lambda^1(\mathfrak{p}^-)_q$ is a morphism of the $U_q \mathfrak{g}$ -modules. This implies $U_q \mathfrak{g}$ -invariance of the element $\sum_{a=1}^n \sum_{\alpha=1}^m dz_a^\alpha \otimes \frac{\partial}{\partial z_a^\alpha} \in \Lambda^1(\mathfrak{p}^-)_q \otimes_{\mathbb{C}[\mathfrak{p}^-]_q} \text{End}_{\mathbb{C}(q^{1/2})}(\mathbb{C}[\mathfrak{p}^-]_q)$, i.e. for all $\xi \in U_q \mathfrak{g}$

$$\sum_{a=1}^n \sum_{\alpha=1}^m \sum_j \xi'_j dz_a^\alpha \otimes \xi''_j \frac{\partial}{\partial z_a^\alpha} = \varepsilon(\xi) \sum_{a=1}^n \sum_{\alpha=1}^m dz_a^\alpha \otimes \frac{\partial}{\partial z_a^\alpha} \quad (3.2)$$

with ε being the counit of $U_q \mathfrak{g}$, $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$ (Δ is the coproduct in $U_q \mathfrak{g}$). As it was proved in [8], $\Lambda^1(\mathfrak{p}^-)_q$ is a free right $\mathbb{C}[\mathfrak{p}^-]_q$ -module with the generators dz_a^α , $a = 1, \dots, n$,

$\alpha = 1, \dots, m$. Thus, for $\xi \in U_q \mathfrak{g}$ there exists a unique set $f_{\beta,a}^{b,\alpha}(\xi) \in \mathbb{C}[\mathfrak{p}^-]_q$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$, $b = 1, \dots, n$, $\beta = 1, \dots, m$, such that

$$\xi dz_a^\alpha = \sum_{b=1}^n \sum_{\beta=1}^m dz_b^\beta f_{\beta,a}^{b,\alpha}(\xi).$$

Using the later equality, we can rewrite (3.2) as follows:

$$\sum_{a,b=1}^n \sum_{\alpha,\beta=1}^m \sum_j dz_b^\beta \otimes f_{\beta,a}^{b,\alpha}(\xi'_j) \xi_j'' \frac{\partial}{\partial z_a^\alpha} = \varepsilon(\xi) \sum_{a=1}^n \sum_{\alpha=1}^m dz_a^\alpha \otimes \frac{\partial}{\partial z_a^\alpha}. \quad (3.3)$$

Now one can obtain formulae (2.16) - (2.21) (and thus prove (3.1) for $T = \frac{\partial}{\partial z_a^\alpha}$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$) via applying (3.3) to the generators E_i, F_i, K_i, K_i^{-1} of $U_q \mathfrak{g}$.

4 A generalization: q-differential operators in holomorphic q-bundles.

Let Γ be a finitely generated free right $\mathbb{C}[\mathfrak{p}^-]_q$ -module, i.e. there exists an isomorphism of the right $\mathbb{C}[\mathfrak{p}^-]_q$ -modules

$$\pi : \Gamma \rightarrow V \bigotimes \mathbb{C}[\mathfrak{p}^-]_q,$$

with V being a finite dimensional vector space. Elements of Γ are q-analogues of sections of a holomorphic vector bundle over $\mathfrak{p}^-$. *Finitely generated free right $\mathbb{C}[\mathfrak{p}^-]_q$ -modules will be called vector q-bundles.* The isomorphism π will be called a trivialization of Γ .

Let Γ_1, Γ_2 be two vector q-bundles, $\pi_1 : \Gamma_1 \rightarrow V_1 \bigotimes \mathbb{C}[\mathfrak{p}^-]_q$, $\pi_2 : \Gamma_2 \rightarrow V_2 \bigotimes \mathbb{C}[\mathfrak{p}^-]_q$ their trivializations. Set

$$D(\Gamma_1, \Gamma_2)_q = \left\{ D \in \text{Hom}(\Gamma_1, \Gamma_2) \mid \pi_2 \cdot D \cdot \pi_1^{-1} \in \text{Hom}(V_1, V_2) \bigotimes D(\mathfrak{p}^-)_q \right\}.$$

Elements of $D(\Gamma_1, \Gamma_2)_q$ can be treated as q-analogues of differential operators in sections of holomorphic vector bundles.

We need to verify that $D(\Gamma_1, \Gamma_2)_q$ is independent of the choice of trivializations. This follows from the observation that for two trivializations $\pi_1 : \Gamma \rightarrow V_1 \bigotimes \mathbb{C}[\mathfrak{p}^-]_q$, $\pi_2 : \Gamma \rightarrow V_2 \bigotimes \mathbb{C}[\mathfrak{p}^-]_q$ of a q-bundle Γ

$$\pi_2 \cdot (\pi_1)^{-1} \in \text{Hom}(V_1, V_2) \bigotimes \widehat{\mathbb{C}[\mathfrak{p}^-]_q}$$

with $\widehat{\mathbb{C}[\mathfrak{p}^-]_q}$ being the unital subalgebra in $D(\mathfrak{p}^-)_q$ generated by $\widehat{z}_a^\alpha$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$.

Suppose that Γ is a $U_q \mathfrak{g}$ -module q-bundle. It means that Γ is a vector q-bundle and a $U_q \mathfrak{g}$ -module, and the multiplication map $\Gamma \bigotimes \mathbb{C}[\mathfrak{p}^-]_q \rightarrow \Gamma$ is a morphism of the $U_q \mathfrak{g}$ -modules.

For $U_q \mathfrak{g}$ -module vector q-bundles a result analogous to the main theorem (Section 2) can be obtained. Let us formulate it.

If V_1, V_2 are modules over a Hopf algebra A then the space $\text{Hom}(V_1, V_2)$ admits the following "canonical" structure of an A -module: for $\xi \in A$, $T \in \text{Hom}(V_1, V_2)$

$$\xi(T) = \sum_j \xi_j' \cdot T \cdot S(\xi_j''), \quad (4.1)$$

where $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$ (Δ is the coproduct), S is the antipode, and the product in the right-hand side means the composition of the maps $S(\xi''_j) \in \text{End}(V_1)$, $T \in \text{Hom}(V_1, V_2)$, $\xi'_j \in \text{End}(V_2)$. It is well known that this action makes $\text{Hom}(V_1, V_2)$ into an A -module left $\text{End}(V_2)$ -module and an A -module right $\text{End}(V_1)$ -module, i.e. the composition map

$$\text{End}(V_2) \otimes \text{Hom}(V_1, V_2) \otimes \text{End}(V_1) \rightarrow \text{Hom}(V_1, V_2)$$

is a morphism of the A -modules.

Let Γ_1, Γ_2 be $U_q\mathfrak{g}$ -module vector q -bundles. Using our main theorem, one can prove the following

Proposition 4.1 *The subspace $D(\Gamma_1, \Gamma_2)_q \subset \text{Hom}(\Gamma_1, \Gamma_2)$ is $U_q\mathfrak{g}$ -invariant; thus, the composition map*

$$D(\Gamma_2)_q \otimes D(\Gamma_1, \Gamma_2)_q \otimes D(\Gamma_1)_q \rightarrow D(\Gamma_1, \Gamma_2)_q$$

(here $D(\Gamma)_q$ denotes $D(\Gamma, \Gamma)_q$) makes $D(\Gamma_1, \Gamma_2)_q$ into a $U_q\mathfrak{g}$ -module left $D(\Gamma_2)_q$ -module and a $U_q\mathfrak{g}$ -module right $D(\Gamma_1)_q$ -module.

5 Concluding notes

The space of $m \times n$ matrices considered in the present paper is the simplest example of a prehomogeneous vector space of commutative parabolic type [7]. Such vector spaces are closely related to non-compact Hermitian symmetric spaces. Specifically, any non-compact Hermitian symmetric space can be realized (via the Harish-Chandra embedding) as a bounded symmetric domain in some prehomogeneous vector space of commutative parabolic type.

In [10] a q -analogue of an arbitrary prehomogeneous vector space of commutative parabolic type was constructed. More precisely, let U be a bounded symmetric domain, $\mathfrak{p}^-$ the corresponding prehomogeneous vector space, and $\mathfrak{g}$ the complexification of the Lie algebra of the automorphism group of U . In the paper [10] a $U_q\mathfrak{g}$ -module algebra $\mathbb{C}[\mathfrak{p}^-]_q$ and a covariant first order differential calculus $(\Lambda^1(\mathfrak{p}^-), d)$ over $\mathbb{C}[\mathfrak{p}^-]_q$ were introduced (the notation $\mathfrak{g}_{-1}$ was used in [10] instead of $\mathfrak{p}^-$). Using the first order differential calculus, one can define an algebra $D(\mathfrak{p}^-)_q$ of q -differential operators in $\mathbb{C}[\mathfrak{p}^-]_q$ just as it was done in Section 2 in the case of the matrix space.

In this general setting it can also be proved that $D(\mathfrak{p}^-)_q$ is a $U_q\mathfrak{g}$ -module subalgebra in the $U_q\mathfrak{g}$ -module algebra $\text{End}(\mathbb{C}[\mathfrak{p}^-]_q)$. Indeed, it is easy to see that the proof of our main theorem (Section 3) does not use a specific nature of the case when $\mathfrak{p}^-$ is the matrix space.

6 Appendix: Constant coefficient q -differential operators.

Let $D(\mathfrak{p}^-)_q^{\text{const}}$ be the unital subalgebra in $D(\mathfrak{p}^-)_q$ generated by $\frac{\partial}{\partial z_a^\alpha}$, $a = 1, \dots, n$, $\alpha = 1, \dots, m$. By (2.16), (2.19), (2.20), (2.21), it is a $U_q\mathfrak{k}$ -module subalgebra in $D(\mathfrak{p}^-)_q$. Its elements are q -analogues of the constant coefficient differential operators in the space of holomorphic polynomials. It turns out that the natural action of $D(\mathfrak{p}^-)_q^{\text{const}}$ in $\mathbb{C}[\mathfrak{p}^-]_q$ is related to the action of the quantized universal enveloping algebra. Turn to precise formulations.

Denote by $\check{U}_q\mathfrak{g}$ the Hopf algebra derived from $U_q\mathfrak{g}$ by adjoining the pairwise commuting generators $L_i^{\pm 1}$, $i = 1, \dots, N-1$, such that $K_i = \prod_{j=1}^{N-1} L_j^{a_{ij}}$ (with (a_{ij}) being the Cartan matrix for $\mathfrak{g}$), and

$$L_i L_i^{-1} = L_i^{-1} L_i = 1, \quad L_i E_j = q^{\delta_{ij}} E_j L_i, \quad L_i F_j = q^{-\delta_{ij}} F_j L_i,$$

$$\Delta(L_i) = L_i \otimes L_i, \quad S(L_i) = L_i^{-1}, \quad \varepsilon(L_i) = 1.$$

Endow $\check{U}_q\mathfrak{g}$ with a structure of $U_q\mathfrak{g}$ -module algebra via the usual quantum adjoint action: $\text{ad}\xi(\eta) = \sum_j \xi'_j \cdot \eta \cdot S(\xi''_j)$, with $\Delta(\xi) = \sum_j \xi'_j \otimes \xi''_j$.

As it was proved in [5], the $U_q\mathfrak{g}$ -submodule $\text{ad}U_q\mathfrak{g}(L_1^{-2}) \subset \check{U}_q\mathfrak{g}$ is finite-dimensional. Let F be its primitive vector $\text{ad}F_{N-1}\text{ad}F_{N-2}\dots\text{ad}F_1(L_1^{-2})$. Evidently, $\text{ad}U_q\mathfrak{k}(F)$ is a finite dimensional $U_q\mathfrak{k}$ -submodule in $\check{U}_q\mathfrak{g}$. Let $U_q\mathfrak{p}^-$ be the minimal subalgebra in $\check{U}_q\mathfrak{g}$ containing $\text{ad}U_q\mathfrak{k}(F)$. The following statement can be proved.

Proposition 6.1 *For any $\xi \in U_q\mathfrak{p}^-$ there exists a unique $\partial_\xi \in D(\mathfrak{p}^-)_q^{\text{const}}$ such that*

$$\xi(f) = \partial_\xi(f),$$

for any $f \in \mathbb{C}[\mathfrak{p}^-]_q$. The map $\phi : U_q\mathfrak{p}^- \rightarrow D(\mathfrak{p}^-)_q^{\text{const}}$, $\phi : \xi \mapsto \partial_\xi$, is an isomorphism of the $U_q\mathfrak{k}$ -module algebras.

Let us produce a notion of constant coefficient differential operators for vector q -bundles. First of all, we need to distinguish a class of vector q -bundles for which this notion is well defined.

Let $U_q(\mathfrak{p}^- + \mathfrak{k})$ be the Hopf subalgebra in $U_q\mathfrak{g}$ generated by F_n and $U_q\mathfrak{k}$. Suppose that Γ is a $U_q(\mathfrak{p}^- + \mathfrak{k})$ -module vector q -bundle (it means that Γ is a vector q -bundle and a $U_q(\mathfrak{p}^- + \mathfrak{k})$ -module, and the multiplication map $\Gamma \otimes \mathbb{C}[\mathfrak{p}^-]_q \rightarrow \Gamma$ is a morphism of the $U_q(\mathfrak{p}^- + \mathfrak{k})$ -modules). A trivialization $\pi : \Gamma \rightarrow V \otimes \mathbb{C}[\mathfrak{p}^-]_q$ is called good trivialization if it satisfies the following conditions:

i) V is a finite dimensional $U_q(\mathfrak{p}^- + \mathfrak{k})$ -module with the property $F_n v = 0$ for any $v \in V$;

ii) π is a morphism of the $U_q(\mathfrak{p}^- + \mathfrak{k})$ -modules (here $V \otimes \mathbb{C}[\mathfrak{p}^-]_q$ is endowed with $U_q(\mathfrak{p}^- + \mathfrak{k})$ -module structure via the coproduct $\Delta : U_q(\mathfrak{p}^- + \mathfrak{k}) \rightarrow U_q(\mathfrak{p}^- + \mathfrak{k}) \otimes U_q(\mathfrak{p}^- + \mathfrak{k})$).

It can be proved that for any two good trivializations $\pi_1 : \Gamma \rightarrow V_1 \otimes \mathbb{C}[\mathfrak{p}^-]_q$, $\pi_2 : \Gamma \rightarrow V_2 \otimes \mathbb{C}[\mathfrak{p}^-]_q$ of a vector q -bundle Γ

$$\pi_2 \cdot \pi_1^{-1} = T \otimes 1 \tag{6.1}$$

with $T \in \text{Hom}_{U_q(\mathfrak{p}^- + \mathfrak{k})}(V_1, V_2)$.

The set of $U_q(\mathfrak{p}^- + \mathfrak{k})$ -module vector q -bundles admitting good trivializations is the class of vector q -bundles for which the notion of a q -differential operator with constant coefficients is well-defined: if Γ_1, Γ_2 admit good trivializations $\pi_1 : \Gamma_1 \rightarrow V_1 \otimes \mathbb{C}[\mathfrak{p}^-]_q$, $\pi_2 : \Gamma_2 \rightarrow V_2 \otimes \mathbb{C}[\mathfrak{p}^-]_q$, then the elements of the space

$$\left\{ D \in D(\Gamma_1, \Gamma_2)_q \mid \pi_2 \cdot D \cdot \pi_1^{-1} \in \text{Hom}(V_1, V_2) \otimes D(\mathfrak{p}^-)_q^{\text{const}} \right\}$$

can be treated as q -analogues of the constant coefficient differential operators in sections of holomorphic vector bundles. By (6.1), this space is independent of good trivializations.

References

- [1] V. Chari, A. Pressley, *A Guide to Quantum Groups*, Cambridge Univ. Press, 1995.
- [2] E. E. Demidov, *Modules over quantum Weyl algebras*, Vestnik MGU, Mathematics and Mechanics, **1** (1993), 53 – 56.
- [3] H. P. Jakobsen, *Quantized Hermitian Symmetric Spaces*, In "Lie theory and its applications in physics" (Clausthal, 1995), 105 – 116.
- [4] H. P. Jakobsen, *Q-Differential Operators*, math.QA/9907009, 1999.
- [5] A. Joseph, G. Letzter, *Local finiteness of the adjoint action for quantized enveloping algebras*, J. Algebra, **153** (1992), 289 – 318.
- [6] A. Kamita, Y. Morita, and T. Tanisaki, *Quantum deformations of certain prehomogeneous spaces I*, Hiroshima Math. J., **28** (1998), 527 – 540.
- [7] H. Rubenthaler, *Les paires duales dans les algèbres de Lie réductives*, Asterisque, **219**, 1994.
- [8] D. Shklyarov, S. Sinel'shchikov, and L. Vaksman, *Quantum matrix ball: differential and integral calculi*, math.QA/9905035, 1999.
- [9] S. Sinel'shchikov and L. Vaksman, *Hidden symmetry of the differential calculus on the quantum matrix space*, J. Phys. A. **30** (1997), 23 – 26 (see also this volume).
- [10] S. Sinel'shchikov and L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, Math. Phys., Anal., and Geom., **1** (1998), 75 – 100; q-alg/9703005, 1997.